



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁷ : C07D 209/40, 487/06, 513/06, 405/04, 401/04, 405/14, 409/04, A61K 31/40, 31/415, 31/425, 31/41, 31/47, A61P 35/00 // (C07D 487/06, 209:00, 235:00) (C07D 487/06, 209:00, 253:00) (C07D 513/06, 209:00, 277:00) (C07D 487/06, 209:00, 221:00)</p>	A1	<p>(11) International Publication Number: WO 00/56710</p> <p>(43) International Publication Date: 28 September 2000 (28.09.00)</p>
<p>(21) International Application Number: PCT/US00/05057</p> <p>(22) International Filing Date: 28 February 2000 (28.02.00)</p> <p>(30) Priority Data: 9904933.0 4 March 1999 (04.03.99) GB</p> <p>(71) Applicant (for all designated States except US): GLAXO GROUP LIMITED [GB/GB]; Glaxo Wellcome House, Berkeley Avenue, Greenford, Middlesex UB6 0NN (GB).</p> <p>(72) Inventors; and</p> <p>(75) Inventors/Applicants (for US only): GLENNON, Kimberley, Caroline [US/US]; Glaxo Wellcome Inc., Five Moore Drive, P.O. Box 13398, Research Triangle Park, NC 27709 (US). KUYPER, Lee, Frederick [US/US]; Glaxo Wellcome Inc., Five Moore Drive, P.O. Box 13398, Research Triangle Park, NC 27709 (US). LACKEY, Karen, Elizabeth [US/US]; Glaxo Wellcome Inc., Five Moore Drive, P.O. Box 13398, Research Triangle Park, NC 27709 (US). MCNUTT, Robert,</p>		<p>Walton, Jr. [US/US]; 301 Dunhill Drive, Durham, NC 27713 (US).</p> <p>(74) Agents: LEVY, David, J.; Glaxo Wellcome Inc., Five Moore Drive, P.O. Box 13398, Research Triangle Park, NC 27709-3398 (US) et al.</p> <p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</p>
<p>(54) Title: 3-(ANILINOMETHYLENE) OXINDOLES AS PROTEIN TYROSINE KINASE AND PROTEIN SERINE/THREONINE KINASE INHIBITORS</p> <div data-bbox="552 1260 1136 1554"> <p style="text-align: right;">(I)</p> </div> <p>(57) Abstract</p> <p>Compounds of formula (I) wherein R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, A, D, X, Y and Z have the meaning as defined in the claims exhibit protein tyrosine kinase and protein serin/threonine kinase inhibitory activity.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

3-(ANILINOMETHYLENE) OXINDOLES AS PROTEIN TYROSINE KINASE AND PROTEIN SERINE/THREONINE KINASE INHIBITORS

The present invention relates generally to novel amine substituted oxindole compounds and compositions having utility as pharmacological agents in treating diseases or conditions alleviated by the inhibition or
5 antagonism of protein kinase activated signalling pathways in general, and in particular in the pathological processes which involve aberrant cellular proliferation, such as tumor growth, restenosis, atherosclerosis, and thrombosis and methods for using and manufacturing such compounds. In particular, the present invention relates to a series of substituted oxindole
10 compounds, which exhibit protein tyrosine kinase and protein serine/threonine kinase inhibition, and which are useful in inhibiting tumor growth via inhibition of tumor-related angiogenesis.

BACKGROUND OF THE INVENTION

15 Protein kinases play a critical role in the control of cell growth and differentiation and are key mediators of cellular signals leading to the production of growth factors and cytokines. See, for example, Schlessinger and Ullrich, *Neuron* **1992**, 9, 383. A partial non-limiting list of such kinases includes abl, ARaf, ATK, ATM, bcr-abl, Blk, BRaf, Brk, Btk, CDK1, CDK2,
20 CDK3, CDK4, CDK5, CDK6, CDK7, CDK8, CDK9, cfms, c-fms, c-kit, c-met, cRaf1, CSF1R, CSK, c-src, EGFR, ErbB2, ErbB3, ErbB4, ERK, ERK1, ERK2, Fak, fes, FGFR1, FGFR2, FGFR3, FGFR4, FGFR5, Fgr, FLK-4, Fps, Frk, Fyn, GSK, gsk3a, gsk3b, Hck, IGF-1R, IKK, IKK1, IKK2, IKK3, INS-R, Integrin-linkedkinase, Jak, JAK1, JAK2, JAK3, JNK, JNK, Lck, Lyn, MEK,
25 MEK1, MEK2, p38, PDGFR, PIK, PKB1, PKB2, PKB3, PKC, PKC α , PKC β , PKC δ , PKC ϵ , PKC γ , PKC λ , PKC μ , PKC ζ , PLK1, Polo-like kinase, PYK2, tie1, tie2, TrkA, TrkB, TrkC, UL13, UL97, VEGF-R1, VEGF-R2, Yes and Zap70. Protein kinases have been implicated as targets in central nervous system disorders such as Alzheimer's (Mandelkow, E. M. et al. *FEBS Lett.*
30 **1992**, 314, 315. Sengupta, A. et al. *Mol. Cell. Biochem.* **1997**, 167,99), pain


sensation (Yashpal, K. *J. Neurosci.* **1995**, *15*, 3263-72), inflammatory disorders such as arthritis (Badger, *J. Pharm. Exp. Ther.* **1996**, *279*, 1453), psoriasis (Dvir, et al, *J. Cell Biol.* **1991**, *113*, 857), and chronic obstructive pulmononary disease, bone diseases such as osteoporosis (Tanaka et al, *Nature*, **1996**, *383*, 528), cancer (Hunter and Pines, *Cell* **1994**, *79*, 573), atherosclerosis (Hajjar and Pomerantz, *FASEB J.* **1992**, *6*, 2933), thrombosis (Salari, *FEBS* **1990**, *263*, 104), metabolic disorders such as diabetes (Borthwick, A.C. et al. *Biochem. Biophys. Res. Commun.* **1995**, *210*, 738), blood vessel proliferative disorders such as angiogenesis (Strawn et al *Cancer Res.* **1996**, *56*, 3540; Jackson et al *J. Pharm. Exp. Ther.* **1998**, *284*, 687), restenosis (Buchdunger et al, *Proc. Nat. Acad. Sci USA* **1991**, *92*, 2258), autoimmune diseases and transplant rejection (Bolen and Brugge, *Ann. Rev. Immunol.* **1997**, *15*, 371) and infectious diseases such as viral (Littler, E. *Nature* **1992**, *358*, 160), and fungal infections (Lum, R. T. PCT Int. Appl., WO 9805335 A1 980212).

The VEGF-R2 kinase is a receptor tyrosine kinase found in endothelial cells and is involved in angiogenesis--the growth and proliferation of blood vessels from existing capillaries. Angiogenesis plays an important role in development, homeostasis, wound healing, the female reproductive cycle, and in pathological conditions such as rheumatoid arthritis, diabetic retinopathy, macular degeneration, psoriasis and cancer. VEGF-R2 kinase transmits the signal initiated by binding of Vascular Endothelial Growth Factor (VEGF) to the extracellular receptor. Signal transmission to the cell interior is accomplished via tyrosine phosphorylation by VEGF-R2, which prompts proliferation of endothelial cells and the release of cytokines and other cellular processes that result in the growth of new blood vessels. Angiogenesis is critical to the growth of cancerous tumors. Solid tumors will not grow beyond 1-2 mm in size without the support of additional vascularization. Most tumor types, if not all, secrete VEGF in order to stimulate angiogenesis. Inhibition of VEGF-R2 kinase would

therefore interrupt a critical process involved in tumor growth and metastasis as well as other pathologic angiogenic conditions.

SUMMARY OF THE INVENTION

- 5 In brief summary, the invention comprises compounds of the formula (I):



wherein

- 10 Y, Z, A, and D are independently selected from the group consisting of: carbon and nitrogen, with the provisos that: (1) Z and D may be nitrogen, but otherwise no more than one of Y, Z, A, and D may be nitrogen, and (2) when Y, Z, or A are nitrogen, substituent R¹, R², or R³ designated for the respective nitrogen atom is non-existent;
- 15 X is selected from the group consisting of: N, CH, CCF₃, and C(C₁₋₁₂ aliphatic);
- R¹ is selected from the group consisting of: hydrogen, C₁₋₁₂ aliphatic, thiol, hydroxy, hydroxy-C₁₋₁₂ aliphatic, Aryl, Aryl-C₁₋₁₂ aliphatic, R⁹-Aryl-C₁₋₁₂ aliphatic, Cyc, Cyc-C₁₋₆ aliphatic, Het, Het-C₁₋₁₂ aliphatic, C₁₋₁₂ alkoxy,
- 20 Aryloxy, amino, C₁₋₁₂ aliphatic amino, di-C₁₋₁₂ aliphatic amino, di-C₁₋₁₂ aliphatic aminocarbonyl, di-C₁₋₁₂ aliphatic aminosulfonyl, C₁₋₁₂ alkoxy carbonyl, fluoro, bromo, iodo, cyano, sulfonamide, or nitro, where R⁹, Aryl, Cyc and Het are as defined below;

- R^2 is selected from the group consisting of: hydrogen, C_{1-12} aliphatic, N-hydroxyimino- C_{1-12} aliphatic, C_{1-12} alkoxy, hydroxy- C_{1-12} aliphatic, C_{1-12} alkoxy-carbonyl, carboxyl C_{1-12} aliphatic, Aryl, R^9 -Aryl-oxycarbonyl, R^9 -oxycarbonyl-Aryl, Het, aminocarbonyl, C_{1-12} aliphatic-aminocarbonyl, Aryl-
- 5 C_{1-12} aliphatic-aminocarbonyl, R^9 -Aryl- C_{1-12} aliphatic-aminocarbonyl, Het- C_{1-12} aliphatic-aminocarbonyl, hydroxy- C_{1-12} aliphatic-aminocarbonyl, C_{1-12} -alkoxy- C_{1-12} aliphatic-aminocarbonyl, C_{1-12} alkoxy- C_{1-12} aliphatic-amino, di- C_{1-12} aliphatic amino, di- C_{1-12} aliphatic aminocarbonyl, di- C_{1-12} aliphatic aminosulfonyl, halogen, hydroxy, C_{1-12} aliphatic-sulfonyl, aminosulfonyl, and
- 10 C_{1-12} aliphatic-aminosulfonyl, where R^9 , Aryl and Het are as defined below, with the proviso that where X is nitrogen, R^2 is not chloro or 3,6-dihydro-6-methyl-2-oxo-2H-1,3,4-thiadiazin-5-yl;
- R^1 and R^2 are optionally joined to form a fused ring selected from the group as defined for Het below, and said fused ring is optionally substituted by a
- 15 substituent selected from the group consisting of: C_{1-12} aliphatic, halogen, nitro, cyano, C_{1-12} alkoxy, amino, hydroxyl, (R^{10} , R^{11})-amino, and oxo;
- R^3 is selected from the group consisting of: hydrogen, C_{1-12} aliphatic, hydroxy, hydroxy C_{1-12} aliphatic, di- C_{1-12} aliphatic amino, di- C_{1-12} aliphatic aminocarbonyl, di- C_{1-12} aliphatic aminosulfonyl, C_{1-12} alkoxy, Aryl, Aryloxy,
- 20 hydroxy-Aryl, Het, hydroxy-Het, Het-oxy, or halogen, where Aryl and Het are as defined below, with the proviso that where X is nitrogen R^3 is not fluoro;
- R^2 and R^3 are optionally joined to form a fused ring selected from the group as defined for Het below, and said fused ring is optionally substituted by C_{1-6} aliphatic or C_{1-6} aliphatic-carbonyl;
- 25 with the proviso that R^1 , R^2 , and R^3 cannot simultaneously be hydrogen;
- R^4 , R^5 and R^6 may be the same or different and are independently selected from the group consisting of: hydrogen, C_{1-12} aliphatic, thiol, C_{1-6} aliphatic-thio, di(trifluoromethyl)hydroxymethyl, carboxamide, mono- C_{1-12} aliphatic aminocarbonyl, hydroxy, hydroxy- C_{1-12} aliphatic, Aryl, Aryl- C_{1-12} aliphatic, R^9 -
- 30 Aryl- C_{1-12} aliphatic, Cyc, Cyc- C_{1-6} aliphatic, Het, Het- C_{1-12} aliphatic, C_{1-12}

- alkoxy, Aryloxy, Het-oxy, amino, (R¹⁰,R¹¹)-amino-C₁₋₁₂ aliphatic aminocarbonyl, (R¹⁰,R¹¹)-amino-C₁₋₁₂ aliphatic alkoxycarbonyl, (R¹⁰,R¹¹)-amino-C₁₋₁₂ aliphatic aminocarbonylamino, (R¹⁰,R¹¹)-amino-C₁₋₆ aliphatic alkoxycarbonylamino, (R¹⁰,R¹¹)-amino-C₁₋₆ aliphatic sulfonyl, Het-C₁₋₆
- 5 aliphatic aminocarbonyl, Het-C₁₋₆ aliphatic aminocarbonylamino, Het-C₁₋₆ alkoxycarbonylamino, Het-C₁₋₆ aliphatic carbonyl, Het-C₁₋₆ alkoxycarbonyl, C₁₋₆ aliphatic sulfonyl-C₁₋₆ aliphatic aminoalkyl, C₁₋₆ aliphatic sulfonyl-C₁₋₆ aliphatic aminoalkyl-Het-, C₁₋₆ alkoxycarbonyl, C₁₋₆ aliphatic carbonylamino, (C₁₋₆ aliphatic carbonyl)(C₁₋₆ aliphatic)amino, (R¹⁰,R¹¹)-amino-C₁₋₆ aliphatic
- 10 carbonylamino, [(R¹⁰,R¹¹)-amino-C₁₋₆ aliphatic carbonyl][C₁₋₆ aliphatic]amino, (R¹⁰,R¹¹)-amino-C₁₋₆ aliphatic sulfonylamino, [(R¹⁰,R¹¹)-amino-C₁₋₆ aliphatic sulfonyl][C₁₋₆ aliphatic]amino, halogen, cyano, diethoxyphosphorylmethyl, nitro, trifluoromethyl, and trifluoromethoxy, where R⁹, R¹⁰, R¹¹, Aryl, Cyc and Het are as defined below, with the proviso that
- 15 where X is nitrogen, R⁴, R⁵ and R⁶ is not nitro;
 R⁷ and R⁸ may be the same or different and are independently selected from the group consisting of: hydrogen, halogen, C₁₋₂ alkoxy, hydroxy, C₁₋₃-aliphatic, and C₁₋₃ aliphatic;
 with the proviso that R⁴, R⁵, R⁶, R⁷, and R⁸ cannot simultaneously be
- 20 hydrogen;
 wherein R⁷ may additionally be optionally fused to R⁵ so as to form a fused benzo ring from the R⁵ to the R⁷ positions;
 R⁹ is selected from the group consisting of: C₁₋₁₂ aliphatic, hydroxy, C₁₋₁₂ alkoxy and halogen;
- 25 R¹⁰ and R¹¹ may be the same or different and are independently selected from the group consisting of: hydrogen, C₁₋₆ aliphatic and Het;
 Aryl is selected from the group consisting of: phenyl, naphthyl, phenanthryl and anthracenyl;


Cyc is selected from the group consisting of: cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl, and optionally has one or more degrees of unsaturation;

Het is a saturated or unsaturated heteroatom ring system selected from the
5 group consisting of: benzimidazole, dihydrothiophene, dioxin, dioxane, dioxolane, dithiane, dithiazine, dithiazole, dithiolane, furan, imidazole, isoquinoline, morpholine, oxazole, oxadiazole, oxathiazole, oxathiazolidine, oxazine, oxadiazine, piperazine, piperidine, pyran, pyrazine, pyrazole, pyridine, pyrimidine, pyrrole, pyrrolidine, quinoline, tetrahydrofuran, tetrazine,
10 thiadiazine, thiadiazole, thiatriazole, thiazine, thiazole, thiomorpholine, thiophene, thiopyran, triazine and triazole, where any of said heterocyclic rings may be optionally substituted by a substituent selected from the group consisting of: C₁₋₁₂ aliphatic, hydroxy, C₁₋₁₂ alkoxy, (R¹⁰,R¹¹)-amino, (R¹⁰,R¹¹)-amino-C₁₋₁₂ aliphatic, (R¹⁰,R¹¹)-amino-C₁₋₁₂ aliphatic amino, oxo
15 and dioxo;

and the pharmaceutically acceptable salts, polymorphs, esters, amides, carbamates, solvates, hydrates, affinity reagents and prodrugs thereof in either crystalline or amorphous form. The esters, amides and carbamates, are preferably hydrolyzable and more preferably are biohydrolyzable.

20

A more preferred genus of compounds of the present invention includes compounds of formula (II), defined as follows:



wherein

- R¹ is selected from the group consisting of: hydrogen, C₁₋₁₂ aliphatic, thiol,
 5 hydroxy, hydroxy-C₁₋₁₂ aliphatic, Aryl, Aryl-C₁₋₁₂ aliphatic, R⁹-Aryl-C₁₋₁₂
 aliphatic, Cyc, Cyc-C₁₋₆ aliphatic, Het, Het-C₁₋₁₂ aliphatic, C₁₋₁₂ alkoxy,
 Aryloxy, amino, C₁₋₁₂ aliphatic amino, di-C₁₋₁₂ aliphatic amino, di-C₁₋₁₂
 aliphatic aminocarbonyl, di-C₁₋₁₂ aliphatic aminosulfonyl, C₁₋₁₂
 alkoxycarbonyl, fluoro, bromo, iodo, cyano, sulfonamide, or nitro, where R⁹,
 10 Aryl, Cyc and Het are as defined below;
 R² is selected from the group consisting of: hydrogen, C₁₋₁₂ aliphatic, N-
 hydroxyimino-C₁₋₁₂ aliphatic, C₁₋₁₂ alkoxy, hydroxy-C₁₋₁₂ aliphatic, C₁₋₁₂
 alkoxycarbonyl, carboxyl C₁₋₁₂ aliphatic, Aryl, R⁹-Aryl-oxycarbonyl, R⁹-
 oxycarbonyl-Aryl, Het, aminocarbonyl, C₁₋₁₂ aliphatic-aminocarbonyl, Aryl-
 15 C₁₋₁₂ aliphatic-aminocarbonyl, R⁹-Aryl-C₁₋₁₂ aliphatic-aminocarbonyl, Het-
 C₁₋₁₂ aliphatic-aminocarbonyl, hydroxy-C₁₋₁₂ aliphatic-aminocarbonyl, C₁₋₁₂-
 alkoxy-C₁₋₁₂ aliphatic-aminocarbonyl, C₁₋₁₂ alkoxy-C₁₋₁₂ aliphatic-amino, di-
 C₁₋₁₂ aliphatic amino, di-C₁₋₁₂ aliphatic aminocarbonyl, di-C₁₋₁₂ aliphatic
 aminosulfonyl, halogen, hydroxy, C₁₋₁₂ aliphatic-sulfonyl, aminosulfonyl, or
 20 one or more substituents selected from the group consisting of: C₁₋₁₂
 aliphatic-aminosulfonyl, where R⁹, Aryl and Het are as defined below;
 R¹ and R² are optionally joined to form a fused ring selected from the group
 as defined for Het below, and said fused ring is optionally substituted by

C₁₋₁₂ aliphatic, halogen, nitro, cyano, C₁₋₁₂ alkoxy, amino, hydroxyl, (R¹⁰, R¹¹)-amino, or oxo;

R³ is selected from the group consisting of: hydrogen, C₁₋₁₂ aliphatic, hydroxy, hydroxy C₁₋₁₂ aliphatic, di-C₁₋₁₂ aliphatic amino, di-C₁₋₁₂ aliphatic
 5 aminocarbonyl, di-C₁₋₁₂ aliphatic aminosulfonyl, C₁₋₁₂ alkoxy, Aryl, Aryloxy, hydroxy-Aryl, Het, hydroxy-Het, Het-oxy, or halogen, where Aryl and Het are as defined below;

R² and R³ are optionally joined to form a fused ring selected from the group as defined for Het below, and said fused ring is optionally substituted by C₁₋₆
 10 aliphatic or C₁₋₆ aliphatic-carbonyl;

with the proviso that R¹, R² and R³ cannot simultaneously be hydrogen;

R⁴, R⁵ and R⁶ may be the same or different and are independently selected from the group consisting of: hydrogen, C₁₋₁₂ aliphatic, thiol, C₁₋₆ aliphatic-thio, di(trifluoromethyl)hydroxymethyl, carboxamide, mono-C₁₋₁₂ aliphatic
 15 aminocarbonyl, hydroxy, hydroxy-C₁₋₁₂ aliphatic, Aryl, Aryl-C₁₋₁₂ aliphatic, R⁹-Aryl-C₁₋₁₂ aliphatic, Cyc, Cyc-C₁₋₆ aliphatic, Het, Het-C₁₋₁₂ aliphatic, C₁₋₁₂ alkoxy, Aryloxy, Het-oxy, amino, (R¹⁰, R¹¹)-amino-C₁₋₁₂ aliphatic aminocarbonyl, (R¹⁰, R¹¹)-amino-C₁₋₁₂ aliphatic alkoxy carbonyl, (R¹⁰, R¹¹)-amino-C₁₋₁₂ aliphatic aminocarbonylamino, (R¹⁰, R¹¹)-amino-C₁₋₆ aliphatic
 20 alkoxy carbonylamino, (R¹⁰, R¹¹)-amino-C₁₋₆ aliphatic sulfonyl, Het-C₁₋₆ aliphatic aminocarbonyl, Het-C₁₋₆ aliphatic aminocarbonylamino, Het-C₁₋₆ alkoxy carbonylamino, Het-C₁₋₆ aliphatic carbonyl, Het-C₁₋₆ alkoxy carbonyl, C₁₋₆ aliphatic sulfonyl-C₁₋₆ aliphatic aminoalkyl, C₁₋₆ aliphatic sulfonyl-C₁₋₆ aliphatic aminoalkyl-Het-, C₁₋₆ alkoxy carbonyl, C₁₋₆ aliphatic carbonylamino,
 25 (C₁₋₆ aliphatic carbonyl)(C₁₋₆ aliphatic)amino, (R¹⁰, R¹¹)-amino-C₁₋₆ aliphatic carbonylamino, [(R¹⁰, R¹¹)-amino-C₁₋₆ aliphatic carbonyl][C₁₋₆ aliphatic]amino, (R¹⁰, R¹¹)-amino-C₁₋₆ aliphatic sulfonylamino, [(R¹⁰, R¹¹)-amino-C₁₋₆ aliphatic sulfonyl][C₁₋₆ aliphatic]amino, halogen, cyano, diethoxyphosphorylmethyl, nitro, trifluoromethyl, or trifluoromethoxy, where
 30 R⁹, R¹⁰, R¹¹, Aryl, Cyc and Het are as defined below;

R^7 and R^8 may be the same or different and are independently selected from the group consisting of: hydrogen, halogen, C_{1-2} alkoxy, hydroxy, C_{1-3} -aliphatic and C_{1-3} aliphatic;

with the proviso that R^4 , R^5 , R^6 , R^7 and R^8 cannot simultaneously be
5 hydrogen;

wherein R^7 may additionally be optionally fused to R^5 so as to form a fused benzo ring from the R^5 to the R^7 positions;

R^9 is selected from the group consisting of: C_{1-12} aliphatic, hydroxy, C_{1-12} alkoxy, or halogen;

10 R^{10} and R^{11} may be the same or different and are independently selected from the group consisting of: hydrogen, C_{1-6} aliphatic and Het;

Aryl is selected from the group consisting of: phenyl, naphthyl, phenanthryl or anthracenyl;

Cyc is selected from the group consisting of: cyclopropyl, cyclobutyl,
15 cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl, and optionally has one or more degrees of unsaturation;

Het is a saturated or unsaturated heteroatom ring system selected from the group consisting of: benzimidazole, dihydrothiophene, dioxin, dioxane, dioxolane, dithiane, dithiazine, dithiazole, dithiolane, furan, imidazole,


20 isoquinoline, morpholine, oxazole, oxadiazole, oxathiazole, oxathiazolidine, oxazine, oxadiazine, piperazine, piperidine, pyran, pyrazine, pyrazole, pyridine, pyrimidine, pyrrole, pyrrolidine, quinoline, tetrahydrofuran, tetrazine, thiadiazine, thiadiazole, thiatriazole, thiazine, thiazole, thiomorpholine, thiophene, thiopyran, triazine and triazole, where any of said heterocyclic

25 rings may be optionally substituted by a substituent selected from the group consisting of: C_{1-12} aliphatic, hydroxy, C_{1-12} alkoxy, (R^{10}, R^{11}) -amino, (R^{10}, R^{11}) -amino- C_{1-12} aliphatic, (R^{10}, R^{11}) -amino- C_{1-12} aliphatic amino, oxo or dioxo;

and the pharmaceutically acceptable salts, esters, amides, carbamates,
30 solvates, hydrates, affinity reagents and prodrugs thereof in either crystalline

or amorphous form. The esters, amides and carbamates, are preferably hydrolyzable and more preferably are biohydrolyzable.

Another preferred genus of compounds of the present invention includes compounds of formula (III), defined as follows:



10 wherein

R¹ is selected from the group consisting of: hydrogen, C₁₋₁₂ aliphatic, thiol, hydroxy, hydroxy-C₁₋₁₂ aliphatic, Aryl, Aryl-C₁₋₁₂ aliphatic, R⁹-Aryl-C₁₋₁₂ aliphatic, Cyc, Cyc-C₁₋₆ aliphatic, Het, Het-C₁₋₁₂ aliphatic, C₁₋₁₂ alkoxy, Aryloxy, amino, C₁₋₁₂ aliphatic amino, di-C₁₋₁₂ aliphatic amino, di-C₁₋₁₂ aliphatic aminocarbonyl, di-C₁₋₁₂ aliphatic aminosulfonyl, C₁₋₁₂ alkoxy carbonyl, fluoro, bromo, iodo, cyano, sulfonamide, or nitro, where R⁹, Aryl, Cyc and Het are as defined below;

R² is selected from the group consisting of: hydrogen, C₁₋₁₂ aliphatic, N-hydroxyimino-C₁₋₁₂ aliphatic, C₁₋₁₂ alkoxy, hydroxy-C₁₋₁₂ aliphatic, C₁₋₁₂ alkoxy carbonyl, carboxyl C₁₋₁₂ aliphatic, Aryl, R⁹-Aryl-oxycarbonyl, R⁹-oxycarbonyl-Aryl, Het, aminocarbonyl, C₁₋₁₂ aliphatic-aminocarbonyl, Aryl-C₁₋₁₂ aliphatic-aminocarbonyl, R⁹-Aryl-C₁₋₁₂ aliphatic-aminocarbonyl, Het-C₁₋₁₂ aliphatic-aminocarbonyl, hydroxy-C₁₋₁₂ aliphatic-aminocarbonyl, C₁₋₁₂-

alkoxy-C₁₋₁₂ aliphatic-aminocarbonyl, C₁₋₁₂ alkoxy-C₁₋₁₂ aliphatic-amino, di-C₁₋₁₂ aliphatic amino, di-C₁₋₁₂ aliphatic aminocarbonyl, di-C₁₋₁₂ aliphatic aminosulfonyl, halogen, hydroxy, C₁₋₁₂ aliphatic-sulfonyl, aminosulfonyl, or one or more substituents selected from the group consisting of: C₁₋₁₂ aliphatic-aminosulfonyl, where R⁹, Aryl and Het are as defined below, with the proviso that R² is not chloro or 3,6-dihydro-6-methyl-2-oxo-2H-1,3,4-thiadiazin-5-yl;

R¹ and R² are optionally joined to form a fused ring selected from the group as defined for Het below, and said fused ring is optionally substituted by C₁₋₁₂ aliphatic, halogen, nitro, cyano, C₁₋₁₂ alkoxy, amino, hydroxyl, (R¹⁰, R¹¹)-amino, or oxo;

R³ is selected from the group consisting of: hydrogen, C₁₋₁₂ aliphatic, hydroxy, hydroxy C₁₋₁₂ aliphatic, di-C₁₋₁₂ aliphatic amino, di-C₁₋₁₂ aliphatic aminocarbonyl, di-C₁₋₁₂ aliphatic aminosulfonyl, C₁₋₁₂ alkoxy, Aryl, Aryloxy, hydroxy-Aryl, Het, hydroxy-Het, Het-oxy, or halogen, where Aryl and Het are as defined below, with the proviso R³ is not fluoro;

R² and R³ are optionally joined to form a fused ring selected from the group as defined for Het below, and said fused ring is optionally substituted by C₁₋₆ aliphatic or C₁₋₆ aliphatic-carbonyl;

with the proviso that R¹, R² and R³ cannot simultaneously be hydrogen;

R⁴, R⁵ and R⁶ may be the same or different and are independently selected from the group consisting of: hydrogen, C₁₋₁₂ aliphatic, thiol, C₁₋₆ aliphatic-thio, di(trifluoromethyl)hydroxymethyl, carboxamide, mono-C₁₋₁₂ aliphatic aminocarbonyl, hydroxy, hydroxy-C₁₋₁₂ aliphatic, Aryl, Aryl-C₁₋₁₂ aliphatic, R⁹-Aryl-C₁₋₁₂ aliphatic, Cyc, Cyc-C₁₋₆ aliphatic, Het, Het-C₁₋₁₂ aliphatic, C₁₋₁₂ alkoxy, Aryloxy, Het-oxy, amino, (R¹⁰, R¹¹)-amino-C₁₋₁₂ aliphatic aminocarbonyl, (R¹⁰, R¹¹)-amino-C₁₋₁₂ aliphatic alkoxycarbonyl, (R¹⁰, R¹¹)-amino-C₁₋₁₂ aliphatic aminocarbonylamino, (R¹⁰, R¹¹)-amino-C₁₋₆ aliphatic alkoxycarbonylamino, (R¹⁰, R¹¹)-amino-C₁₋₆ aliphatic aminosulfonyl, Het-C₁₋₆ aliphatic aminocarbonyl, Het-C₁₋₆ aliphatic aminocarbonylamino, Het-C₁₋₆


- alkoxycarbonylamino, Het-C₁₋₆ aliphatic carbonyl, Het-C₁₋₆ alkoxycarbonyl, C₁₋₆ aliphatic sulfonyl-C₁₋₆ aliphatic aminoalkyl, C₁₋₆ aliphatic sulfonyl-C₁₋₆ aliphatic aminoalkyl-Het-, C₁₋₆ alkoxycarbonyl, C₁₋₆ aliphatic carbonylamino, (C₁₋₆ aliphatic carbonyl)(C₁₋₆ aliphatic)amino, (R¹⁰, R¹¹)-amino-C₁₋₆ aliphatic carbonylamino, [(R¹⁰, R¹¹)-amino-C₁₋₆ aliphatic carbonyl][C₁₋₆ aliphatic]amino, (R¹⁰, R¹¹)-amino-C₁₋₆ aliphatic sulfonylamino, [(R¹⁰, R¹¹)-amino-C₁₋₆ aliphatic sulfonyl][C₁₋₆ aliphatic]amino, halogen, cyano, diethoxyphosphorylmethyl, nitro, trifluoromethyl, or trifluoromethoxy, where R⁹, R¹⁰, R¹¹, Aryl, Cyc and Het are as defined below, with the proviso that
- 10 R⁴, R⁵ and R⁶ is not nitro;
- R⁷ and R⁸ may be the same or different and are independently selected from the group consisting of: hydrogen, halogen, C₁₋₂ alkoxy, hydroxy, C₁₋₃-aliphatic and C₁₋₃ aliphatic;
- with the proviso that R⁴, R⁵, R⁶, R⁷ and R⁸ cannot simultaneously be
- 15 hydrogen;
- R⁹ is selected from the group consisting of: C₁₋₁₂ aliphatic, hydroxy, C₁₋₁₂ alkoxy, or halogen;
- R¹⁰ and R¹¹ may be the same or different and are independently selected from the group consisting of: hydrogen, C₁₋₆ aliphatic and Het;
- 20 Aryl is selected from the group consisting of: phenyl, naphthyl, phenanthryl or anthracenyl;
- Cyc is selected from the group consisting of: cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or cyclooctyl, any one of which may have one or more degrees of unsaturation;
- 25 Het is a saturated or unsaturated heteroatom ring system selected from the group consisting of: benzimidazole, dihydrothiophene, dioxin, dioxane, dioxolane, dithiane, dithiazine, dithiazole, dithiolane, furan, imidazole, isoquinoline, morpholine, oxazole, oxadiazole, oxathiazole, oxathiazolidine, oxazine, oxadiazine, piperazine, piperidine, pyran, pyrazine, pyrazole,
- 30 pyridine, pyrimidine, pyrrole, pyrrolidine, quinoline, tetrahydrofuran, tetrazine,

thiadiazine, thiadiazole, thiatriazole, thiazine, thiazole, thiomorpholine, thiophene, thiopyran, triazine and triazole, where any of said heterocyclic rings may be optionally substituted by a substituent selected from the group consisting of: C₁₋₁₂ aliphatic, hydroxy, C₁₋₁₂ alkoxy, (R¹⁰,R¹¹)-amino, (R¹⁰,R¹¹)-amino-C₁₋₁₂ aliphatic, (R¹⁰,R¹¹)-amino-C₁₋₁₂ aliphatic amino, oxo or dioxo;

and the pharmaceutically acceptable salts, esters, amides, carbamates, solvates, hydrates, affinity reagents and prodrugs thereof in either crystalline or amorphous form. The esters, amides and carbamates, are preferably hydrolyzable and more preferably are biohydrolyzable.

Another preferred genus of compounds of the present invention includes compounds of formula (IV), defined as follows:

15



wherein

Y, Z, A, and D are independently selected from the group consisting of: carbon and nitrogen, with the provisos that: (1) Z and D may be nitrogen, but otherwise no more than one of Y, Z, A, and D may be nitrogen, and (2) when Y, Z, or A are nitrogen, substituent R¹, R², or R³ designated for the respective nitrogen atom is non-existent;

R¹ is selected from the group consisting of: hydrogen, C₁₋₁₂ aliphatic, thiol, hydroxy, hydroxy-C₁₋₁₂ aliphatic, Aryl, Aryl-C₁₋₁₂ aliphatic, R⁹-Aryl-C₁₋₁₂

aliphatic, Cyc, Cyc-C₁₋₆ aliphatic, Het, Het-C₁₋₁₂ aliphatic, C₁₋₁₂ alkoxy, Aryloxy, amino, C₁₋₁₂ aliphatic amino, di-C₁₋₁₂ aliphatic amino, di-C₁₋₁₂ aliphatic aminocarbonyl, di-C₁₋₁₂ aliphatic aminosulfonyl, C₁₋₁₂ alkoxycarbonyl, fluoro, bromo, iodo, cyano, sulfonamide, or nitro, where R⁹,

5 Aryl, Cyc and Het are as defined below;

R² is selected from the group consisting of: hydrogen, C₁₋₁₂ aliphatic, N-hydroxyimino-C₁₋₁₂ aliphatic, C₁₋₁₂ alkoxy, hydroxy-C₁₋₁₂ aliphatic, C₁₋₁₂ alkoxycarbonyl, carboxyl C₁₋₁₂ aliphatic, Aryl, R⁹-Aryl-oxycarbonyl, R⁹-oxycarbonyl-Aryl, Het, aminocarbonyl, C₁₋₁₂ aliphatic-aminocarbonyl, Aryl-

10 C₁₋₁₂ aliphatic-aminocarbonyl, R⁹-Aryl-C₁₋₁₂ aliphatic-aminocarbonyl, Het-C₁₋₁₂ aliphatic-aminocarbonyl, hydroxy-C₁₋₁₂ aliphatic-aminocarbonyl, C₁₋₁₂-alkoxy-C₁₋₁₂ aliphatic-aminocarbonyl, C₁₋₁₂ alkoxy-C₁₋₁₂ aliphatic-amino, di-C₁₋₁₂ aliphatic amino, di-C₁₋₁₂ aliphatic aminocarbonyl, di-C₁₋₁₂ aliphatic aminosulfonyl, halogen, hydroxy, C₁₋₁₂ aliphatic-sulfonyl, aminosulfonyl, or
15 one or more substituents selected from the group consisting of: C₁₋₁₂ aliphatic-aminosulfonyl, where R⁹, Aryl and Het are as defined below;

R¹ and R² are optionally joined to form a fused ring selected from the group as defined for Het below, and said fused ring is optionally substituted by C₁₋₁₂ aliphatic, halogen, nitro, cyano, C₁₋₁₂ alkoxy, amino, hydroxyl, (R¹⁰,

20 R¹¹)-amino, or oxo;

R³ is selected from the group consisting of: hydrogen, C₁₋₁₂ aliphatic, hydroxy, hydroxy C₁₋₁₂ aliphatic, di-C₁₋₁₂ aliphatic amino, di-C₁₋₁₂ aliphatic aminocarbonyl, di-C₁₋₁₂ aliphatic aminosulfonyl, C₁₋₁₂ alkoxy, Aryl, Aryloxy, hydroxy-Aryl, Het, hydroxy-Het, Het-oxy, or halogen, where Aryl and Het are

25 as defined below;

R² and R³ are optionally joined to form a fused ring selected from the group as defined for Het below, and said fused ring is optionally substituted by C₁₋₆ aliphatic or C₁₋₆ aliphatic-carbonyl;

with the proviso that R¹, R² and R³ cannot simultaneously be hydrogen;

- R⁴, R⁵ and R⁶ may be the same or different and are independently selected from the group consisting of: hydrogen, C₁₋₁₂ aliphatic, thiol, C₁₋₆aliphatic-thio, di(trifluoromethyl)hydroxymethyl, carboxamide, mono-C₁₋₁₂aliphatic aminocarbonyl, hydroxy, hydroxy-C₁₋₁₂ aliphatic, Aryl, Aryl-C₁₋₁₂ aliphatic, R⁹-
- 5 Aryl-C₁₋₁₂ aliphatic, Cyc, Cyc-C₁₋₆ aliphatic, Het, Het-C₁₋₁₂ aliphatic, C₁₋₁₂ alkoxy, Aryloxy, Het-oxy, amino, (R¹⁰,R¹¹)-amino-C₁₋₁₂ aliphatic aminocarbonyl, (R¹⁰,R¹¹)-amino-C₁₋₁₂ aliphatic alkoxy carbonyl, (R¹⁰,R¹¹)-amino-C₁₋₁₂ aliphatic aminocarbonylamino, (R¹⁰,R¹¹)-amino-C₁₋₆ aliphatic alkoxy carbonylamino, (R¹⁰,R¹¹)-amino-C₁₋₆ aliphatic sulfonyl, Het-C₁₋₆
- 10 aliphatic aminocarbonyl, Het-C₁₋₆ aliphatic aminocarbonylamino, Het-C₁₋₆ alkoxy carbonylamino, Het-C₁₋₆ aliphatic carbonyl, Het-C₁₋₆ alkoxy carbonyl, C₁₋₆ aliphatic sulfonyl-C₁₋₆ aliphatic aminoalkyl, C₁₋₆ aliphatic sulfonyl-C₁₋₆ aliphatic aminoalkyl-Het-, C₁₋₆ alkoxy carbonyl, C₁₋₆ aliphatic carbonylamino, (C₁₋₆ aliphatic carbonyl)(C₁₋₆ aliphatic)amino, (R¹⁰,R¹¹)-amino-C₁₋₆ aliphatic
- 15 carbonylamino, [(R¹⁰,R¹¹)-amino-C₁₋₆ aliphatic carbonyl][C₁₋₆ aliphatic]amino, (R¹⁰,R¹¹)-amino-C₁₋₆ aliphatic sulfonylamino, [(R¹⁰,R¹¹)-amino-C₁₋₆ aliphatic sulfonyl][C₁₋₆ aliphatic]amino, halogen, cyano, diethoxyphosphorylmethyl, nitro, trifluoromethyl, or trifluoromethoxy, where R⁹, R¹⁰, R¹¹, Aryl, Cyc and Het are as defined below;
- 20 R⁷ and R⁸ may be the same or different and are independently selected from the group consisting of: hydrogen, halogen, C₁₋₂ alkoxy, hydroxy, C₁₋₃-aliphatic and C₁₋₃ aliphatic;
- with the proviso that R⁴, R⁵, R⁶, R⁷ and R⁸ cannot simultaneously be hydrogen;
- 25 wherein R⁷ may additionally be optionally fused to R⁵ so as to form a fused benzo ring from the R⁵ to the R⁷ positions;
- R⁹ is selected from the group consisting of: C₁₋₁₂ aliphatic, hydroxy, C₁₋₁₂ alkoxy, or halogen;
- R¹⁰ and R¹¹ may be the same or different and are independently selected
- 30 from the group consisting of: hydrogen, C₁₋₆ aliphatic and Het;

Aryl is selected from the group consisting of: phenyl, naphthyl, phenanthryl or anthracenyl;

Cyc is selected from the group consisting of: cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or cyclooctyl, and optionally has one or
5 more degrees of unsaturation;

Het is a saturated or unsaturated heteroatom ring system selected from the group consisting of: benzimidazole, dihydrothiophene, dioxin, dioxane, dioxolane, dithiane, dithiazine, dithiazole, dithiolane, furan, imidazole, isoquinoline, morpholine, oxazole, oxadiazole, oxathiazole, oxathiazolidine,
10 oxazine, oxadiazine, piperazine, piperidine, pyran, pyrazine, pyrazole, pyridine, pyrimidine, pyrrole, pyrrolidine, quinoline, tetrahydrofuran, tetrazine, thiadiazine, thiadiazole, thiatriazole, thiazine, thiazole, thiomorpholine, thiophene, thiopyran, triazine and triazole, where any of said heterocyclic rings may be optionally substituted by a substituent selected from the group
15 consisting of: C₁₋₁₂ aliphatic, hydroxy, C₁₋₁₂ alkoxy, (R¹⁰, R¹¹)-amino, (R¹⁰, R¹¹)-amino-C₁₋₁₂ aliphatic, (R¹⁰, R¹¹)-amino-C₁₋₁₂ aliphatic amino, oxo or dioxo;

and the pharmaceutically acceptable salts, esters, amides, carbamates, solvates, hydrates, affinity reagents and prodrugs thereof in either crystalline
20 or amorphous form. The esters, amides and carbamates, are preferably hydrolyzable and more preferably are biohydrolyzable.

While the ensuing discussion refers to the compound of formula (I), it will be understood that the compounds of formula (I) includes the compounds of formulas (II), (III) and (IV); accordingly, references hereafter
25 to formula (I) should hereafter be understood to includes the compounds of formulas (II), (III) and (IV) as well as the compounds of formula (I). Furthermore, references in the ensuing discussion to the formula (I) should also be understood as referring to the compounds of Examples 163-212 of Table 5 below which, while not included within the general formula (I), have
30 been found by the inventors to have Kinase inhibiting properties.

Due to the presence of an oxindole exocyclic double bond, also included in the compounds of the invention are their respective pure E and Z geometric isomers as well as mixtures of E and Z isomers. The invention as described and claimed does not set any limiting ratios on prevalence of Z to
5 E isomers.

Likewise, it is understood that compounds of formula (I) as used herein includes all tautomeric forms other than the specific tautomer represented by the formula.

Certain of the compounds as described contain one or more chiral, or
10 asymmetric, centers and are therefore be capable of existing as optical isomers that are either dextrorotatory or levorotatory. Also included in the compounds of the invention are the respective dextrorotatory or levorotatory pure preparations, and mixtures thereof.

Certain compounds of formula (I) above are optionally provided in
15 stereoisomeric forms (e.g. they may contain one or more asymmetric carbon atoms or may exhibit cis-trans isomerism). The individual stereoisomers (enantiomers and diastereoisomers) and mixtures of these are included within the scope of the present invention. Likewise, it is understood that compounds of formula (I) are optionally provided in various tautomeric forms
20 within the scope of the present invention.

The present invention also provides compounds of formula (I) and pharmaceutically acceptable salts thereof (hereafter collectively referred to as the "active compounds") for use in medical therapy, and particularly in the treatment of disorders mediated by a Kinase, such as VEGF-R2 tyrosine
25 kinase, including, for example, as angiogenesis accompanying malignant tumor growth.

A further aspect of the invention provides a method of treatment of a human or animal suffering from a disorder mediated by a protein kinase, said treatment comprising administering an effective amount of an active
30 compound of formula (I) to the human or animal patient.

In a related aspect the present invention comprises a method for inhibiting a kinase comprising bringing said kinase into contact with a compound of formula (I).

Another aspect of the present invention provides a method for using
5 an active compound of formula (I), in the preparation of a medicament for the treatment of malignant tumors, or for the treatment of disorders involving abnormal angiogenesis, such as arthritis, diabetic retinopathy, macular degeneration and psoriasis. Alternatively, compounds of formula (I) can be used in the preparation of a medicament for the treatment of a disease
10 mediated by a kinase selected from the group consisting of: abl, ARaf, ATK, ATM, bcr-abl, Blk, BRaf, Brk, Btk, CDK1, CDK2, CDK3, CDK4, CDK5, CDK6, CDK7, CDK8, CDK9, cfms, c-fms, c-kit, c-met, cRaf1, CSF1R, CSK, c-src, EGFR, ErbB2, ErbB3, ErbB4, ERK, ERK1, ERK2, Fak, fes, FGFR1, FGFR2, FGFR3, FGFR4, FGFR5, Fgr, FLK-4, Fps, Frk, Fyn, GSK, gsk3a,
15 gsk3b, Hck, IGF-1R, IKK, IKK1, IKK2, IKK3, INS-R, Integrin-linkedkinase, Jak, JAK1, JAK2, JAK3, JNK, JNK, Lck, Lyn, MEK, MEK1, MEK2, p38, PDGFR, PIK, PKB1, PKB2, PKB3, PKC, PKC α , PKC β , PKC δ , PKC ϵ , PKC γ , PKC λ , PKC μ , PKC ζ , PLK1, Polo-like kinase, PYK2, tie1, tie2, TrkA, TrkB, TrkC, UL13, UL97, VEGF-R1, VEGF-R2, Yes and Zap70.

20 Additionally, compounds of formula (I) can be used in the preparation of a medicament for the treatment of organ transplant rejection, tumor growth, chemotherapy-induced mucositis, radiation-induced mucositis, plantar-palmar syndrome, chemotherapy-induced alopecia, chemotherapy-induced thrombocytopenia, chemotherapy-induced leukopenia and hirsutism
25 or of treating a disease state selected from the group consisting of: mucocitis, restenosis, atherosclerosis, rheumatoid arthritis, angiogenesis, hepatic cirrhosis, glomerulonephritis, diabetic nephropathy, malignant nephrosclerosis, chronic obstructive pulmonary disease, thrombotic microangiopathy, aglomerulopathy, psoriasis, diabetes mellitus,

inflammation, a neurodegenerative disease, macular degeneration, actinic keratosis and hyperproliferative disorders.


Another aspect of the present invention provides the use of an active compound of formula (I), in coadministration or alternating administration
5 with previously known anti-tumor therapies for more effective treatment of such tumors.

Another aspect of the present invention provides the use of an active compound of formula (I) in the preparation of a medicament for the treatment of viral or eukaryotic infections.

10 Other aspects of the present invention related to the inhibition of protein kinases are discussed in more detail below.

Compounds synthesized as part of the present invention which are currently preferred are listed in Tables 1 - 3 below. Compounds are identified by the numbers shown in the first column; variables below in the
15 rest of the columns are with reference to the generic structure (I). Corresponding IUPAC nomenclature are disclosed in Table 4. Since all substituents at each point of substitution are capable of independent synthesis of each other, the tables are to be read as a matrix in which any combination of substituents is within the scope of the disclosure and claims
20 of the invention.

(II)



5 Table 1

Example	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	R ⁸
1	H	H	Br	4-morpholino	H	H	H	H
2	H	H	Br	4-pyridylmethyl	H	H	H	H
3		-NH-C≡N-		CH ₃	H	H	H	H
4		-NH-N≡N-		NO ₂	H	H	H	H
5		-NH-N≡N-		NO ₂	H	H	H	H
6		-NH-N≡N-		NO ₂	H	H	H	H
7		-NH-N≡N-		NO ₂	H	H	H	H
8		-NH-N≡N-		-C(OH)(CF ₃) ₂	H	H	H	H
9		-NH-N≡N-		-CH ₂ CH ₂ OH	H	H	H	H
10		-S-CH=N-		Sme	H	H	H	H
11		-S-CH=N-		H	H	H	H	H

Example	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷
12		-S-CH=N-	H	OH	H	H	H
13		-CH=CH-CH=N-	H	H	H	H	H
14		-CH=CH-CH=N-	H	H	H	H	H
15		-CH=CH-CH=N-	H	CH ₃	H	H	H
16		-CH=CH-CH=N-	H	OMe	H	H	H
17		-NH-N=N-	H	H	H	H	H
18		-NH-N=N-	H	CH ₃	H	H	H
19		-NH-N=N-	H	OMe	H	H	H
20		-S-CH=N-	H	4-morpholino	H	H	H
21		-S-CH=N-	H	NHAc	H	H	H
22		-S-CH=N-	H	-CH ₂ CH ₂ OH	H	H	H
23		-S-CH=N-	H	4-pyridylmethyl	H	H	H
24		-S-CH=N-	H	CONH ₂	H	H	H
25	H	H	Br	4-morpholino	H	H	H
26	H	H	Br	CONH ₂	H	H	H
27	H	H	Br	NHAc	H	H	H
28	H	H	Br	-CH ₂ CH ₂ OH	H	H	H
29		-CH=CH-CH=N-	H	4-morpholino	H	H	H
30		-CH=CH-CH=N-	H	4-pyridylmethyl	H	H	H
31		-CH=CH-CH=N-	H	NHAc	H	H	H
32		-CH=CH-CH=N-	H	-CH ₂ CH ₂ OH	H	H	H
33		-CH=CH-CH=N-	H	CONH ₂	H	H	H
34		-CH=CH-CH=N-	H	OH	H	H	H
35	H	H	2-furanyl	4-morpholino	H	H	H
36	H	H	2-furanyl	NHAc	H	H	H
37	H	H	2-furanyl	-CH ₂ CH ₂ OH	H	H	H
38	H	H	-CH=CH ₂	4-morpholino	H	H	H
39	H	H	-CH=CH ₂	4-pyridylmethyl	H	H	H
40	H	H	-CH=CH ₂	NHAc	H	H	H
41	H	H	-CH=CH ₂	-CH ₂ CH ₂ OH	H	H	H
42	H	H	2-furanyl	4-pyridylmethyl	H	H	H

Example	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	R ⁸
43	H	H	2-furanyl	OH	H	H	H	H
44	H	H	2-thienyl	4-morpholino	H	H	H	H
45	H	H	2-thienyl	NHAc	H	H	H	H
46	H	H	Br	H	-CH ₂ OH	H	H	H
47	H	H	Br	5-methyl-3-pyrazolon-1-yl	H	H	H	H
48	-S-CH=N-	-S-CH=N-	H	3-ethyl-piperidine-2,6-dion-3-yl	H	H	H	H
49	-S-CH=N-	-S-CH=N-	H	OPh	H	H	H	H
50	-S-CH=N-	-S-CH=N-	H	OCH ₂ Ph	H	H	H	H
51	-S-CH=N-	-S-CH=N-	H	4-(methoxycarbonyl)phenoxy	H	H	H	H
52	-S-CH=N-	-S-CH=N-	H	3-(methoxycarbonyl)phenoxy	-CH ₂ OH	H	H	H
53	-S-CH=N-	-S-CH=N-	H	H	CONH ₂	H	H	H
54	-S-CH=N-	-S-CH=N-	H	5-methyl-3-pyrazolon-1-yl	H	H	H	H
55	-S-CH=N-	-S-CH=N-	H	CO ₂ Me	H	H	H	H
56	-S-CH=N-	-S-CH=N-	H	CN	H	H	H	H
57	-S-CH=N-	-S-CH=N-	H	NMeAc	H	H	H	H
58	-S-CH=N-	-S-CH=N-	H	OPh	H	H	H	H
59	-CH=CH-CH=N-	-CH=CH-CH=N-	H	OCH ₂ Ph	H	H	H	H
60	-CH=CH-CH=N-	-CH=CH-CH=N-	H	4-(methoxycarbonyl)phenoxy	H	H	H	H
61	-CH=CH-CH=N-	-CH=CH-CH=N-	H	3-(methoxycarbonyl)phenoxy	H	H	H	H
62	-CH=CH-CH=N-	-CH=CH-CH=N-	H	3-ethyl-piperidine-2,6-dion-3-yl	H	H	H	H
63	-CH=CH-CH=N-	-CH=CH-CH=N-	H	benzoyl	-CH ₂ OH	H	H	H
64	-CH=CH-CH=N-	-CH=CH-CH=N-	H	H	H	H	H	H
65	-CH=CH-CH=N-	-CH=CH-CH=N-	H	5-methyl-3-pyrazolon-1-yl	H	H	H	H
66	-CH=CH-CH=N-	-CH=CH-CH=N-	H	2-(4-hydroxyphenyl)ethenyl	CONH ₂	H	H	H
67	-CH=CH-CH=N-	-CH=CH-CH=N-	H	H	H	H	H	H
68	-CH=CH-CH=N-	-CH=CH-CH=N-	H	CN	H	H	H	H
69	-CH=CH-CH=N-	-CH=CH-CH=N-	H	CO ₂ Me	H	H	H	H
70	-CH=CH-CH=N-	-CH=CH-CH=N-	H	SO ₂ CH ₂ CH ₂ NEt ₂	H	H	H	H
71	-S-CH=N-	-S-CH=N-	H	SO ₂ CH ₂ CH ₂ NEt ₂	H	H	H	H
72	-CH=CH-CH=N-	-CH=CH-CH=N-	H	H	CONH ₂	H	H	H
73	H	H	Ph	H	H	H	H	H

Example	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	R ⁷	R ⁸
74	H	H	Ph	-CH ₂ PO(OEt) ₂	H	H	H	H
75	H	H	Ph	H	CN	H	H	H
76	H	H	Ph	H	-CH ₂ OH	H	-CH ₂ CH ₂ OH	H
77	H	H	Ph	H	H	H	OMe	H
78	H	H	Ph	H	H	H	H	H
79	H	H	Ph	5-methyl-3-pyrazolon-1-yl	H	H	H	H
80	H	H	Ph	I	CONH ₂	H	H	H
81	H	H	2-furanyl	H	H	H	H	H
82	H	H	2-furanyl	-CH ₂ PO(OEt) ₂	CN	H	-CH ₂ CH ₂ OH	H
83	H	H	2-furanyl	H	H	H	OMe	H
84	H	H	2-furanyl	H	-CH ₂ OH	H	H	H
85	H	H	2-furanyl	H	H	H	H	H
86	H	H	2-furanyl	5-methyl-3-pyrazolon-1-yl	H	H	H	H
87	H	H	2-furanyl	I	H	H	H	H
88	H	H	2-furanyl	H	H	H	H	H

24


R^6 H H H H H H H H H H

R^7 H H H H H H H H H H

R^8 H H H H H H H H H H

R^9 H H H H OMe H H F H H

(III)



R^4 OMe
5-oxazolyl
CH₃
2-(2-pyridyl)ethenyl
H
CH₃
CF₃
H
F
Br

R^3 H H H H H CH₃ H H H H

R^2
-NH-N=N-
-NH-N=N-
-NH-N=N-
-NH-N=N-
-S-CH=N-
OH
-NH-N=N-
-S-CH=N-
-S-CH=N-
-S-CH=N-

R^1 CH₃

Table 2

Example	R^1	R^2
89		-NH-N=N-
90		-NH-N=N-
91		-NH-N=N-
92		-NH-N=N-
93		-S-CH=N-
94	CH ₃	OH
95		-NH-N=N-
96		-S-CH=N-
97		-S-CH=N-
98		-S-CH=N-




Table 3

Example	Y	Z	A	D	R ¹	R ²	R ³	R ⁴	R ⁵	R ⁷	R ⁶	R ⁸
99	C	C	C	N	H	Ph	H	H	CONH ₂	H	H	H
100	C	C	C	N	H	Ph	H	-CH ₂ PO(OEt) ₂	H	H	H	H
101	C	C	C	N	H	Ph	H	H	CN	H	H	H
102	C	C	C	N	H	Ph	H	H	H	-CH ₂ CH ₂ OH	H	H
103	C	C	C	N	H	Ph	H	H	-CH ₂ OH	OMe	H	H
104	C	C	C	N	H	Ph	H	H	H	H	H	H
105	C	C	C	N	H	Ph	H	5-methyl-3-pyrazolon-1-yl	H	H	H	H
106	C	C	C	N	H	Ph	H	H	CONH ₂	H	H	H
107	C	C	C	N	H	2-furanyl	H	-CH ₂ PO(OEt) ₂	H	H	H	H
108	C	C	C	N	H	2-furanyl	H	H	CN	H	H	H
109	C	C	C	N	H	2-furanyl	H	H	H	-CH ₂ CH ₂ OH	H	H
110	C	C	C	N	H	2-furanyl	H	H	-CH ₂ OH	H	H	H
111	C	C	C	N	H	2-furanyl	H	H	H	OMe	H	H
112	C	C	C	N	H	2-furanyl	H	H	H	H	H	H

[illegible]

[illegible]

Standard accepted nomenclature corresponding to the Examples set forth in this specification are set forth below. In some cases nomenclature is
5 given for one or more possible isomers.

TABLE 4

Example 1	6-Bromo-3-[(Z and E)-[4-(4-morpholinyl)anilino]methylidene]-1,3-dihydro-2H-indol-2-one.
Example 2	6-Bromo-3-[(Z and E)-[4-(4-pyridinylmethyl)anilino]methylidene]-1,3-dihydro-2H-indol-2-one;
Example 3	8-[(Z and E)-4-Toluidinomethylidene]-6,8-dihydroimidazo[4,5-e]indol-7(3H)-one;
Example 4	8-[(Z and E)-(3-Methyl-4-nitroanilino)methylidene]-3,6-dihydro[1,2,3]triazolo[4,5-e]indol-7-one;
Example 5	8-[(Z and E)-[4-Nitro-3-(trifluoromethyl)anilino]methylidene]-3,6-dihydro[1,2,3]triazolo[4,5-e]indol-7-one;
Example 6	8-[(Z and E)-(3-Chloro-4-nitroanilino)methylidene]-1,6-dihydro[1,2,3]triazolo[4,5-e]indol-7-one;
Example 7	8-[(Z and E)-(3,5-Dimethyl-4-nitroanilino)methylidene]-1,6-dihydro[1,2,3]triazolo[4,5-e]indol-7-one;
Example 8	8-[(Z and E)-[4-[2,2,2-Trifluoro-1-hydroxy-1-(trifluoromethyl)ethyl]anilino]methylidene]-1,6-dihydro[1,2,3]triazolo[4,5-e]indol-7-one;
Example 9	8-[(Z and E)-[4-(2-Hydroxyethyl)anilino]methylidene]-1,6-dihydro[1,2,3]triazolo[4,5-e]indol-7-one;
Example 10	8-[(Z)-[4-(Methylsulfanyl)anilino]methylidene]-6H-[1,3]thiazolo[5,4-e]indol-7-one;
Example 11	8-[(Z)-(3,5-Dimethoxyanilino)methylidene]-6H-[1,3]thiazolo[5,4-e]indol-7-one;
Example 12	8-[(Z)-(4-Hydroxyanilino)methylidene]-6H-[1,3]thiazolo[5,4-e]indol-7-one;
Example 13	1-[(Z)-(3-Methoxyanilino)methylidene]-1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one;
Example 14	3-[[[(Z)-(2-Oxo-2,3-dihydro-1H-pyrrolo[3,2-f]quinolin-1-ylidene)methyl]amino]benzonitrile;
Example 15	1-[(Z)-4-Toluidinomethylidene]-1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one;
Example 16	1-[(Z)-(4-Methoxyanilino)methylidene]-1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one;
Example 17	3-[[[(Z and E)-7-Oxo-6,7-dihydro[1,2,3]triazolo[4,5-e]indol-8(1H)-ylidene]methyl]amino]benzonitrile;
Example 18	8-[(Z and E)-4-Toluidinomethylidene]-1,6-dihydro[1,2,3]triazolo[4,5-e]indol-7-one;

- Example 19** 8-[(Z and E)-(4-Methoxyanilino)methylidene]-1,6-dihydro[1,2,3]thiazolo[4,5-e]indol-7-one;
- Example 20** 8-[(Z)-[4-(4-Morpholinyl)anilino)methylidene]-6H-[1,3]thiazolo[5,4-e]indol-7-one;
- Example 21** N-(4-[(Z)-(7-Oxo-6,7-dihydro-8H-[1,3]thiazolo[5,4-e]indol-8-ylidene)methyl]amino}phenyl)acetamide;
- Example 22** 8-[(Z)-[4-(2-Hydroxyethyl)anilino)methylidene]-6H-[1,3]thiazolo[5,4-e]indol-7-one;
- Example 23** 8-[(Z)-[4-(4-Pyridinylmethyl)anilino)methylidene]-6H-[1,3]thiazolo[5,4-e]indol-7-one;
- Example 24** 4-[(Z)-(7-Oxo-6,7-dihydro-8H-[1,3]thiazolo[5,4-e]indol-8-ylidene)methyl]amino}benzamide;
- Example 25** 6-Bromo-3-[(Z and E)-[4-(4-morpholinyl)anilino)methylidene]-1,3-dihydro-2H-indol-2-one;
- Example 26** 4-[(Z and E)-(6-Bromo-2-oxo-1,2-dihydro-3H-indol-3-ylidene)methyl]amino}benzamide;
- Example 27** N-(4-[(Z and E)-(6-Bromo-2-oxo-1,2-dihydro-3H-indol-3-ylidene)methyl]amino}phenyl)acetamide;
- Example 28** 6-Bromo-3-[(Z and E)-[4-(2-hydroxyethyl)anilino)methylidene]-1,3-dihydro-2H-indol-2-one;
- Example 29** 1-[(Z)-[4-(4-Morpholinyl)anilino)methylidene]-1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one;
- Example 30** 1-[(Z)-[4-(4-Pyridinylmethyl)anilino)methylidene]-1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one;
- Example 31** N-(4-[(Z)-(2-Oxo-2,3-dihydro-1H-pyrrolo[3,2-f]quinolin-1-ylidene)methyl]amino}phenyl)acetamide;
- Example 32** 1-[(Z)-[4-(2-Hydroxyethyl)anilino)methylidene]-1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one;
- Example 33** 4-[(Z)-(2-Oxo-2,3-dihydro-1H-pyrrolo[3,2-f]quinolin-1-ylidene)methyl]amino}benzamide;
- Example 34** 1-[(Z)-(4-Hydroxyanilino)methylidene]-1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one;
- Example 35** 6-(2-Furyl)-3-[(Z and E)-[4-(4-morpholinyl)anilino)methylidene]-1,3-dihydro-2H-indol-2-one;
- Example 36** N-[4-[(Z and E)-[6-(2-Furyl)-2-oxo-1,2-dihydro-3H-indol-3-ylidene)methyl]amino}phenyl]acetamide;
- Example 37** 6-(2-Furyl)-3-[(Z and E)-[4-(2-hydroxyethyl)anilino)methylidene]-1,3-dihydro-2H-indol-2-one;
- Example 38** 3-[(Z and E)-[4-(4-Morpholinyl)anilino)methylidene]-6-vinyl-1,3-dihydro-2H-indol-2-one;
- Example 39** 3-[(Z and E)-[4-(4-Pyridinylmethyl)anilino)methylidene]-6-vinyl-1,3-dihydro-2H-indol-2-one;
- Example 40** N-(4-[(Z and E)-[(2-Oxo-6-vinyl-1,2-dihydro-3H-indol-3-ylidene)methyl]amino}phenyl)acetamide;

- Example 41** 3-((Z and E)-[4-(2-Hydroxyethyl)anilino]methylidene)-6-vinyl-1,3-dihydro-2H-indol-2-one;
- Example 42** 6-(2-Furyl)-3-((Z and E)-[4-(4-pyridinylmethyl)anilino]methylidene)-1,3-dihydro-2H-indol-2-one;
- Example 43** 6-(2-Furyl)-3-((Z and E)-(4-hydroxyanilino)methylidene)-1,3-dihydro-2H-indol-2-one;
- Example 44** 3-((Z and E)-[4-(4-Morpholinyl)anilino]methylidene)-6-(2-thienyl)-1,3-dihydro-2H-indol-2-one;
- Example 45** N-[4-((Z and E)-[2-Oxo-6-(2-thienyl)-1,2-dihydro-3H-indol-3-ylidene]methyl)amino]phenyl]acetamide;
- Example 46** 6-Bromo-3-((Z and E)-[3-(hydroxymethyl)anilino]methylidene)-1,3-dihydro-2H-indol-2-one;
- Example 47** 6-Bromo-3-((Z and E)-[4-(5-methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene)-1,3-dihydro-2H-indol-2-one;
- Example 48** 3-Ethyl-3-(4-((Z and E)-[(7-oxo-6,7-dihydro-8H-[1,3]thiazolo[5,4-e]indol-8-ylidene)methyl]amino)phenyl)-2,6-piperidinedione;
- Example 49** 8-((Z)-(4-Phenoxyanilino)methylidene)-6H-[1,3]thiazolo[5,4-e]indol-7-one;
- Example 50** 8-((Z)-[4-(Benzyloxy)anilino]methylidene)-6H-[1,3]thiazolo[5,4-e]indol-7-one;
- Example 51** Methyl 4-(4-(((Z)-7-oxo-6,7-dihydro-8H-[1,3]thiazolo[5,4-e]indol-8-ylidene)methyl)amino)phenoxy)benzoate;
- Example 52** Methyl 3-(4-((Z)-[(7-oxo-6,7-dihydro-8H-[1,3]thiazolo[5,4-e]indol-8-ylidene)methyl]amino)phenoxy)benzoate;
- Example 53** 8-((Z)-[3-(Hydroxymethyl)anilino]methylidene)-6H-[1,3]thiazolo[5,4-e]indol-7-one;
- Example 54** 3-((Z)-[(7-Oxo-6,7-dihydro-8H-[1,3]thiazolo[5,4-e]indol-8-ylidene)methyl]amino)benzamide;
- Example 55** 8-((Z)-[4-(5-Methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene)-6H-[1,3]thiazolo[5,4-e]indol-7-one;
- Example 56** Methyl 4-((Z)-[(7-Oxo-6,7-dihydro-8H-[1,3]thiazolo[5,4-e]indol-8-ylidene)methyl]amino)benzoate;
- Example 57** 4-((Z)-[(7-Oxo-6,7-dihydro-8H-[1,3]thiazolo[5,4-e]indol-8-ylidene)methyl]amino)benzonitrile;
- Example 58** N-Methyl-N-(4-((Z)-[(7-oxo-6,7-dihydro-8H-[1,3]thiazolo[5,4-e]indol-8-ylidene)methyl]amino)phenyl)acetamide;
- Example 59** 1-((Z)-(4-Phenoxyanilino)methylidene)-1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one;
- Example 60** 1-((Z)-[4-(Benzyloxy)anilino]methylidene)-1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one;
- Example 61** Methyl 4-(4-((Z)-[(2-Oxo-2,3-dihydro-1H-pyrrolo[3,2-f]quinolin-1-ylidene)methyl]amino)phenoxy)benzoate;
- Example 62** Methyl 3-(4-((Z)-[(2-Oxo-2,3-dihydro-1H-pyrrolo[3,2-f]quinolin-1-ylidene)methyl]amino)phenoxy)benzoate;

- Example 63** 3-Ethyl-3-(4-((Z)-[(2-oxo-2,3-dihydro-1H-pyrrolo[3,2-f]quinolin-1-ylidene)methyl]amino)phenyl)-2,6-piperidinedione;
- Example 64** 1-((Z)-(4-Benzoylanilino)methylidene)-1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one;
- Example 65** 1-((Z)-[3-(Hydroxymethyl)anilino]methylidene)-1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one;
- Example 66** 1-((Z)-[4-(5-Methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene)-1H-pyrrolo[3,2-f]quinolin-2(3H)-one
- Example 67** 1-((E)-{4-[(E)-2-(4-Hydroxyphenyl)ethenyl]anilino}methylidene)-1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one;
- Example 68** 3-((Z)-[(2-Oxo-2,3-dihydro-1H-pyrrolo[3,2-f]quinolin-1-ylidene)methyl]amino)benzamide;
- Example 69** 4-((Z)-[(2-Oxo-2,3-dihydro-1H-pyrrolo[3,2-f]quinolin-1-ylidene)methyl]amino)benzonitrile;
- Example 70** Methyl 4-((Z)-[(2-oxo-2,3-dihydro-1H-pyrrolo[3,2-f]quinolin-1-ylidene)methyl]amino)benzoate;
- Example 71** 8-((Z)-(4-{[2-(Diethylamino)ethyl]sulfonyl}anilino)methylidene)-6H-[1,3]thiazolo[5,4-e]indol-7-one;
- Example 72** 1-((Z)-(4-{[2-(Diethylamino)ethyl]sulfonyl}anilino)methylidene)-1H-pyrrolo[3,2-f]quinolin-2(3H)-one;
- Example 73** 3-(((Z and E)-(2-Oxo-6-phenyl-1,2-dihydro-3H-indol-3-ylidene)methyl]amino)benzamide;
- Example 74** Diethyl 4-(((Z and E)-(2-oxo-6-phenyl-1,2-dihydro-3H-indol-3-ylidene)methyl]amino)benzylphosphonate;
- Example 75** 3-(((Z and E)-(2-Oxo-6-phenyl-1,2-dihydro-3H-indol-3-ylidene)methyl]amino)benzonitrile;
- Example 76** 3-((Z and E)-[2-(2-Hydroxyethyl)anilino]methylidene)-6-phenyl-1,3-dihydro-2H-indol-2-one;
- Example 77** 3-((Z and E)-[3-(Hydroxymethyl)anilino]methylidene)-6-phenyl-1,3-dihydro-2H-indol-2-one;
- Example 78** 3-((Z and E)-(2-Methoxyanilino)methylidene)-6-phenyl-1,3-dihydro-2H-indol-2-one;
- Example 79** 3-((Z and E)-[4-(5-Methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene)-6-phenyl-1H-indol-2-one;
- Example 80** 3-((Z and E)-(4-Iodoanilino)methylidene)-6-phenyl-1H-indol-2-one;
- Example 81** 3-(((Z and E)-[6-(2-Furyl)-2-oxo-1,2-dihydro-3H-indol-3-ylidene)methyl]amino)benzamide;
- Example 82** Diethyl 4-(((Z and E)-[6-(2-furyl)-2-oxo-1,2-dihydro-3H-indol-3-ylidene)methyl]amino)benzylphosphonate;
- Example 83** 3-(((Z and E)-[6-(2-Furyl)-2-oxo-1,2-dihydro-3H-indol-3-ylidene)methyl]amino)benzonitrile;
- Example 84** 6-(2-Furyl)-3-((Z and E)-[2-(2-hydroxyethyl)anilino]methylidene)-1,3-dihydro-2H-indol-2-one;

- Example 85** 6-(2-Furyl)-3-((Z and E)-[3-(hydroxymethyl)anilino]methylidene)-1,3-dihydro-2H-indol-2-one;
- Example 86** 6-(2-Furyl)-3-[(Z and E)-(2-methoxyanilino)methylidene]-1,3-dihydro-2H-indol-2-one;
- Example 87** 6-(2-Furyl)-3-((Z and E)-[4-(5-methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene)-1H-indol-2-one;
- Example 88** 6-(2-Furyl)-3-[(Z and E)-(4-iodoanilino)methylidene]-1H-indol-2-one;
- Example 89** 3,6-Dihydro[1,2,3]triazolo[4,5-e]indole-7,8-dione8-[N-(4-methoxyphenyl)hydrazone];
- Example 90** 3,6-Dihydro[1,2,3]triazolo[4,5-e]indole-7,8-dione8-[N-[4-(1,3-oxazol-5-yl)phenyl]hydrazone];
- Example 91** 3,6-Dihydro[1,2,3]triazolo[4,5-e]indole-7,8-dione8-[N-(4-methylphenyl)hydrazone];
- Example 92** 3,6-Dihydro[1,2,3]triazolo[4,5-e]indole-7,8-dione8-(N-[4-[(E)-2-(2-pyridinyl)ethenyl]phenyl]hydrazone);
- Example 93** 6H-[1,3]Thiazolo[5,4-e]indole-7,8-dione8-[N-(3-methoxyphenyl)hydrazone];
- Example 94** 5-Hydroxy-4,6-dimethyl-1H-indole-2,3-dione3-[N-(4-methylphenyl)hydrazone];
- Example 95** 3,6-Dihydro[1,2,3]triazolo[4,5-e]indole-7,8-dione8-[N-[4-(trifluoromethyl)phenyl]hydrazone];
- Example 96** 6H-[1,3]Thiazolo[5,4-e]indole-7,8-dione8-[N-(3-fluorophenyl)hydrazone];
- Example 97** 6H-[1,3]Thiazolo[5,4-e]indole-7,8-dione8-[N-(4-fluorophenyl)hydrazone];
- Example 98** 6H-[1,3]Thiazolo[5,4-e]indole-7,8-dione8-[N-(4-bromophenyl)hydrazone];
- Example 99** 3-[[2-Oxo-5-phenyl-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene)methyl]amino}benzamide;
- Example 100** Diethyl 4-[[2-oxo-5-phenyl-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene)methyl]amino}benzylphosphonate;
- Example 101** 3-[[2-Oxo-5-phenyl-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene)methyl]amino}benzonitrile;
- Example 102** 3-[[2-(2-Hydroxyethyl)anilino]methylidene]-5-phenyl-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;
- Example 103** 3-[[3-(Hydroxymethyl)anilino]methylidene]-5-phenyl-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;
- Example 104** 3-[(2-Methoxyanilino)methylidene]-5-phenyl-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;
- Example 105** 3-[[4-(5-Methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene]-5-phenyl-1H-pyrrolo[2,3-b]pyridin-2-one;
- Example 106** 3-[(4-Iodoanilino)methylidene]-5-phenyl-1H-pyrrolo[2,3-b]pyridin-2-one;

- Example 107** 3-({[5-(2-Furyl)-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene]methyl}amino)benzamide;
- Example 108** Diethyl 4-({[5-(2-furyl)-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene]methyl}amino)benzylphosphonate;
- Example 109** 3-({[5-(2-Furyl)-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene]methyl}amino)benzonitrile;
- Example 110** 5-(2-Furyl)-3-({[2-(2-hydroxyethyl)anilino]methylidene}-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;
- Example 111** 5-(2-Furyl)-3-({[3-(hydroxymethyl)anilino]methylidene}-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;
- Example 112** 5-(2-Furyl)-3-({[2-methoxyanilino]methylidene}-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;
- Example 113** 5-(2-Furyl)-3-({[4-(5-methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene}-1H-pyrrolo[2,3-b]pyridin-2-one;
- Example 114** 5-(2-Furyl)-3-({[4-iodoanilino]methylidene}-1H-pyrrolo[2,3-b]pyridin-2-one;
- Example 115** 3-({[2-Oxo-5-(3-thienyl)-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene]methyl}amino)benzamide;
- Example 116** Diethyl 4-({[2-oxo-5-(3-thienyl)-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene]methyl}amino)benzylphosphonate;
- Example 117** 3-({[2-Oxo-5-(3-thienyl)-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene]methyl}amino)benzonitrile;
- Example 118** 3-({[2-(2-Hydroxyethyl)anilino]methylidene}-5-(3-thienyl)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;
- Example 119** 3-({[3-(Hydroxymethyl)anilino]methylidene}-5-(3-thienyl)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;
- Example 120** 3-({[2-Methoxyanilino]methylidene}-5-(3-thienyl)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;
- Example 121** 3-({[4-(5-Methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene}-5-(3-thienyl)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;
- Example 122** 3-({[4-Iodoanilino]methylidene}-5-(3-thienyl)-1H-pyrrolo[2,3-b]pyridin-2-one;
- Example 123** 3-({[5-Bromo-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene]methyl}amino)benzamide;
- Example 124** Diethyl 4-({[5-Bromo-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene]methyl}amino)benzylphosphonate;
- Example 125** 3-({[5-Bromo-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene]methyl}amino)benzonitrile;
- Example 126** 5-Bromo-3-({[2-(2-hydroxyethyl)anilino]methylidene}-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;
- Example 127** 5-Bromo-3-({[3-(hydroxymethyl)anilino]methylidene}-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

- Example 128** 5-Bromo-3-[(2-methoxyanilino)methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;
- Example 129** 5-Bromo-3-[[4-(5-methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;
- Example 130** 5-Bromo-3-[(4-iodoanilino)methylidene]-1H-pyrrolo[2,3-b]pyridin-2-one;
- Example 131** 3-[[[(6-Chloro-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene)methyl]amino]benzamide;
- Example 132** Diethyl 4-[[[(6-chloro-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene)methyl]amino]benzylphosphonate;
- Example 133** 3-[[[(6-Chloro-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene)methyl]amino]benzonitrile;
- Example 134** 6-Chloro-3-[[2-(2-hydroxyethyl)anilino]methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;
- Example 135** 6-Chloro-3-[[3-(hydroxymethyl)anilino]methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;
- Example 136** 6-Chloro-3-[(2-methoxyanilino)methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;
- Example 137** 6-Chloro-3-[[4-(5-methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;
- Example 138** 6-Chloro-3-[(4-iodoanilino)methylidene]-1H-pyrrolo[2,3-b]pyridin-2-one;
- Example 139** Ethyl 3-[[3-(aminocarbonyl)anilino]methylidene]-2-oxo-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine-5-carboxylate;
- Example 140** Ethyl 3-[[4-[(diethoxyphosphoryl)methyl]anilino]methylidene]-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridine-5-carboxylate;
- Example 141** Ethyl 3-[[3-(cyanoanilino)methylidene]-2-oxo-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine-5-carboxylate;
- Example 142** Ethyl 3-[[2-(2-hydroxyethyl)anilino]methylidene]-2-oxo-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine-5-carboxylate;
- Example 143** Ethyl 3-[[3-(hydroxymethyl)anilino]methylidene]-2-oxo-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine-5-carboxylate;
- Example 144** Ethyl 3-[(2-methoxyanilino)methylidene]-2-oxo-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine-5-carboxylate;
- Example 145** Ethyl 3-[[4-(5-methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene]-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridine-5-carboxylate;
- Example 144** Ethyl 3-[(4-iodoanilino)methylidene]-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridine-5-carboxylate;
- Example 147** 3-[[[(2-Oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene)methyl]amino]benzamide;
- Example 148** Diethyl 4-[[[(2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene)methyl]amino]benzylphosphonate;

- Example 149** 3-[[2-Oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene)methyl]amino}benzonitrile;
- Example 150** 3-{[2-(2-Hydroxyethyl)anilino]methylidene}-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;
- Example 151** 3-{[3-(Hydroxymethyl)anilino]methylidene}-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;
- Example 152** 3-[(2-Methoxyanilino)methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;
- Example 153** 3-{[4-(5-Methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene}-1H-pyrrolo[2,3-b]pyridin-2-one;
- Example 154** 3-[(4-Iodoanilino)methylidene]-1H-pyrrolo[2,3-b]pyridin-2-one;
- Example 155** 3-[[2-Oxo-1,2-dihydro-3H-pyrrolo[3,2-b]pyridin-3-ylidene)methyl]amino}benzamide;
- Example 156** Diethyl 4-[[2-oxo-1,2-dihydro-3H-pyrrolo[3,2-b]pyridin-3-ylidene)methyl]amino}benzylphosphonate;
- Example 157** 3-[[2-Oxo-1,2-dihydro-3H-pyrrolo[3,2-b]pyridin-3-ylidene)methyl]amino}benzonitrile;
- Example 158** 3-{[2-(2-Hydroxyethyl)anilino]methylidene}-1,3-dihydro-2H-pyrrolo[3,2-b]pyridin-2-one;
- Example 159** 3-{[3-(Hydroxymethyl)anilino]methylidene}-1,3-dihydro-2H-pyrrolo[3,2-b]pyridin-2-one;
- Example 160** 3-[(2-Methoxyanilino)methylidene]-1,3-dihydro-2H-pyrrolo[3,2-b]pyridin-2-one;
- Example 161** 3-{[4-(5-Methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene}-1H-pyrrolo[3,2-b]pyridin-2-one; and
- Example 162** 3-[(4-Iodoanilino)methylidene]-1H-pyrrolo[3,2-b]pyridin-2-one.
- Example 163** 3-{(Z and E)-[(4-Hydroxy-1-naphthyl)amino]methylidene}-5-phenyl-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;
- Example 164** 5-(2-Furyl)-3-{(Z and E)-[(4-hydroxy-1-naphthyl)amino]methylidene}-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;
- Example 165** 3-{(Z and E)-[(4-Hydroxy-1-naphthyl)amino]methylidene}-5-(3-thienyl)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;
- Example 166** 5-Bromo-3-{(Z and E)-[(4-hydroxy-1-naphthyl)amino]methylidene}-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;
- Example 167** 6-Chloro-3-{(Z and E)-[(4-hydroxy-1-naphthyl)amino]methylidene}-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;
- Example 168** Ethyl 3-{(Z and E)-[(4-hydroxy-1-naphthyl)amino]methylidene}-2-oxo-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine-5-carboxylate;
- Example 169** 3-{(Z and E)-[(4-Hydroxy-1-naphthyl)amino]methylidene}-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;
- Example 170** 3-{(Z and E)-[(4-Hydroxy-1-naphthyl)amino]methylidene}-1,3-dihydro-2H-pyrrolo[3,2-b]pyridin-2-one;
- Example 171** 6-(2-Furyl)-3-{(Z and E)-[(4-hydroxy-1-naphthyl)amino]methylidene}-1,3-dihydro-2H-indol-2-one;

Example 172 3-((Z and E)-[(4-Hydroxy-1-naphthyl)amino]methylidene)-6-phenyl-1,3-dihydro-2H-indol-2-one;

The invention discloses thirteen different points of substitution on structural formula (I). Each of these points of substitution bears a substituent
5 whose selection and synthesis as part of this invention is independent of all other points of substitution on formula (I). Each point of substitution is now further described.

Preferred substitutions for Y, Z, A and D are nitrogen or carbon. The most preferred substituents for D are nitrogen and carbon. The most highly
10 preferred substituent for Y, Z, A and D is carbon.

Preferred substitutions at the R¹ position include hydrogen, fluoro, bromo, iodo, lower alkyl, cyano and nitro. Alternatively, R¹ is optionally joined with R² to form a fused ring structure selected from the group consisting of: thiazole, imidazole, triazole and pyridine. Further, such fused
15 ring structures are optionally substituted by one or more substituents selected from the group consisting of: halogen, amino, lower alkyl substituted amino, lower alkyl and lower alkyl carbonyl. In a preferred embodiment, R¹ is selected from the group which includes hydrogen and methyl or R¹ is fused with R² to form a ring structure selected from the group
20 which includes fused thiazole and fused pyridine. In a most highly preferred embodiment, R¹ is fused with R² to form a ring structure selected from the group which includes fused thiazole, pyridine and pyridine substituted by halogen or methyl.

Preferred substitutions at the R² position include hydrogen, lower
25 alkyl, lower alkoxy, hydroxy lower alkyl, C₁₋₁₂ alkoxy carbonyl, Aryl, Het, aminocarbonyl, lower alkyl aminocarbonyl, halogen and hydroxy. Alternatively, R² is fused with R¹ to form a fused ring selected from the group which includes thiazole, imidazole, triazole and pyridine. Such fused rings are optionally substituted by a substituent selected from the group which

includes halogen, amino, lower alkyl substituted amino, lower alkyl and lower alkyl carbonyl. Most preferably, R^2 is selected from the group which includes hydroxyl, hydroxy and lower alkyl, or is fused with R^1 to form a ring structure selected from the group which includes fused thiazole and fused pyridine. In
5 a most highly preferred embodiment, R^2 is selected from the group which includes hydroxy and hydroxymethyl, or is fused with R^1 to form a fused ring from the group which includes fused thiazole, pyridine and pyridine substituted by halogen or methyl.

Preferred substitutions at R^3 include hydrogen, lower alkyl, lower
10 alkenyl, halogen, phenyl, Het and alkoxy. Most preferred are hydrogen, halogen, ethenyl and methyl. Most highly preferred substitutions at R^3 are hydrogen and bromo.

Preferred substitutions at R^4 include hydrogen, lower alkyl, hydroxy, hydroxy-lower alkyl, carboxamide, mono- lower alkyl aminocarbonyl,
15 substituted Aryl-lower alkyl, Het, Het-lower alkyl, lower alkoxy, Aryloxy, Het-oxy, amino, mono- or di- lower alkyl-amino lower alkyl aminocarbonyl, mono- or di- lower alkyl-amino lower alkoxycarbonyl, mono- or di-lower alkyl-amino lower alkyl aminocarbonylamino, mono- or di- lower alkyl-amino lower alkoxycarbonylamino, lower alkyl carbonylamino, (lower alkyl carbonyl)(lower
20 alkyl)amino, mono- or di-lower alkyl-amino lower alkyl carbonylamino, [mono- or di-lower alkyl-amino lower alkyl carbonyl][lower alkyl]amino, mono- or di-lower alkyl-amino lower alkyl sulfonylamino, [mono- or di-lower alkyl-amino lower alkyl sulfonyl][lower alkyl]amino, mono- or di- lower alkyl-amino lower alkyl sulfonyl, Het lower alkyl aminocarbonyl, Het lower alkyl
25 aminocarbonylamino, Het lower alkoxycarbonylamino, Het lower alkyl carbonyl, Het lower alkoxycarbonyl, lower alkyl sulfonyl lower alkyl aminoalkyl, lower alkyl sulfonyl-lower alkyl-aminoalkyl-Het-, lower alkoxycarbonyl, halogen, cyano, diethoxyphosphorylmethyl, trifluoromethyl and trifluoromethoxy. The most preferred substitutions are lower alkyl,
30 hydroxy, hydroxy-lower alkyl, carboxamide, mono- lower alkyl

aminocarbonyl, substituted Aryl-lower alkyl, Het, Het-lower alkyl, Het-oxy, mono- or di- lower alkyl-amino lower alkyl aminocarbonyl, mono- or di- lower alkyl-amino lower alkoxy carbonyl, mono- or di-lower alkyl-amino lower alkyl aminocarbonylamino, mono- or di- lower alkyl-amino lower
 5 alkoxy carbonylamino, lower alkyl carbonylamino, (lower alkyl carbonyl)(lower alkyl)amino, mono- or di-lower alkyl-amino lower alkyl carbonylamino, [mono- or di-lower alkyl-amino lower alkyl carbonyl][lower alkyl]amino, mono- or di-lower alkyl-amino lower alkyl sulfonylamino, [mono- or di-lower alkyl-amino lower alkyl sulfonyl][lower alkyl]amino, mono- or di- lower alkyl-amino
 10 lower alkyl sulfonyl, Het lower alkyl aminocarbonyl, Het lower alkyl carbonyl, lower alkyl sulfonyl lower alkyl aminoalkyl, lower alkyl sulfonyl-lower alkyl-aminoalkyl-Het-, halogen, cyano and trifluoromethyl. Most highly preferred are hydroxymethyl, hydroxyethyl, 4-pyridylmethyl, 4-morpholino, acetamido, N-methylacetamido, carboxamide, diethylaminoethylsulfonyl, 5-methyl-3-
 15 pyrazolon-1-yl and 3-ethyl-piperidine-2,6-dione-3-yl.

Preferred substitutions at R⁵ include hydrogen, lower alkyl, hydroxy, hydroxy-lower alkyl, carboxamide, mono- lower alkyl aminocarbonyl, substituted Aryl-lower alkyl, Het, Het-lower alkyl, lower alkoxy, Aryloxy, Het-oxy, amino, mono- or di- lower alkyl-amino lower alkyl aminocarbonyl, mono-
 20 or di- lower alkyl-amino lower alkoxy carbonyl, mono- or di-lower alkyl-amino lower alkyl aminocarbonylamino, mono- or di- lower alkyl-amino lower alkoxy carbonylamino, lower alkyl carbonylamino, (lower alkyl carbonyl)(lower alkyl)amino, mono- or di-lower alkyl-amino lower alkyl carbonylamino, [mono- or di-lower alkyl-amino lower alkyl carbonyl][lower alkyl]amino, mono-
 25 or di-lower alkyl-amino lower alkyl sulfonylamino, [mono- or di-lower alkyl-amino lower alkyl sulfonyl][lower alkyl]amino, mono- or di- lower alkyl-amino lower alkyl sulfonyl, Het lower alkyl aminocarbonyl, Het lower alkyl aminocarbonylamino, Het lower alkoxy carbonylamino, Het lower alkyl carbonyl, Het lower alkoxy carbonyl, lower alkyl sulfonyl lower alkyl
 30 aminoalkyl, lower alkyl sulfonyl-lower alkyl-aminoalkyl-Het-, lower

alkoxycarbonyl, halogen, cyano, diethoxyphosphorylmethyl, trifluoromethyl and trifluoromethoxy. The most preferred substitutions are lower alkyl, hydroxy, hydroxy-lower alkyl, carboxamide, mono- lower alkyl aminocarbonyl, substituted Aryl-lower alkyl, Het, Het-lower alkyl, Het-oxy, 5 mono- or di- lower alkyl-amino lower alkyl aminocarbonyl, mono- or di- lower alkyl-amino lower alkoxycarbonyl, mono- or di-lower alkyl-amino lower alkyl aminocarbonylamino, mono- or di- lower alkyl-amino lower alkoxycarbonylamino, lower alkyl carbonylamino, (lower alkyl carbonyl)(lower alkyl) amino, mono- or di-lower alkyl-amino lower alkyl carbonylamino, 10 [mono- or di-lower alkyl-amino lower alkyl carbonyl][lower alkyl]amino, mono- or di-lower alkyl-amino lower alkyl sulfonylamino, [mono- or di-lower alkyl-amino lower alkyl sulfonyl][lower alkyl]amino, mono- or di- lower alkyl-amino lower alkyl sulfonyl, Het lower alkyl aminocarbonyl, Het lower alkyl carbonyl, lower alkyl sulfonyl lower alkyl aminoalkyl, lower alkyl sulfonyl-lower alkyl- 15 aminoalkyl-Het-, halogen, cyano and trifluoromethyl. Most highly preferred are hydroxymethyl, hydroxyethyl, 4-pyridylmethyl, 4-morpholino, acetamido, N-methylacetamido, carboxamide, diethylaminoethylsulfonyl, 5-methyl-3-pyrazolon-1-yl and 3-ethyl-piperidine-2,6-dion-3-yl.

The most preferred substitution at R⁶ is hydrogen.





20

Preferred substitutions at R⁷ and R⁸ are hydrogen, halogen and methyl.






Another preferred substitution at R⁷ includes the state in which R⁷ is joined to R⁵ so as to form a fused benzo ring from R⁵ to R⁷.

25 Preferred substitutions at X include N, CH and CCH₃. Most preferred is CH.


Preferred individual compounds of the present invention are selected from the group consisting of:



Highly preferred compounds include
5



; and



DETAILED DESCRIPTION OF THE INVENTION

Salts encompassed within the term "pharmaceutically acceptable salts" refer to non-toxic salts of the compounds of this invention which are generally prepared by reacting the free base with a suitable organic or inorganic acid or by reacting the acid with a suitable organic or inorganic base. Representative salts include the following salts: Acetate, Benzenesulfonate, Benzoate, Bicarbonate, Bisulfate, Bitartrate, Borate, Bromide, Calcium Edetate, Camsylate, Carbonate, Chloride, Clavulanate, Citrate, Diethanolamine, Dihydrochloride, Edetate, Edisylate, Estolate, Esylate, Fumarate, Gluceptate, Gluconate, Glutamate, Glycolylarsanilate, Hexylresorcinate, Hydrabamine, Hydrobromide, Hydrochloride, Hydroxynaphthoate, Iodide, Isethionate, Lactate, Lactobionate, Laurate, Malate, Maleate, Mandelate, Mesylate, Metaphosphoric, Methylbromide, Methylnitrate, Methylsulfate, Monopotassium Maleate, Mucate, Napsylate, Nitrate, N-methylglucamine, Oxalate, Pamoate (Embonate), Palmitate, Pantothenate, Phosphate/diphosphate, Polygalacturonate, Potassium, Salicylate, Sodium, Stearate, Subacetate, Succinate, Tannate, Tartrate, Teoclate, Tosylate, Trifluoroacetate, Triethiodide, Trimethylammonium and Valerate.

Other salts which are not pharmaceutically acceptable may be useful in the preparation of compounds of formula (I) and these form a further aspect of the invention.

Also included within the scope of the invention are the individual isomers of the compounds represented by formula (I) above as well as any wholly or partially equilibrated mixtures thereof. The present invention also covers the individual isomers of the compounds represented by formula above as mixtures with isomers thereof in which one or more chiral asymmetric centers are inverted.

As used herein, the term "aliphatic" refers to the terms alkyl, alkylene, alkenyl, alkenylene, alkynyl and alkynylene.

As used herein, the term "lower" refers to a group having between one and six carbons.

As used herein, the term "alkyl" refers to a straight or branched chain hydrocarbon having from one to twelve carbon atoms, optionally substituted
5 with substituents selected from the group which includes lower alkyl, lower alkoxy, lower alkylsulfanyl, lower alkylsulfenyl, lower alkylsulfonyl, oxo, hydroxy, mercapto, amino optionally substituted by alkyl, carboxy, carbamoyl optionally substituted by alkyl, aminosulfonyl optionally substituted by a substituent selected from the group including alkyl, nitro, cyano, halogen and
10 lower perfluoroalkyl, multiple degrees of substitution being allowed. Examples of "alkyl" as used herein include, but are not limited to, n-butyl, n-pentyl, isobutyl, isopropyl and the like.

As used herein, the term "alkylene" refers to a straight or branched chain divalent hydrocarbon radical having from one to ten carbon atoms,
15 optionally substituted with substituents selected from the group which includes lower alkyl, lower alkoxy, lower alkylsulfanyl, lower alkylsulfenyl, lower alkylsulfonyl, oxo, hydroxy, mercapto, amino optionally substituted by alkyl, carboxy, carbamoyl optionally substituted by alkyl, aminosulfonyl optionally substituted by alkyl, nitro, cyano, halogen and lower perfluoroalkyl,
20 multiple degrees of substitution being allowed. Examples of "alkylene" as used herein include, but are not limited to, methylene, ethylene, and the like.

As used herein, the term "alkenyl" refers to a hydrocarbon radical having from two to ten carbons and at least one carbon-carbon double bond, optionally substituted with substituents selected from the group which
25 includes lower alkyl, lower alkoxy, lower alkylsulfanyl, lower alkylsulfenyl, lower alkylsulfonyl, oxo, hydroxy, mercapto, amino optionally substituted by alkyl, carboxy, carbamoyl optionally substituted by alkyl, aminosulfonyl optionally substituted by alkyl, nitro, cyano, halogen and lower perfluoroalkyl, multiple degrees of substitution being allowed.

As used herein, the term "alkenylene" refers to an straight or branched chain divalent hydrocarbon radical having from two to ten carbon atoms and one or more carbon - carbon double bonds, optionally substituted with substituents selected from the group which includes lower alkyl, lower
5 alkoxy, lower alkylsulfanyl, lower alkylsulfenyl, lower alkylsulfonyl, oxo, hydroxy, mercapto, amino optionally substituted by alkyl, carboxy, carbamoyl optionally substituted by alkyl, aminosulfonyl optionally substituted by alkyl, nitro, cyano, halogen and lower perfluoroalkyl, multiple degrees of substitution being allowed. Examples of "alkenylene" as used herein
10 include, but are not limited to, ethene-1,2-diyl, propene-1,3-diyl, methylene-1,1-diyl, and the like.

As used herein, the term "alkynyl" refers to a hydrocarbon radical having from two to ten carbons and at least one carbon-carbon triple bond, optionally substituted with substituents selected from the group which
15 includes lower alkyl, lower alkoxy, lower alkylsulfanyl, lower alkylsulfenyl, lower alkylsulfonyl, oxo, hydroxy, mercapto, amino optionally substituted by alkyl, carboxy, carbamoyl optionally substituted by alkyl, aminosulfonyl optionally substituted by alkyl, nitro, cyano, halogen and lower perfluoroalkyl, multiple degrees of substitution being allowed.

20 As used herein, the term "alkynylene" refers to a straight or branched chain divalent hydrocarbon radical having from two to ten carbon atoms and one or more carbon - carbon triple bonds, optionally substituted with substituents selected from the group which includes lower alkyl, lower alkoxy, lower alkylsulfanyl, lower alkylsulfenyl, lower alkylsulfonyl, oxo,
25 hydroxy, mercapto, amino optionally substituted by alkyl, carboxy, carbamoyl optionally substituted by alkyl, aminosulfonyl optionally substituted by alkyl, nitro, cyano, halogen and lower perfluoroalkyl, multiple degrees of substitution being allowed. Examples of "alkynylene" as used herein include, but are not limited to, ethyne-1,2-diyl, propyne-1,3-diyl, and the like.

As used herein, the term "cycloaliphatic" includes the terms cycloalkyl, cycloalkylene, cycloalkenyl, cycloalkenylene, cycloalkynyl and cycloalkylnylene.

As used herein, "cycloalkyl" refers to a alicyclic hydrocarbon group
5 with one or more degrees of unsaturation, having from three to twelve carbon atoms, optionally substituted with substituents selected from the group which includes lower alkyl, lower alkoxy, lower alkylsulfanyl, lower alkylsulfenyl, lower alkylsulfonyl, oxo, hydroxy, mercapto, amino optionally substituted by alkyl, carboxy, carbamoyl optionally substituted by alkyl, aminosulfonyl
10 optionally substituted by alkyl, nitro, cyano, halogen and lower perfluoroalkyl, multiple degrees of substitution being allowed. "Cycloalkyl" includes by way of Example cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, and the like.

As used herein, the term "cycloalkylene" refers to a non-aromatic
15 alicyclic divalent hydrocarbon radical having from three to twelve carbon atoms, optionally substituted with substituents selected from the group which includes lower alkyl, lower alkoxy, lower alkylsulfanyl, lower alkylsulfenyl, lower alkylsulfonyl, oxo, hydroxy, mercapto, amino optionally substituted by alkyl, carboxy, carbamoyl optionally substituted by alkyl, aminosulfonyl
20 optionally substituted by alkyl, nitro, cyano, halogen, lower perfluoroalkyl, multiple degrees of substitution being allowed. Examples of "cycloalkylene" as used herein include, but are not limited to, cyclopropyl-1,1-diyl, cyclopropyl-1,2-diyl, cyclobutyl-1,2-diyl, cyclopentyl-1,3-diyl, cyclohexyl-1,4-diyl, cycloheptyl-1,4-diyl, or cyclooctyl-1,5-diyl, and the like.

25 As used herein, the term "cycloalkenyl" refers to a substituted alicyclic hydrocarbon radical having from three to twelve carbon atoms and at least one carbon-carbon double bond in the ring system, optionally substituted with substituents selected from the group which includes lower alkyl, lower alkoxy, lower alkylsulfanyl, lower alkylsulfenyl, lower alkylsulfonyl, oxo,
30 hydroxy, mercapto, amino optionally substituted by alkyl, carboxy, carbamoyl

optionally substituted by alkyl, aminosulfonyl optionally substituted by alkyl, nitro, cyano, halogen and lower perfluoroalkyl, multiple degrees of substitution being allowed. Examples of "cycloalkenylene" as used herein include, but are not limited to, 1-cyclopentene-3-yl, 1-cyclohexene-3-yl, 1-
5 cycloheptene-4-yl, and the like.

As used herein, the term "cycloalkenylene" refers to a substituted alicyclic divalent hydrocarbon radical having from three to twelve carbon atoms and at least one carbon-carbon double bond in the ring system, optionally substituted with substituents selected from the group which
10 includes lower alkyl, lower alkoxy, lower alkylsulfanyl, lower alkylsulfenyl, lower alkylsulfonyl, oxo, hydroxy, mercapto, amino optionally substituted by alkyl, carboxy, carbamoyl optionally substituted by alkyl, aminosulfonyl optionally substituted by alkyl, nitro, cyano, halogen, or lower perfluoroalkyl, multiple degrees of substitution being allowed. Examples of
15 "cycloalkenylene" as used herein include, but are not limited to, 4,5-cyclopentene-1,3-diyl, 3,4-cyclohexene-1,1-diyl, and the like.

As used herein, the term "heteroatom ring system" refers to the terms heterocyclic, heterocyclyl, heteroaryl and heteroarylene. Non-limiting examples of such heteroatom ring systems are recited in the Summary of
20 the Invention, above.

As used herein, the term "heterocyclic" or the term "heterocyclyl" refers to a three to twelve-membered heterocyclic ring having one or more degrees of unsaturation containing one or more heteroatomic substitutions selected from S, SO, SO₂, O, or N, optionally substituted with substituents
25 selected from the group which includes lower alkyl, lower alkoxy, lower alkylsulfanyl, lower alkylsulfenyl, lower alkylsulfonyl, oxo, hydroxy, mercapto, amino optionally substituted by alkyl, carboxy, carbamoyl optionally substituted by alkyl, aminosulfonyl optionally substituted by alkyl, nitro, cyano, halogen, or lower perfluoroalkyl, multiple degrees of substitution
30 being allowed. Such a ring may be optionally fused to one or more of

another "heterocyclic" ring(s) or cycloalkyl ring(s). Examples of "heterocyclic" include, but are not limited to, tetrahydrofuran, pyran, 1,4-dioxane, 1,3-dioxane, piperidine, pyrrolidine, morpholine, tetrahydrothiopyran, tetrahydrothiophene, and the like.

5 As used herein, the term "heterocyclylene" refers to a three to twelve-membered heterocyclic ring diradical having one or more degrees of unsaturation containing one or more heteroatoms selected from S, SO, SO₂, O, or N, optionally substituted with substituents selected from the group which includes lower alkyl, lower alkoxy, lower alkylsulfanyl, lower
10 alkylsulfenyl, lower alkylsulfonyl, oxo, hydroxy, mercapto, amino optionally substituted by alkyl, carboxy, carbamoyl optionally substituted by alkyl, aminosulfonyl optionally substituted by alkyl, nitro, cyano, halogen and lower perfluoroalkyl, multiple degrees of substitution being allowed. Such a ring may be optionally fused to one or more benzene rings or to one or more of
15 another "heterocyclic" rings or cycloalkyl rings. Examples of "heterocyclylene" include, but are not limited to, tetrahydrofuran-2,5-diyl, morpholine-2,3-diyl, pyran-2,4-diyl, 1,4-dioxane-2,3-diyl, 1,3-dioxane-2,4-diyl, piperidine-2,4-diyl, piperidine-1,4-diyl, pyrrolidine-1,3-diyl, morpholine-2,4-diyl, and the like.

20 As used herein, the term "aryl" refers to a benzene ring or to an optionally substituted benzene ring system fused to one or more optionally substituted benzene rings to form ring systems such as anthracene, phenanthrene and naphthalene, optionally substituted with substituents selected from the group which includes lower alkyl, lower alkoxy, lower
25 alkylsulfanyl, lower alkylsulfenyl, lower alkylsulfonyl, oxo, hydroxy, mercapto, amino optionally substituted by alkyl, carboxy, tetrazolyl, carbamoyl optionally substituted by alkyl, aminosulfonyl optionally substituted by alkyl, acyl, aroyl, heteroaroyl, acyloxy, aroyloxy, heteroaroyloxy, alkoxycarbonyl, nitro, cyano, halogen, lower perfluoroalkyl, heteroaryl and aryl, multiple

degrees of substitution being allowed. Examples of aryl include, but are not limited to, phenyl, 2-naphthyl, 1-naphthyl, biphenyl, and the like.

As used herein, the term "arylene" refers to a benzene ring diradical or to a benzene ring system diradical fused to one or more optionally substituted benzene rings, optionally substituted with substituents selected from the group which includes lower alkyl, lower alkoxy, lower alkylsulfanyl, lower alkylsulfenyl, lower alkylsulfonyl, oxo, hydroxy, mercapto, amino optionally substituted by alkyl, carboxy, tetrazolyl, carbamoyl optionally substituted by alkyl, aminosulfonyl optionally substituted by alkyl, acyl, aroyl, heteroaroyl, acyloxy, aroyloxy, heteroaroyloxy, alkoxycarbonyl, nitro, cyano, halogen, lower perfluoroalkyl, heteroaryl and aryl, multiple degrees of substitution being allowed. Examples of "arylene" include, but are not limited to, benzene-1,4-diyl, naphthalene-1,8-diyl, anthracene-1,4-diyl, and the like.

As used herein, the term "heteroaryl" refers to a five - to seven - membered aromatic ring, or to a polycyclic heterocyclic aromatic ring, containing one or more nitrogen, oxygen, or sulfur heteroatoms at any position, where N-oxides and sulfur monoxides and sulfur dioxides are permissible heteroaromatic substitutions, optionally substituted with substituents selected from the group which includes lower alkyl, lower alkoxy, lower alkylsulfanyl, lower alkylsulfenyl, lower alkylsulfonyl, oxo, hydroxy, mercapto, amino optionally substituted by alkyl, carboxy, tetrazolyl, carbamoyl optionally substituted by alkyl, aminosulfonyl optionally substituted by alkyl, acyl, aroyl, heteroaroyl, acyloxy, aroyloxy, heteroaroyloxy, alkoxycarbonyl, nitro, cyano, halogen, lower perfluoroalkyl, heteroaryl, or aryl, multiple degrees of substitution being allowed. For polycyclic aromatic ring systems, one or more of the rings may contain one or more heteroatoms. Examples of "heteroaryl" used herein are furan, thiophene, pyrrole, imidazole, pyrazole, triazole, tetrazole, thiazole, oxazole, isoxazole, oxadiazole, thiadiazole, isothiazole, pyridine, pyridazine, pyrazine,

pyrimidine, quinoline, isoquinoline, benzofuran, benzothiophene, indole and indazole, and the like.

As used herein, the term "heteroarylene" refers to a five - to seven - membered aromatic ring diradical, or to a polycyclic heterocyclic aromatic ring diradical, containing one or more nitrogen, oxygen, or sulfur heteroatoms, where N-oxides and sulfur monoxides and sulfur dioxides are permissible heteroaromatic substitutions, optionally substituted with substituents selected from the group consisting of: lower alkyl, lower alkoxy, lower alkylsulfanyl, lower alkylsulfenyl, lower alkylsulfonyl, oxo, hydroxy, mercapto, amino optionally substituted by alkyl, carboxy, tetrazolyl, carbamoyl optionally substituted by alkyl, aminosulfonyl optionally substituted by alkyl, acyl, aroyl, heteroaroyl, acyloxy, aroyloxy, heteroaroyloxy, alkoxycarbonyl, nitro, cyano, halogen, lower perfluoroalkyl, heteroaryl, or aryl, multiple degrees of substitution being allowed. For polycyclic aromatic ring system diradicals, one or more of the rings may contain one or more heteroatoms. Examples of "heteroarylene" used herein are furan-2,5-diyl, thiophene-2,4-diyl, 1,3,4-oxadiazole-2,5-diyl, 1,3,4-thiadiazole-2,5-diyl, 1,3-thiazole-2,4-diyl, 1,3-thiazole-2,5-diyl, pyridine-2,4-diyl, pyridine-2,3-diyl, pyridine-2,5-diyl, pyrimidine-2,4-diyl, quinoline-2,3-diyl, and the like.

As used herein, the term "alkoxy" refers to the group R_aO- , where R_a is aliphatic.

As used herein, the term "alkylsulfanyl" refers to the group R_aS- , where R_a is aliphatic.

As used herein, the term "alkylsulfenyl" refers to the group $R_aS(O)-$, where R_a is aliphatic.

As used herein, the term "alkylsulfonyl" refers to the group R_aSO_2- , where R_a is aliphatic.

As used herein, the term "acyl" refers to the group $R_aC(O)-$, where R_a is aliphatic, cycloaliphatic, or heterocyclyl.

As used herein, the term "aroyl" refers to the group $R_aC(O)-$, where R_a is aryl.

As used herein, the term "heteroaroyl" refers to the group $R_aC(O)-$, where R_a is heteroaryl.

5 As used herein, the term "alkoxycarbonyl" refers to the group $R_aOC(O)-$, where R_a is aliphatic.

As used herein, the term "acyloxy" refers to the group $R_aC(O)O-$, where R_a is aliphatic, cycloaliphatic, or heterocyclyl.

As used herein, the term "aroyloxy" refers to the group $R_aC(O)O-$,
10 where R_a is aryl.

As used herein, the term "heteroaroyloxy" refers to the group $R_aC(O)O-$, where R_a is heteroaryl.

As used herein, the term "optionally" is inclusive of circumstances in which described condition is present and circumstances in which the
15 described condition is not present, for example, where the term is used with reference to a chemical substituent, it indicates the inclusion of embodiments in which the specified substituent is present as well as embodiments in which the specified substituent is not present.

As used herein, the term "substituted" indicates the presence of the
20 named substituent or substituents, and includes multiple degrees of substitution.

As used herein, the terms "contain" or "containing" with reference to alkyl, alkenyl, alkynyl or cycloalkyl substituents indicates in-line substitution(s) with one or more substituents at any position along the alkyl,
25 alkenyl, alkynyl or cycloalkyl substituents, such as one or more of any of O, S, SO, SO₂, N, or N-alkyl, including, for example, -CH₂-O-CH₂-, -CH₂-SO₂-CH₂-, -CH₂-NH-CH₃ and so forth.

As used herein, the term "solvate" is a complex of variable stoichiometry formed by a solute (in this invention, a compound of formula
30 (I)) and a solvent. Such solvents for the purpose of the invention may not

interfere with the biological activity of the solute. Solvents may be, by way of example, water, ethanol, or acetic acid.

The compounds of the present invention have the ability to crystallize in more than one form, a characteristic which is known as polymorphism, 5 and such polymorphic forms ("polymorphs") are within the scope of the present invention. Polymorphism generally can occur as a response to changes in temperature or pressure or both and can also result from variations in the crystallization process. Polymorphs can be distinguished by various physical characteristics known in the art such as x-ray diffraction 10 patterns, solubility, and melting point.

As used herein, the terms "biohydrolyzable carbonate", "biohydrolyzable ureide" and "biohydrolyzable carbamate" include carbonates, ureides, and carbamates, respectively, of a compound of the general formula (I) which carbonates, ureides, and carbamates, do not 15 completely diminish the biological activity of the parent substance. Such carbonates, ureides, and carbamates may confer on the parent compound of the general formula (I) advantageous properties *in vivo*, such as improved duration of action, onset of action, and the like. Also included are compounds which are relatively biologically inactive but which are converted 20 *in vivo* by the subject to the biologically active principle. An advantage of such biohydrolyzable forms is that, for example, they facilitate improved oral administration because the carbonates, ureides, and carbamates are more readily absorbed from the gut and are then transformed to a compound of formula (I) in plasma. Many examples of such biohydrolyzable compounds 25 are known in the art and include, by way of example, lower alkyl carbamates.

As used herein, the term "biohydrolyzable ester" is an ester of a compound of general formula which does not completely diminish the biological activity of the parent substance. Such esters may confer on the parent compound of the general formula (I) advantageous properties *in vivo*, 30 such as improved duration of action, onset of action, and the like. Also

included are esters which are relatively biologically inactive but which are converted in vivo by the subject to the biologically active principle. An advantage of such biohydrolyzable forms is that, for example, they facilitate improved oral administration because they are more readily absorbed from
5 the gut and are then transformed to a compound of formula (I) in plasma. Many examples of such biohydrolyzable esters are known in the art and include, by way of example, lower alkyl esters, lower acyloxy-alkyl esters, lower alkoxyacyloxyalkyl esters, alkoxyacyloxy esters, alkyl acylamino alkyl esters and choline esters.

10 As used herein, the term "biohydrolyzable amide" is an amide of a compound of general formula which does not completely diminish the biological activity of the parent substance. Such amides may confer on the parent compound of the general formula (I) advantageous properties *in vivo*, such as improved duration of action, onset of action, and the like. Also
15 included are amides which are relatively biologically inactive but which are converted in vivo by the subject to the biologically active principle. An advantage of such biohydrolyzable forms is that, for example, they facilitate improved oral administration because they are more readily absorbed from the gut and are then transformed to a compound of formula (I) in plasma.
20 Many examples of such biohydrolyzable are known in the art and include, by way of example, lower alkyl amides, α -amino acid amides, alkoxyacyl amides and alkylaminoalkylcarbonyl amides.

As used herein, the term "prodrug" includes compounds which are hydrolyzable in vivo to yield an active compound of formula (I), including for
25 example, biohydrolyzable amides, biohydrolyzable esters and biohydrolyzable carbamates. The term "prodrug" also includes compounds in which the biohydrolyzable functionality is encompassed in the compound of formula (I): for example, a lactam formed by a carboxylic group in R₁ and an amine in R₂, and compounds which may be oxidized or reduced biologically
30 at a given functional group to yield drug substances of formula (I). Examples

of such functional groups are, but are not limited to, 1,4-dihydropyridine, N-alkylcarbonyl-1,4-dihydropyridine, 1,4-cyclohexadiene, tert-butyl, and the like.

As used herein, the term "affinity reagent" means a group attached to
5 the compound of formula (I) which does not affect its in vitro biological activity, allowing the compound to bind to a target, yet such a group binds strongly to a third component allowing a) characterization of the target as to localization within a cell or other organism component, perhaps by visualization by fluorescence or radiography, or b) facile separation of the
10 target from an unknown mixture of targets, whether proteinaceous or not proteinaceous. An Example of an affinity reagent according to b) would be biotin either directly attached to (I) or linked with a spacer of one to 50 atoms selected from the group consisting of: C, H, O, N, S, or P in any combination. An Example of an affinity reagent according to a) above would be
15 fluorescein, either directly attached to (I) or linked with a spacer of one to 50 atoms selected from the group consisting of: C, H, O, N, S, or P in any combination.

The term "effective amount" means that amount of a drug or pharmaceutical agent that will elicit the biological or medical response of a
20 tissue, system, animal or human that is being sought by a researcher or clinician. The term "therapeutically effective amount" means any amount which, as compared to a corresponding subject who has not received such amount, results in improved treatment, healing, prevention, or amelioration of a disease or disorder, or a decrease in the rate of advancement of a
25 disease or disorder, and also includes amounts effective to enhance normal physiological function.

Whenever the terms "aliphatic" or "aryl" or either of their prefixes appear in a name of a substituent (e.g. arylalkoxyaryloxy) they include those characteristics given above for "aliphatic" and "aryl". Aliphatic or cycloalkyl
30 substituents are term equivalents to those having one or more degrees of

unsaturation. Designated numbers of carbon atoms (e.g. C₁₋₁₀) refer independently to the number of carbon atoms in an aliphatic or cyclic aliphatic moiety or to the aliphatic portion of a larger substituent in which the term "aliphatic" appears as a prefix (e.g. "al-").

5 As used herein, the term "disubstituted amine" or "disubstituted amino-" includes either one or two substitutions on that particular nitrogen atom.

As used herein, the term "oxo" refers to the substituent =O.

As used herein, the term "halogen" or "halo" shall include iodine,
10 bromine, chlorine and fluorine.

As used herein, the term "mercapto" refers to the substituent -SH.

As used herein, the term "carboxy" refers to the substituent -COOH.

As used herein, the term "cyano" refers to the substituent -CN.

As used herein, the term "aminosulfonyl" refer to the substituent
15 -SO₂NH₂.

As used herein, the term "carbamoyl" refers to the substituent -
C(O)NH₂.


As used herein, the term "sulfanyl" refers to the substituent -S-.


As used herein, the term "sulfenyl" refers to the substituent -S(O)-.


20 As used herein, the term "sulfonyl" refers to the substituent -S(O)₂-.


The compounds of formula (I) can be prepared readily according to the following reaction General Synthesis Schemes (in which all variables are as defined before) and Examples or modifications thereof using readily
25 available starting materials, reagents and conventional synthesis procedures. In these reactions, it is also possible to make use of variants which are themselves known to those of ordinary skill in this art, but are not mentioned in greater detail.


General Synthesis Schemes











The most preferred compounds of the invention are any or all of those specifically set forth in these examples. These compounds are not, however, to be construed as forming the only genus that is considered as
5 the invention, and any combination of the compounds or their moieties may itself form a genus. The following examples further illustrate details for the preparation of the compounds of the present invention. Those skilled in the art will readily understand that known variations of the conditions and processes of the following preparative procedures can be used to prepare
10 these compounds. All temperatures are degrees Celsius unless noted otherwise.

Abbreviations used in the Examples are as follows:

	g	= grams
	mg	= milligrams
15	L	= liters
	mL	= milliliters
	M	= molar
	N	= normal
	mM	= millimolar
20	i.v.	= intravenous
	p.o.	= per oral
	s.c.	= subcutaneous
	Hz	= hertz
	mol	= moles
25	mmol	= millimoles
	mbar	= millibar
	psi	= pounds per square inch
	rt	= room temperature
	min	= minutes
30	h	= hours
	mp	= melting point
	TLC	= thin layer chromatography
	R _f	= relative TLC mobility
	MS	= mass spectrometry
35	NMR	= nuclear magnetic resonance spectroscopy
	APCI	= atmospheric pressure chemical ionization
	ESI	= electrospray ionization
	m/z	= mass to charge ratio
	t _r	= retention time

	Pd/C	= palladium on activated carbon
	ether	= diethyl ether
	MeOH	= methanol
	EtOAc	= ethyl acetate
5	TEA	= triethylamine
	DIEA	= diisopropylethylamine
	THF	= tetrahydrofuran
	DMF	= N, N-dimethylformamide
	DMSO	= dimethylsulfoxide
10	DDQ	= 2,3-dichloro-5,6-dicyano-1,4-benzoquinone
	LAH	= lithium aluminum hydride
	TFA	= trifluoroacetic acid
	LDA	= lithium diisopropylamide
	THP	= tetrahydropyranyl
15	NMM	= N-methylmorpholine, 4-methylmorpholine
	HMPA	= hexamethylphosphoric triamide
	DMPU	= 1,3-dimethoxypropylene urea
	d	= days
	ppm	= parts per million
20	kD	= kiloDalton
	LPS	= lipopolysaccharide
	PMA	= phorbol myristate acetate
	SPA	= scintillation proximity assay
	EDTA	= ethylenediamine tetraacetic acid
25	FBS	= fetal bovine serum
	PBS	= phosphate buffered saline solution
	BrdU	= bromodeoxyuridine
	BSA	= bovine serum albumin
	FCS	= fetal calf serum
30	DMEM	= Dulbeccos modified Eagle's medium
	pfu	= plaque forming units
	MOI	= multiplicity of infection

Reagents are commercially available or are prepared according to
35 procedures in the literature. The physical data given for the compounds
exemplified is consistent with the assigned structure of those compounds. ¹H
NMR spectra were obtained on VARIAN Unity Plus NMR
spectrophotometers at 300 or 400 Mhz. Mass spectra were obtained on
Micromass Platform II mass spectrometers from Micromass Ltd. Altrincham,
40 UK, using either Atmospheric Chemical Ionization (APCI) or Electrospray
Ionization (ESI). Analytical thin layer chromatography (TLC) was used to

verify the purity of some intermediates which could not be isolated or which were too unstable for full characterisation, and to follow the progress of reactions. Unless otherwise stated, this was done using silica gel (Merck Silica Gel 60 F254). Unless otherwise stated, column chromatography for the purification of some compounds, used Merck Silica gel 60 (230-400 mesh), and the stated solvent system under pressure.

Procedure A - First method for 1H-indol-2,3-dione (isatin) formation: preparation of 6-H-1-thia-3,6-diaza-as-indacen-7,8-dione.

10 To a 1-L flask was added a magnetic stir bar, 85 g of sodium sulfate, and 100 mL of water. The mixture was magnetically stirred until all the solids were dissolved. To the resultant aqueous solution was added a solution of 6-aminobenzothiazole (4.96 g, 33.0 mmol) in 50 mL of 1N aqueous hydrochloric acid and 10 mL of ethanol. The mixture was stirred, 15 and chloral (6.0 g, (36 mmol) was added. To the resultant solution was added a solution of hydroxyl amine hydrochloride (7.50 g, 108 mmol) in 30 mL of water. The final mixture was heated with stirring to a gentle boil until all solids disappeared, and heating was continued for an additional 15 min. The flask was removed from the heat, and the solution was poured onto 500 20 g of ice. The mixture was stirred as the product precipitated from solution. The precipitate was collected by suction filtration, washed thoroughly with water, filtered, and air dried to provide 6.9 g (94%) of N-benzothiazol-6-yl-2-hydroxyimino-acetamide: ^1H NMR (DMSO- d_6): δ 12.2 (s, 1H), 10.4 (s, 1H), 9.2 (s, 1H), 8.5 (s, 1H), 7.9 (d, 1H), 7.7 (m, 1H), 7.7 (s, 1H); APCI-MS m/z 25 220 (M-H) $^-$. To a 1-L 3-neck round bottom flask was placed a magnetic stir bar and 100 ml of concentrated sulfuric acid. The flask was fitted with a thermometer to monitor the temperature of the reaction. The sulfuric acid was heated to 100°C, and 10.0 g (45.2 mmol) of N-benzothiazol-6-yl-2-hydroxyimino-acetamide was added slowly. The solution was heated for ~1

h, and the reaction mixture was poured into 750 g of ice and water. The residual reaction mixture in the reaction vessel was washed out with an additional 20 mL of cold water. The aqueous slurry was stirred for about 1 h and filtered. The solid was washed thoroughly with water, filtered, and air
5 dried to yield 4.3 g (46 %) of 6-H-1-thia-3,6-diaza-as-indacen-7,8-dione: ¹H NMR (DMSO-d₆): δ 11.1 (s, 1H), 9.2 (s, 1H), 8.2 (d, 1H), 7.0 (d, 1H); APCI-MS *m/z* 203 (M-H)⁺.

Procedure B - First method for 1,3-dihydro-indol-2-one (oxindole)
10 formation (Gassman and van Bergen, Journal of the American Chemical Society 1974, 96, 5508-12): preparation of 6,8-dihydro-1-thia-3, 6-diaza-as-indacen-7-one.

A 2-L three-neck round bottom flask was fitted with an internal thermometer, 250-mL addition funnel, magnetic stir bar and septa. The flask
15 was charged with nitrogen, 200 mL of dry THF, and 6-aminobenzothiazole (15.2 g, 0.100 mol). The mixture was stirred and cooled in a dry ice-acetone bath to an internal temperature of -74°C. A solution of tert-butyl hypochlorite (11.0 g, 0.103 mol) in 50 mL of dichloromethane was added over a 15 min period. The resultant solution was stirred for an additional 3 h at dry ice-
20 acetone bath temperature. To the reaction was then added by slow, dropwise addition a solution of ethyl methylthioacetate (13.8 g, 0.103 mol) in 50 mL of dichloromethane. The resultant solution was stirred for an additional 3 h at dry ice-acetone bath temperature. A solution of triethyl amine (25.3 g, 0.250 mol) and 50 ml of dichloromethane was added at dry
25 ice-acetone bath temperature, and the solution was stirred for 0.5 h. The cooling bath was removed, and the reaction was allowed to warm to rt. The reaction was then concentrated to a thick residue. The thick oil was resuspended in 200 mL of ether and 600 mL of 0.25 M hydrochloric acid. The mixture was allowed to stir for 24 h. The resulting solid was filtered from

the mixture and triturated with water and ether. The solid was then resuspended in cold MeOH, filtered and dried under vacuum for 16 h to yield 18.7 g (79%) of 8-methylsulfanyl-6, 8-dihydro-1-thia-3,6-diaza-as-indacen-7-one: ^1H NMR (DMSO- d_6) δ 10.8 (s, 1H), 9.2 (s, 1H), 8.0 (d, 1H), 7.1 (d, 5 1H), 1.8 (s, 3H); APCI-MS m/z 235 (M-H) $^-$. To a 500-mL erlenmeyer flask was added a stir bar, 8.1 g (0.034 moles) of 8-methylsulfanyl-6, 8-dihydro-1-thio-3, 6-diaza-as-indacen-7-one and 100 mL of glacial acetic acid. The mixture was stirred until all the starting material had dissolved. The reaction mixture was then diluted with 100 mL of THF. Zinc metal (16 g, 325 mesh) 10 was then added. The heterogeneous mixture was then stirred and heated to 60 °C for 2.5 h. The mixture was vacuum filtered through a one half inch pad of celite. The residue on the filter pad was washed with additional THF. The filtrates were combined and concentrated to a wet solid. The solid was triturated with MeOH, filtered and air dried to yield 4.51 g (70%) of 6,8- 15 dihydro-1-thia-3,6-diaza-as-indacen-7-one as a free-flowing solid: ^1H NMR (DMSO- d_6): δ 10.5 (s, 1H), 9.1 (s, 1H), 7.9 (d, 1H), 7.0 (d, 1H), 3.6 (s, 2H); APCI-MS m/z 191 (M+H) $^+$.

Procedure C - Second method for 1,3-dihydro-indol-2-one (oxindole) 20 formation (Seibert, Chemie Berichte 1947, 80, 494-502): preparation of 3-H-pyrrolo[3,2-f]quinoline-2-one via Wolff-Kishner reduction.

A solution of 2.3 g (12 mmol) of 3-H-pyrrolo[3,2-f]quinoline-1,2-dione (prepared from 6-aminoquinoline according to Procedure A) and 2.0 ml (0.06 mol) of hydrazine in 50 ml of DMF and 50 ml of ethanol was stirred at reflux 25 for 2 h. The resulting suspension was allowed to cool to ambient temperature and was then chilled in an ice bath and filtered. The solid was washed with a small volume of ethanol and allowed to air dry to give 1-hydrazono-1,3-dihydropyrrolo[3,2-f]quinolin-2-one as an orange solid (1.8 g, 73%): ^1H NMR (DMSO- d_6): δ 7.37 (d, J = 8.8 Hz, 1H), 7.47 (dd, J = 8.4, 4.2

Hz, 1H), 7.81 (d, $J = 8.8$ Hz, 1H), 8.71 (dd, $J = 4.2, 1.6$ Hz, 1H), 8.80 (d, $J = 8.4$ Hz, 1H), 9.90 (br d, $J = 14.7$ Hz, 1H), 10.89 (br d, $J = 14.7$ Hz, 1H), 10.95 (br s, 1H); ESI-MS m/z 213 ($M+H$)⁺. A solution 1.8 g (8.5 mmol) of 1-hydrazono-1,3-dihydropyrrolo[3,2-f]quinolin-2-one in 50 ml of freshly
5 prepared 0.5 M sodium ethoxide solution was stirred at reflux for 3 h. The solution was diluted with 50 ml of water, neutralized with acetic acid, and concentrated on a rotary evaporator until cloudy. The solution was stored in a refrigerator overnight. The solid was filtered off, and the filtrate was extracted with three 80-ml portions of EtOAc. A solution of the solid in
10 MeOH/EtOAc was combined with the extracts and passed through a short pad of silica gel, eluting with EtOAc. The solution was then concentrated to a small volume on a rotary evaporator, and the resulting suspension was diluted with an equal volume of ethanol, sonicated, and filtered to give 3-H-pyrrolo[3,2-f]quinoline-2-one as a light green solid (0.52 g, 33%); ¹H NMR
15 (DMSO-*d*₆): δ 3.80 (s, 2H), 7.35 (d, $J = 8.8$ Hz, 1H), 7.44 (dd, $J = 8.4, 4.2$ Hz, 1H), 7.88 (d, $J = 8.8$ Hz, 1H), 8.08 (d, $J = 8.4$ Hz, 1H), 8.70 (dd, $J = 4.2, 1.6$ Hz, 1H), 10.57 (br s, 1H); APCI-MS m/z 183 ($M-H$)⁻.

Procedure D - Third method for 1,3-dihydro-indol-2-one (oxindole)
20 formation (Quallich and Morrissey, Synthesis, 1993, 51-53): preparation of 6-bromooxindole.

Sodium hydride (60% oil dispersion, 4.00 g, 100 mmol) was added to a dry 500 ml flask under nitrogen and washed with three 25 ml portions of hexanes. Anhydrous DMSO (100 ml) was added, followed by dimethyl
25 malonate (11.4 ml, 100 mmol). The reaction was heated briefly to 100° C with stirring, then cooled to room temperature. 2,5-Dibromonitrobenzene (12.9 g, 46.0 mmol) was added and the reaction was heated at 110° C for 2 hrs. After cooling to room temperature, the solution was added in portions to 300 ml of saturated aqueous ammonium chloride with 150 ml of 1:1

hexanes/ethyl acetate. The organic layer was washed with 300 ml of saturated aqueous ammonium chloride, four 200 ml portions of water, and 200 ml of saturated aqueous sodium chloride. The organic layer was dried over magnesium sulfate and the solvent was evaporated to give 13.6 g of
5 crude dimethyl 2-(4-bromo-2-nitrophenyl)malonate as a brown oil. This material (30-40 mmol) was heated to 110 °C in 250 ml DMSO with 3.6 g (84 mmol) of lithium chloride and 750 mg (42 mmol) of water for 4.5 hrs. The reaction was cooled to room temperature and added to 300 ml of ethyl acetate with 300 ml of saturated aqueous sodium chloride. The organic
10 layer was washed with a second portion of 300 ml saturated aqueous sodium chloride, dried over magnesium sulfate, and the solvent was removed to give 11.1 g brown oil. This material was adsorbed on 40 g of silica gel and applied to a column containing another 80 g of silica gel. Elution with 0-10% ethyl acetate in hexanes gave 3.53 g (28% from 2,5-
15 dibromonitrobenzene) of methyl (4-bromo-2-nitrophenyl) acetate as a yellow solid.

This material (3.53 g, 12.8 mmol) was dissolved in ethanol (80 ml) with 50 ml of 50% sulfuric acid and heated to reflux with stirring. Zinc powder (3.40 g, 52 mmol) was added in portions over 1 hr. Heating was
20 continued for another 2 hrs and the reflux condenser was removed to allow ethanol to evaporate from the hot reaction under a stream of nitrogen. The reaction mixture was filtered through celite, washing with 100 ml of ethyl acetate. The water layer was separated from the filtrate and extracted with 100 ml of ethyl acetate. Combined ethyl acetate layers were washed with 30
25 ml of saturated aqueous sodium bicarbonate and 30 ml of saturated aqueous sodium chloride and dried over magnesium sulfate. Evaporation of solvent gave 1.6 g of crude product which was purified by chromatography on 25 g of silica gel with 10-40% ethyl acetate/hexanes to give 0.85 g (31%) of 6-bromooxindole as an off-white solid.

Procedure E - Method for dimethylaminomethinyloxindole formation: preparation of 8-dimethylamino-methylene-6,8-dihydro-1-thia-3,6-diaza-as-indacen-7-one.

To a suspension of 1.0 g (5.3 mmol) of 6,8-dihydro-1-thia-3,6-diaza-
5 as-indacen-7-one (Procedure B) in 7.5 mL of DMF was added 1.38 g (6.80 mmol) of N,N-dimethylformamide-di-t-butyl acetal. The mixture was stirred at ambient temperature for 1 h and diluted with 7.5 mL of Et₂O. The resulting precipitate was isolated filtration to afford 8-dimethylamino-methylene-6,8-dihydro-1-thia-3,6-diaza-as-indacen-7-one as a tan solid (1.0
10 g, 77%): ¹H NMR (DMSO-d₆): δ 3.33 (bs, 3H), 3.59 (bs, 3H), 6.97 (d, J = 8.4, 1H), 7.33 (s, 1H), 7.62 (d, J = 8.4, 1H), 9.13 (s, 1H), 10.29 (s, 1H); APCI-MS: *m/z* 246 (M+H)⁺.

Procedure F - Method for ethoxymethinyloxindole formation:
15 preparation of 8-ethoxymethylene-6,8-dihydro-1-thia-3,6-diaza-as-indacen-7-one.

To a 250-ml round bottom flask was added a stir bar, 6,8-dihydro-1-thia-3,6-diaza-as-indacen-7-one (Procedure B, 4.0 g, 0.021 mol), 40 mL of glacial acetic and diethoxymethyl acetate (17.0 g, 0.105 moles). The flask
20 was fitted with a reflux condensor and charged with nitrogen. The reaction was heated to reflux for 8 h. The flask was cooled, the stir bar was removed and the reaction was concentrated to a wet solid. The solid was triturated with a solution of ether and ethanol. The mixture was filtered, the solid was washed with an ethanol-ether solution, and the solid was dried under
25 vacuum to yield 8-ethoxymethylene-6,8-dihydro-1-thia-3,6-diaza-as-indacen-7-one: ¹H NMR (DMSO-d₆): δ 10.5 (s, 1H), 9.1 (s, 1H), 7.8 (d, 1H), 7.7 (s, 1H), 7.0 (d, 1H), 4.5 (q, 2H), 1.4 (t, 3H); APCI-MS *m/z* 245 (M-H)⁻.

Procedure G - Method for vinylogous urea formation: preparation of N-methyl-N-(4-((Z)-[(7-oxo-6,7-dihydro-8H-[1,3]thiazolo[5,4-e]indol-8-ylidene)methyl]amino)phenyl)acetamide (Example 58).

A mixture of 8-dimethylamino-methylene-6,8-dihydro-1-thia-3,6-diazas-indacen-7-one (Procedure E, 0.040 g, 0.163 mmol) or 8-ethoxymethylene-6,8-dihydro-1-thia-3,6-diazas-indacen-7-one (Procedure F, 0.163 mmol), 4-amino-N-methylacetanilide (0.040 g, 0.244 mmol) in absolute ethanol (5 ml) was heated with stirring at 90°C for 16h. The reaction was diluted with ethanol and diethyl ether and the product collected
10 by filtration to yield 0.038 g (64%) of the title compound. ¹H NMR (DMSO-d₆): δ 11.03 (d, 1H, J=12.3Hz), 10.84 (s, 1H), 9.23 (s, 1H), 8.02 (d, 1H, J=12.3Hz), 7.78 (d, 1H, J=8.4Hz), 7.48 (d, 2H, J=8.2Hz), 7.35 (d, 2H, J=8.2Hz), 7.09 (d, 1H, J=8.4Hz), 3.11 (s, 3H), 1.76 (s, 3H); ES-MS *m/z* 363 (M-H).

15

Procedure H - Method for condensation of a phenylhydrazine with an isatin to form hydrazones: preparation of 3,6-dihydro[1,2,3]triazolo[4,5-e]indole-7,8-dione 8-[N-(4-methoxyphenyl)hydrazone] (Example 89).

3,6-Dihydro[1,2,3]triazolo[4,5-e]indole-7,8-dione was prepared from 5-aminobenzotriazole according to Procedure A in 6% yield: ¹H NMR (DMSO-d₆): δ 7.04 (d, J = 8.4 Hz, 1H), 7.97 (d, J = 2.2 Hz, 1H), 8.01 (dd, J = 2.2, 8.4 Hz, 1H), 8.20 (s, 1H), 9.26 (s, 1H), 11.19 (bs, 1H); APCI-MS *m/z* 215 (M+1)⁺. 3,6-Dihydro[1,2,3]triazolo[4,5-e]indole-7,8-dione (47 mg, 0.25 mmol) was combined with 4-methoxyphenylhydrazine hydrochloride (52 mg, 0.3
25 mmol) in 2 ml of ethanol and heated at 70°C for 3 hrs. The product was collected by filtration of the hot solution, washing with ethanol and diethyl ether, to give 39 mg (50%) of the title compound as a dark red solid. NMR showed ~1:1 Z/E mixture. ¹H NMR (DMSO-d₆): δ 3.80 (s, 3H); 7.0 (m, 3H); 7.24 (d, J = 8.7 Hz, 0.5H); 7.41 (d, J = 8.8 Hz, 0.5H); 7.78 (m, 1.5H); 7.98 (d,

J = 8.5 Hz, 0.5H); 10.8 (s, 0.5H); 11.3 (s, 0.5H); 12.85 (s, 0.5H); 12.95 (s, 0.5H). APCI-MS *m/z* 307 (M-1).

Procedure I - Method for palladium catalyzed coupling of 6-bromooxindole
5 with alkenyl and aromatic tin reagents: preparation of 6-vinyl oxindole.

To a mixture of 6-bromooxindole (Procedure D, 0.50 g, 2.4 mmol), vinyltributylstannane (0.95g, 3.0mmol), lithium chloride (.03g, 7.1mmol), 2,6-di-*tert*-butyl-4-methylphenol (0.01g, 0.05mmol) in acetonitrile (25ml) stirring at 80°C was added dichlorobis(triphenylphosphine)palladium (II). The
10 resulting reaction was stirred with heating for 16h. The reaction was poured into a vigorously stirring mixture of 5M potassium fluoride solution: ethyl acetate/ 1:1 (250mL) and stirred for 0.75h. The resulting biphasic mixture was filtered through a Celite 521 pad and the pad flushed with ethyl acetate (5x200mL). The combined organic phases were washed with water
15 (200mL), saturated sodium chloride (200mL) and filtered through Whatman PS 1 paper and evaporated *in vacuo* to a golden yellow syrup. The syrup was titrated with diethyl ether to yield several crops of tan solid. Pure samples were combined, slurried with diethyl ether, filtered, and air dried to yield 0.12g (31%) of 6-vinyloxindole: ¹H NMR (DMSO-d₆): δ 10.36 (s,1H),
20 7.13 (d,1H, J=7.7Hz), 6.98 (d,1H, J=7.5Hz), 6.66 (dd,1H, J=10.9,17.7Hz), 5.70 (d,1H, J=17.6Hz), 5.18 (d,1H, J=10.9Hz), 3.42 (s,2H).

Example 1:

6-Bromo-3-[(Z and E)-[4-(4-morpholinyl)anilino]methylidene]-1,3-dihydro-2H-indol-2-one.

25 Prepared from the dimethylaminomethylene derivative (Procedure E) of 6-bromo-oxindole (procedure D) and 4-(4-morpholino)aniline according to Procedure G in 87% yield to give ~4:1 isomer mixture. Principal isomer: ¹H NMR (DMSO-d₆): δ 10.67 (d, 1H, J=12.5Hz), 10.56(s,1H), 8.57 (d,1H, J=12.8Hz), 7.48 (d,1H, J=7.9Hz), 7.35 - 7.29 (m,2H), 7.11 - 7.04 (m, 3H),
30 6.93 (s,1H), 3.76 (s,4H), 3.15 (s, 4H); ES-MS *m/z* 400, 402 (M + 1).

Example 2:**6-Bromo-3-[(Z and E)-[4-(4-pyridinylmethyl)anilino]methylidene]-1,3-dihydro-2H-indol-2-one.**

Prepared according to Example 1 in 26% yield to give ~9:1 isomer
5 mixture. Principal isomer: ^1H NMR (DMSO- d_6): δ 10.72 (d, 1H, $J=12.6\text{Hz}$), 10.65 (s, 1H), 8.66 (d, 1H, $J=12.8\text{Hz}$), 8.48 (d, 1H, $J=5.9\text{Hz}$), 7.55 (d, 1H, 8.1Hz), 7.40 - 7.27 (m, 6H), 7.12 (dd, 1H, $J=1.8, 8.1\text{Hz}$), 6.99 (d, 1H, $J=1.8\text{Hz}$), 3.97 (s, 2H); APCI-MS m/s 404, 406 (M-1).

10 Example 3:**8-[(Z and E)-4-Toluidinomethylidene]-6,8-dihydroimidazo[4,5-e]indol-7(3H)-one.**

Sodium hydride (7.2 g, 60% in mineral oil, 0.18 mol) was added in portions over 30 min to a stirred solution of 5-aminobenzimidazole (15.0 g, 0.105 mol) in dry DMF (100 mL) and the mixture was stirred at room
15 temperature for 30 min. The reaction mixture was cooled to 0° C and a solution of di-tert-butyl dicarbonate (24.0 g, 0.110 mol) in dry DMF (25 mL) was added over 10 min. The reaction mixture was stirred at room temperature for 18 h. The solvent was evaporated under reduced pressure and the residue was partitioned between water (200 mL) and diethyl ether
20 (200 mL). The organic phase was separated and the aqueous layer was extracted with diethyl ether. The combined organic extracts were washed with water and brine and dried over anhydrous magnesium sulfate. The drying agent was removed by filtration through a pad of silica gel and the filtrate evaporated to give a mixture of 5- and 6-amino-1-tert-
25 butoxycarbonylbenzimidazole, 21.5 g (84%). ^1H NMR (DMSO- d_6): δ 8.40 (s, 1H), 8.23 (s, 1H), 7.55 (d, 1H, $J = 8.8\text{ Hz}$), 7.34 (d, 1H, $J = 8.4\text{ Hz}$), 7.13 (d, 1H, $J = 1.6\text{ Hz}$), 6.83 (d, 1H, $J = 2\text{ Hz}$), 6.68 (dd, 1H, $J = 8.8, 1.6\text{ Hz}$), 6.60 (dd, 1H, $J = 8.4, 2\text{ Hz}$), 1.61 (s, 9H), 1.60 (s, 9H).

To a cold (-78° C) solution of ethyl methylthioacetate (11.87 g, 88
30 mmol) in dry dichloromethane (300 mL) was added, dropwise, sulfuric

- chloride (7.1 mL, 11.93 g, 88 mmol) over 5 min. The solution was stirred for 30 min and then a solution of a mixture of 5- and 6-amino-1-*tert*-butoxycarbonylbenzimidazole (21.5 g, 88 mmol) and Proton Sponge (18.9 g, 88 mmol) in dry dichloromethane (150 mL) was added dropwise over 30 min.
- 5 The mixture was stirred at $\approx 78^{\circ}\text{C}$ for about 1 h and then triethylamine (12.3 mL, 8.94 g, 88 mmol) was added dropwise and the mixture was allowed to warm to room temperature over 18 h. The reaction mixture was washed with water (3 X 100 mL) and brine and then the organic phase was dried over anhydrous magnesium sulfate. The drying agent was removed by filtration
- 10 through a pad of silica gel and the filtrate evaporated. The residue was triturated with a small amount of diethyl ether and the resulting solid was collected by filtration. The filtrate was stirred with 2N aqueous HCl (10 mL) for 18 h to yield further white solid which was collected by filtration. The combined white solids were dried under vacuum to give a mixture of 1 and 3-
- 15 *tert*-butoxycarbonyl-8-methylthio-7-oxo-7,8-dihydropyrrolo[2,3-g]benzimidazole, 8.74 g (30%). ^1H NMR ($\text{DMSO}-d_6$): δ 10.882 (s, 1H), 10.72 (s, 1H), 9.41 (s, 1H), 8.52 (s, 1H), 7.45 (d, 1H, $J = 8.4\text{ Hz}$), 7.65 (d, 1H, $J = 8.4\text{ Hz}$), 7.11 (d, 1H, $J = 8.4\text{ Hz}$), 7.93 (d, 1H, $J = 8.4\text{ Hz}$), 4.84 (s, 1H), 4.67 (s, 1H), 1.93 (s, 3H), 1.81 (s, 3H), 1.66 (s, 9H).
- 20 A solution of a mixture of 1 and 3-*tert*-butoxycarbonyl-8-methylthio-7-oxo-7,8-dihydropyrrolo[2,3-g]benzimidazole (2.0 g, 6 mmol) in THF (50 mL) was stirred at room temperature and a saturated aqueous solution of ammonium chloride (50 mL) was added. Activated zinc dust (8.0 g) was added and the resulting mixture was stirred vigorously for about 18 h. The
- 25 solids were removed by filtration through a pad of Celite and the organic layer was separated. The organic phase was dried over anhydrous magnesium sulfate, filtered and the solvent evaporated under reduced pressure to leave a yellow solid. Trituration with a small amount of solid gave a pale solid which was collected by filtration and dried under vacuum to
- 30 give a mixture of 1 and 3-*tert*-butoxycarbonyl-7-oxo-7,8-dihydropyrrolo[2,3-

g]benzimidazole, 1.32 g (77%). ^1H NMR ($\text{DMSO}-d_6$): δ 10.51 (br s, 1H), 10.21 (br s, 1H), 8.46 (s, 1H), 8.02 (s, 1H), 7.67 (m, 1H), 7.55 (d, 1H), 6.87 (d, 1H), 6.70 (m, 1H), 3.84 (s, 2H), 1.62 (s, 9H).

Dimethylformamide di-*tert*-butyl acetal (0.34 mL, 0.29 g, 1.4 mmol) 5 was added dropwise to a stirred solution of a mixture of 1 and 3-*tert*-butoxycarbonyl-7-oxo-7,8-dihydropyrrolo[2,3-*g*]benzimidazole (0.2 g, 0.7 mmol) in dry DMF (2 mL) and the mixture was stirred at room temperature for about 6 h. The solvent was evaporated under reduced pressure. The residue was dissolved in chloroform:methanol (10:1, 10 mL) and passed 10 through a pad of silica gel. The filtrate was evaporated to give a mixture of 1 and 3-*tert*-butoxycarbonyl-8-(dimethylaminomethylidiny)-7-oxo-7H-pyrrolo[2,3-*g*]benzimidazole as a brown solid, 0.042 g (17%). ^1H NMR ($\text{DMSO}-d_6$): δ 10.10 (s, 1H), 8.40 (s, 1H), 7.40 (s, 1H), 7.22 (d, 1H, $J = 8.4$ Hz), 6.82 (d, 1H, $J = 8.4$ Hz), 1.64 (s, 9H).

15 A solution of a mixture of 1 and 3-*tert*-butoxycarbonyl-8-(dimethylaminomethylidiny)-7-oxo-7H-pyrrolo[2,3-*g*]benzimidazole (0.02g, 0.061 mmol) and *p*-toluidine (0.01 g, 0.089 mmol) in acetic acid (1 mL) was heated at 120° C for 3 h. The solvent was evaporated under vacuum and the residue was purified using silica gel chromatography with 20 chloroform:methanol (10:1) as eluent to afford the title compound as a mixture of *E/Z* isomers, 0.012g (68%). ^1H NMR of principal isomer ($\text{DMSO}-d_6$): δ 12.57 (s, 1H), 12.38 (d, 1H, $J = 12.8$ Hz), 10.01 (s, 1H), 8.33 (s, 1H), 7.95 (d, 1H, $J = 12.8$ Hz), 7.24 (m, 5H), 6.80 (d, 1H, $J = 8$ Hz), 2.28 (s, 3H). MS (AP^+) 289 (100) ($\text{M}^+ - \text{H}$).

25

Example 4: 8-[(*Z* and *E*)-(3-Methyl-4-nitroanilino)methylidene]-3,6-dihydro[1,2,3]triazolo[4,5-*e*]indol-7-one

Prepared as a 1:1 mixture of *Z* and *E* isomers in 95% yield from 8-dimethylaminomethylene-1-*tert*-butyloxycarbonyl-1,6-

dihydro[1,2,3]triazolo[4,5-e]indol-7-one and 3-methyl-4-nitroaniline according to the procedure of Example 5. ¹H NMR (DMSO-d₆): δ 2.65 (s, 6H); 7.07 (d, J = 8.5 Hz, 0.5 H); 7.18 (d, J = 8.5 Hz, 0.5H); 7.44 (dd, J = 9.0 Hz and 1.5 Hz, 0.5H); 7.53 (m, 1H); 7.6 (m, 1H); 7.78 (brd d, J = 8.2 Hz, 0.5H); 8.17 (d, J = 9.0 Hz, 0.5H); 8.19 (d, J = 9.5 Hz, 0.5H); 8.29 (brd d, J = 12.6 Hz, 0.5H); 8.70 (d, J = 12.1 Hz, 0.5 H); 10.49 (s, 0.5H); 11.0 (s, 0.5H); 11.16 (d, J = 12 Hz, 0.5H); 11.8 (brd d, J ~ 8 Hz, 0.5H).

APCI-MS: *m/z* 335 (M-H)⁺.

10 Example 5: 8-[(E/Z)-[4-nitro-3-(trifluoromethyl)anilino]methylidene]-1,6-dihydro[1,2,3]triazolo[4,5-e]indol-7-one

5-Aminobenzotriazole (Lancaster Chemical, 10.14 g, 75 mmol) was dissolved in 200 ml of anhydrous DMF under nitrogen and 3.00 g (75 mmol) of sodium hydride (60% oil dispersion) was added in one portion. Hydrogen evolution and mild exothermicity was observed. The reaction was stirred at room temperature for 20 minutes and then cooled in an ice bath. A solution of di-tert-butylidicarbonate (16.4 g, 75 mmol) in 100 ml of anhydrous DMF was added via siphon. Stirring was continued for 2 hrs at ice bath temperature. The solvent was removed by rotary evaporation under high vacuum at 50 °C to give 32 g of viscous liquid. The crude product was dissolved in a minimum volume of chloroform and filtered through a short column of 600 ml silica gel, eluting with 10% methanol in chloroform. The collected product was evaporated to dryness, redissolved in 400 ml of diethyl ether, and washed three times with water and once with saturated sodium chloride solution. The ether solution was dried over magnesium sulfate and the solvent was removed to give 17.7 g of a mixture of 1- and 3-tert-butyloxycarbonyl-5-aminobenzotriazole contaminated with approx. 1 g of residual mineral oil. This material was then cyclized to the corresponding 3-methylthio-oxindole by the method of Procedure (Gassman). The resultant

product (9.6 g of gray solid) was shown to be partially deprotected by NMR. This material was dissolved 200 ml of THF and treated with 50 g of zinc dust (activated by stirring for 10 min in 150 ml of 1 M HCl, followed by washing with three 100 ml portions of water). Saturated aqueous ammonium chloride
5 (150 ml) was added and the reaction was stirred overnight at room temperature. The solution was filtered through Celite, washing with THF and ethyl acetate to give 4.0 g of gray solid which was primarily 1-tert-butyloxycarbonyl-1,6-dihydro[1,2,3]triazolo[4,5-e]indol-7-one. This material (2.04 g, 7.4 mmol) was suspended in 10 ml of anhydrous DMF under
10 nitrogen, cooled in an ice bath, and treated with 4.0 ml (3.4 g, 2.2 equiv) of dimethylformamide di-tert-butyl acetal. The reaction was allowed to warm to room temperature and was stirred overnight. The solvent was removed by rotary evaporation under high vacuum. The residue was filtered through a short column of 100 ml silica gel with 30% ethanol in dichloromethane.
15 Evaporation of solvent provided 1.74 g of yellow solid which was primarily 8-dimethylaminomethylene-1-tert-butyloxycarbonyl-1,6-dihydro[1,2,3]triazolo[4,5-e]indol-7-one containing some product lacking the tert-butyloxycarbonyl protecting group.

The dimethylamino-oxindole derivative from above (33 mg, ~ 0.1
20 mmol) was combined in a reaction vial with 4-nitro-3-trifluoromethylaniline (25 mg, 0.12 mmol) in 1-2 ml of glacial acetic acid and stirred overnight in an oil bath at 110 °C. The residue was triturated with 2 ml of ethanol, heating briefly to reflux. After cooling to room temperature, 2-3 ml of diethyl ether was added and the resulting precipitate was collected by filtration to give 7.8
25 mg (20%) of the title compound as a brown solid that was shown by NMR to be ~1:1 mixture of E/Z isomers. ¹H NMR (DMSO-d₆): δ 7.07 (d, J = 9 Hz, 0.5 H); 7.20 (d, J = 8 Hz, 0.5 H); 7.63 (d, J = 8 Hz, 0.5 H); 7.8 - 8.0 (m, 2H); 8.13 (s, 0.5 H); 8.25 - 8.42 (m, 1.5 H); 8.67 (d, J = 11 Hz, 0.5 H); 10.53 (s,

0.5 H); 11.0 (s, 0.5 H); 11.24 (d, J = 11 Hz); 12.0 (brd d, J ~ 9 Hz, 0.5 H).

APCI-MS: m/z 389 (M-H)⁻.

Example 6: 8-[(Z and E)-(3-Chloro-4-nitroanilino)methylidene]-1,6-dihydro[1,2,3]triazolo[4,5-e]indol-7-one.

Prepared as a ~1:1 mixture of Z and E isomers in 70% yield from 8-dimethylaminomethylene-1-tert-butyloxycarbonyl-1,6-dihydro[1,2,3]triazolo[4,5-e]indol-7-one and 3-chloro-4-nitroaniline according to the procedure of Example 5. ¹H NMR (DMSO-d₆): δ 7.06 (d, J = 8.8 Hz, 0.5H); 7.20 (d, J = 8.5 Hz, 0.5H); 7.3 - 7.66 (m, 1.5H); 7.80 (brd m, 0.5H); 7.85 (d, J = 2.5 Hz, 0.5H); 7.91 (d, J = 2.5 Hz, 0.5H); 8.23 (d, J = 9.1 Hz, 1H); 8.32 (brd m, 0.5H); 8.63 (d, J = 11.7 Hz, 0.5H); 10.5 (s 0.5H); 11.0 (s, 0.5H); 11.1 (d, J = 11.7 Hz, 0.5H); 11.8 (brd, 0.5H). APCI-MS: m/z 355 (M-H)⁻.

15

Example 7: 8-[(Z and E)-(3,5-Dimethyl-4-nitroanilino)methylidene]-1,6-dihydro[1,2,3]triazolo[4,5-e]indol-7-one.

Prepared as a ~1:1 mixture of Z and E isomers in 74% yield from 8-dimethylaminomethylene-1-tert-butyloxycarbonyl-1,6-dihydro[1,2,3]triazolo[4,5-e]indol-7-one and 3,5-dimethyl-4-nitroaniline according to the procedure of Example 5. ¹H NMR (DMSO-d₆): δ 2.36 (s, 6H); 7.07 (d, J = 8.5 Hz, 0.5H); 7.17 (d, J = 8.5 Hz, 0.5H); 7.37 (s, 2H); 7.57 (d, J = 8.5 Hz, 0.5H); 7.76 (brd d, J = 8.5 Hz, 0.5H); 8.24 (brd d, J = 12 Hz, 0.5H); 8.69 (brd d, J = 12 Hz, 0.5H); 10.44 (s, 0.5H); 10.97 (s, 0.5H); 11.07 (brd d, J = 12 Hz, 0.5H); 11.6 (brd d, J = 12 Hz, 0.5H). APCI-MS: m/z 349 (M-H)⁻.

25

Example 8: 8-((Z and E)-{4-[2,2,2-Trifluoro-1-hydroxy-1-(trifluoromethyl)ethyl]anilino}methylidene)-1,6-dihydro[1,2,3]triazolo[4,5-e]indol-7-one.

The title compound was prepared as a ~1:1 mixture of Z and E isomers in 54% yield from 8-dimethylaminomethylene-1-tert-butyloxycarbonyl-1,6-dihydro[1,2,3]triazolo[4,5-e]indol-7-one and 4-[2,2,2-trifluoro-1-hydroxy-1-(trifluoromethyl)ethyl]aniline according to the procedure of Example 5. ¹H NMR (DMSO-d₆): δ 7.07 (d, J= 8.7 Hz, 0.5H); 7.17 (d, J= 8.7 Hz, 0.5H); 7.56 (d, J= 8.7 Hz, 0.5H); 7.6 and 7.75 (2 overlapping Abq, 4H); 8.22 (d, J= 13 Hz, 0.5H); 8.75 (m, 1H); 10.4 (s, 0.5H); 10.9 (s, 0.5H); 11.08 (d, J= 13 Hz, 0.5H); 11.7 (brd d, J= 13 Hz, 0.5H). APCI-MS: m/z 442 (M-H)⁺.

Example 9: 8-((Z and E)-[4-(2-Hydroxyethyl)anilino]methylidene)-1,6-dihydro[1,2,3]triazolo[4,5-e]indol-7-one.

The acetate ester of the title compound was obtained as a ~1:1 mixture of Z and E isomers in 71% yield (26 mg) from 8-dimethylaminomethylene-1-tert-butyloxycarbonyl-1,6-dihydro[1,2,3]triazolo[4,5-e]indol-7-one (25 mg) and 4-(2-hydroxyethyl)aniline (17 mg) in acetic acid according to the procedure of Example 5. The product was stirred in 4 ml methanol with 0.2 ml of 2M NaOH overnight at room temperature. The reaction solution was acidified with 0.2 ml of 2M sulfuric acid, and ethanol and diethyl ether were added. The resulting precipitate was collected by filtration and stirred in 3 ml of water for several hours to dissolve Na₂SO₄, then filtered and dried to give the title compound (16 mg dark green solid, 70%). ¹H NMR (DMSO-d₆): δ 2.69 (t, J~ 6 Hz, 2H); 3.58 (m, 2H); 4.62 (m, 1H); 7.01 (d, J = 8.5 Hz, 0.5H); 7.12 (d, J = 8.5 Hz, 0.5H); 7.26 and 7.32 (overlapping Abq, 4H); 7.46 (d, J = 8.5 Hz, 0.5H); 7.64 (brd,

0.5H); 8.13 (d, J = 13 Hz, 0.5H); 10.3 (s, 0.5H); 10.8 (s, 0.5H); 11.0 (d, J = 13 Hz, 0.5H); 11.45 (brd, 0.5H). APCI-MS: m/z 320 (M-H)⁺.

Example 10: **8-[(Z)-[4-(Methylsulfanyl)anilino]methylidene]-6H-5 [1,3]thiazolo[5,4-e]indol-7-one.**

This compound was prepared with 4-methylthioaniline according to Procedure G on a 0.5 mmol scale. The yield was 0.089 grams (53%) for C₁₇H₁₃N₃O₁S₂.

¹H NMR (400 MHz, DMSO-d₆) δ: 10.9(d, 1H), 10.78(s, 1H), 9.21(s, 1H),
10 7.9(d, 1H), 7.7(d, 1H), 7.25(d, 2H), 7.1(d, 1H), 6.8(d, 2H), 2.4(s, 3H).
ESI-MS m/z 338(M-1).

Example 11: **8-[(Z)-(3,5-Dimethoxyanilino)methylidene]-6H-[1,3]thiazolo[5,4-e]indol-7-one.**

15 This compound was prepared 3,5-dimethoxyaniline according to Procedure G on a 0.5 mmol scale. The yield was 0.076 grams (43%) for C₁₈H₁₅N₃O₃S₁.

¹H NMR (400 MHz, DMSO-d₆) δ: 10.9(d, 1H), 10.83(s, 1H), 9.21(s, 1H),
7.98(d, 1H), 7.77(d, 1H), 7.08(d, 1H), 6.59(s, 2H), 6.25(s, 1H), 3.77(s, 6H).
20 ESI-MS m/z 352(M-1).

Example 12: 8-[(Z)-(4-Hydroxyanilino)methylidene]-6H-[1,3]thiazolo[5,4-e]indol-7-one.

This compound was prepared with 4-hydroxyaniline according to
25 Procedure G on a 0.5 mmol scale. The yield was 0.081 grams (53%) for C₁₆H₁₁N₃O₂S₁.

¹H NMR (400 MHz, DMSO-d₆) δ: 10.95(d, 1H), 10.75(s, 1H), 9.42(s, 1H),
9.2(s, 1H), 7.91(d, 1H), 7.72(d, 1H), 7.23(d, 2H), 7.07(d, 1H), 6.79(d, 2H).
ESI-MS m/z 308(M-1).

Example 13: 1-[(Z)-(3-Methoxyanilino)methylidene]-1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one.

Prepared from the dimethylaminomethylene derivative (Procedure E) of 1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one (Procedure C) with 3-methoxyaniline according to Procedure G to give the title compound in 63% yield. ¹H NMR (DMSO-d₆) δ: 3.80 (s, 3H); 6.68 (dd, J = 1.8, 8.2 Hz, 1H); 7.06 (brd d, J = 8.0 Hz, 1H); 7.11 (brd s, 1H); 7.29 (t, J = 8.1 Hz, 1H); 7.40 (d, J = 8.8 Hz, 1H); 7.46 (dd, J = 8.5, 4.1 Hz, 1H); 7.71 (d, J = 8.8 Hz, 1H); 8.70 (d, J = 4.1 Hz, 1H); 8.8 (2 overlapping d, J = 8.5, 12 Hz, 2H); 10.94 (s, 1H); 11.74 (d, J = 12 Hz, 1H). ES-MS: *m/z* 316 (M-H)⁻.

Example 14: 3-[[[(Z)-(2-Oxo-2,3-dihydro-1H-pyrrolo[3,2-f]quinolin-1-ylidene)methyl]amino]benzonitrile.

Prepared from the dimethylaminomethylene derivative (Procedure E) of 1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one (Procedure C) with 3-aminobenzonitrile according to Procedure G in acetic acid to give the title compound in 78% yield as the acetate salt. ¹H NMR (DMSO-d₆) δ: 1.88 (s, 3H); 7.39 (d, J = 8.5 Hz, 1H); 7.45-7.6 (m, 3H); 7.78 (m, 2H); 8.20 (s, 1H); 8.75 (d, J = 4 Hz, 1H); 8.8-8.9 (m, 2H); 11.0 (s, 1H); 11.8 (d, J = 12 Hz, 1H); 11.95 (s, 1H). ES-MS: *m/z* 313 (M+H)⁺.

Example 15: 1-[(Z)-4-Toluidinomethylidene]-1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one.

Prepared from the dimethylaminomethylene derivative (Procedure E) of 1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one (Procedure C) with p-toluidine according to Procedure G to give the title compound in 98% yield. ¹H NMR (DMSO-d₆) δ: 2.25 (s, 3H); 7.2 (m, 2H); 7.35-7.5 (m, 4H); 7.70 (d, J = 8.5 Hz,

1H); 8.70 (d, J = 4.0 Hz, 1H); 7.8 (m, 2H); 10.9 (s, 1H); 11.8 (d, J = 12 Hz, 1H). ES-MS: m/z 302 (M+H)⁺.

Example 16: 1-[(Z)-(4-Methoxyanilino)methylidene]-1,3-dihydro-2H-5 pyrrolo[3,2-f]quinolin-2-one .

Prepared from the dimethylaminomethylene derivative (Procedure E) of 1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one (Procedure C) with 4-methoxyaniline according to Procedure G in acetic acid to give the title compound in 54% yield as the acetate salt. ¹H NMR (DMSO-d₆) δ: 1.88 (s, 3H); 3.75 (s, 3H); 6.97 (d, J = 8.9 Hz, 2H); 7.39 (d, J = 8.7 Hz, 1H); 7.43-7.5 (m, 3H); 7.68 (d, J = 8.5 Hz, 1H); 8.70 (d, J = 4.0 Hz, 1H); 8.75-8.82 (m, 2H); 10.9 (s, 1H); 11.75 (d, J = 12 Hz, 1H); 11.9 (s, 1H). ES-MS: m/z 318 (M+H)⁺.

Example 17: 3-({[(Z and E)-7-Oxo-6,7-dihydro[1,2,3]triazolo[4,5-e]indol-8(1H)-ylidene)methyl]amino)benzonitrile.

Prepared as a ~3:2 mixture of geometrical isomers in 88% yield from 8-dimethylaminomethylene-1-tert-butyloxycarbonyl-1,6-dihydro[1,2,3]triazolo[4,5-e]indol-7-one and 3-aminobenzonitrile according to the procedure of Example 5. ¹H NMR (DMSO-d₆): δ 6.88 and 7.02 (2 d, 3:2 ratio, J = 8.7 Hz, 1H); 7.45-7.6 (m, ~2.6H); 7.65-7.8 (m, ~1.4H); 7.90 and 7.95 (2 s, 1H); 8.15 and 8.83 (2 s, 1H); 10.2 and 10.85 (2 s, 1H). ES-MS: m/z 303 (M+H)⁺.

Example 18: 8-[(Z and E)-4-Toluidinomethylidene]-1,6-dihydro[1,2,3]triazolo[4,5-e]indol-7-one.

Prepared as a ~1:1 mixture Z and E isomers in 78% yield from 8-dimethylaminomethylene-1-tert-butyloxycarbonyl-1,6-dihydro[1,2,3]triazolo[4,5-e]indol-7-one and p-toluidine according to the

procedure of Example 5. ^1H NMR (DMSO- d_6): δ 2.28 (s, 3H); 7.01 and 7.11 (2 d, 1:1 ratio, $J = 8.4$ Hz, 1H); 7.22 and 7.32 (2 overlapping ABq, 4H); 7.45 and 7.64 (2 d, 1:1 ratio, $J = 8.4$ Hz, 1H); 8.11 and 8.68 (2 d, 1:1 ratio, $J = 13$ Hz, 1H); 10.3 and 10.8 (2 s, 1:1 ratio, 1H); 11.0 and 11.45 (2 d, 1:1 ratio, $J = 13$ Hz, 1H). ES-MS: m/z 292 ($M+H$) $^+$.

Example 19: 8-[(Z and E)-(4-Methoxyanilino)methylidene]-1,6-dihydro[1,2,3]triazolo[4,5-e]indol-7-one.

Prepared as a ~1:1 mixture Z and E isomers in 88% yield from 8-dimethylaminomethylene-1-tert-butyloxycarbonyl-1,6-dihydro[1,2,3]triazolo[4,5-e]indol-7-one and 4-methoxyaniline according to the procedure of Example 5. ^1H NMR (DMSO- d_6): δ 3.78 (s, 3H); 6.95-7.1 (m, 3H); 7.3-7.45 (m, 2.5H); 7.60 (d, $J = 8.5$ Hz, 0.5H); 8.05 and 8.65 (2 d, $J = 12$ Hz, 1H), 10.2 and 10.75 (2 s, 1H); 10.95 and 11.55 (2 brd d, $J = 12$ Hz, 1H). ES-MS: m/z 308 ($M+H$) $^+$.

Example 20: 8-[(Z)-[4-(4-Morpholinyl)anilino]methylidene]-6H-[1,3]thiazolo[5,4-e]indol-7-one.

Prepared according to Procedure G with 4-(4-morpholino)aniline in 78% yield. ^1H NMR (DMSO- d_6): δ 10.96 (d, 1H, $J=12.6$ Hz), 10.76 (s, 1H), 9.20 (s, 1H), 7.95 (d, 1H, $J=12.5$ Hz), 7.74 (d, 1H, $J= 8.4$ Hz), 7.31 (d, 2H, $J=8.8$ Hz), 7.08 (d, 1H, $J=8.4$ Hz), 6.99 (d, 2H, $J=9.0$ Hz), 3.72 (t, 4H, $J=4.7$ Hz), 3.07 (t, 4H, $J=4.8$ Hz); ES-MS m/z 379 ($M+H$).

Example 21: N-(4-[(Z)-(7-Oxo-6,7-dihydro-8H-[1,3]thiazolo[5,4-e]indol-8-ylidene)methyl]amino}phenyl)acetamide.

Prepared according to Procedure G with N-(4-aminophenyl)acetamide in 87% yield. ^1H NMR (DMSO- d_6): δ 11.01 (d, 1H, $J=12.5$ Hz), 10.80 (s, 1H), 9.95 (s, 1H), 9.22 (s, 1H), 7.99 (d, 1H, $J=12.5$ Hz), 7.75 (d, 1H, $J=8.4$ Hz),

7.60(d, 2H, J=8.8Hz), 7.35 (d, 2H, J=8.8Hz), 7.09 (d, 1H, J=8.4Hz), 2.01(s, 3H).

Example 22: 8-[(Z)-[4-(2-Hydroxyethyl)anilino]methylidene}-6H-5 [1,3]thiazolo[5,4-e]indol-7-one.

Prepared according to Procedure G with 4-(2-hydroxyethyl)aniline in 41% yield. ¹H NMR (DMSO-d₆): δ 10.98 (d, 1H, J=12.5Hz), 10.81 (s, 1H), 9.22 (s, 1H), 8.00 (d, 1H, J=12.5Hz), 7.76 (d, 1H, J=8.4Hz), 7.31 (d, 2H, J=8.4Hz), 7.24 (d, 2H, J=8.4Hz), 7.09 (d, 1H, J=8.4Hz), 4.62 (t, 1H, J=5.1Hz), 3.59 - 3.54 (m, 2H), 2.69 (t, 2H, J= 7.0Hz); ES-MS *m/z* 338 (M+H).

Example 23: 8-((Z)-[4-(4-Pyridinylmethyl)anilino]methylenidene)-6H-[1,3]thiazolo[5,4-e]indol-7-one.

Prepared according to Procedure G with 4-(4-pyridylmethyl)aniline in
 15 66% yield. ^1H NMR (DMSO- d_6): δ 10.99 (d, 1H, $J=12.5\text{Hz}$), 10.84 (s, 1H),
 9.23 (s, 1H), 8.74 (d, 2H, $J=6.4\text{Hz}$), 8.00 (d, 1H, $J=12.3\text{Hz}$), 7.82 (d, 2H,
 $J=6.2\text{Hz}$), 7.77 (d, 1H, $J=8.4\text{Hz}$), 7.40 (d, 2H, $J=8.6\text{Hz}$), 7.35 (d, 2H, $J=8.4\text{Hz}$)
 7.09 (s, 1H), 4.20 (s, 2H); ES-MS m/z 384 (M+H).

20 Example 24: 4-[[*(Z)*-(7-Oxo-6,7-dihydro-8H-[1,3]thiazolo[5,4-*e*]indol-8-ylidene)methyl]amino}benzamide.

Prepared according to Procedure G with 4-aminobenzamide in 73% yield. ¹H NMR (DMSO-d₆): δ 11.14 (d, 1H, J=12.3Hz), 10.90 (s, 1H), 9.27 (s, 1H), 8.10 (d, 1H, J=12.3Hz), 7.93 (d, 3H, J=8.4Hz), 7.82 (d, 1H, J=8.4Hz), 7.49 (d, 2H, J=8.6Hz), 7.30 (s, 1H), 7.12 (d, 1H, J=8.4Hz); ES-MS *m/z* 337 (M+H).

Example 25: 6-Bromo-3-[(Z and E)-[4-(4-morpholinyl)anilino]methylidene]-1,3-dihydro-2H-indol-2-one.

Prepared according to Example 1 in 87% yield to give ~4:1 isomer mixture. Principal isomer: ^1H NMR ($\text{DMSO}-d_6$): δ 10.67 (d, 1H, $J=12.5\text{Hz}$), 10.56(s,1H), 8.57 (d,1H, $J=12.8\text{Hz}$), 7.48 (d,1H, $J=7.9\text{Hz}$), 7.35 - 7.29 (m,2H), 7.11 - 7.04 (m, 3H), 6.93 (s,1H), 3.76 (s,4H), 3.15 (s, 4H); ES-MS m/z 400, 402 ($M + 1$).

Example 26: 4- $\{[(Z)-(6\text{-Bromo-2-oxo-1,2-dihydro-3H-indol-3-ylidene)methyl}]\text{amino}\}$ benzamide.

Prepared according to Example 1 with 4-aminobenzamide in 70%
10 yield. ^1H NMR ($\text{DMSO}-d_6$): δ 10.75 (d,1H, $J=12.5\text{Hz}$), 10.65 (s,1H), 8.69 (d,1H, $J=12.5\text{Hz}$), 7.87 (d,3H, $J=8.6\text{Hz}$), 7.54 (d,1H, $J=7.9\text{Hz}$), 7.44 (d,2H, $J=8.6\text{Hz}$), 7.24 (s,1H), 7.09 (d,1H, $J=8.2\text{Hz}$), 6.95 (d,1H, $J=1.4\text{Hz}$); ES-MS m/z 356, 358 ($M - 1$).

15 **Example 27:** N-(4- $\{[(Z \text{ and } E)-(6\text{-Bromo-2-oxo-1,2-dihydro-3H-indol-3-ylidene)methyl}]\text{amino}\}$ phenyl)acetamide.

Prepared according to Example 1 with N-(4-aminophenyl)acetamide in 79% yield to give ~9:1 isomer mixture. Principal isomer: ^1H NMR ($\text{DMSO}-d_6$): δ 10.75 (d,1H, $J=12.5\text{Hz}$), 10.65 (s,1H), 8.69 (d,1H, $J=12.5\text{Hz}$), 7.87
20 (d,3H, $J=8.6\text{Hz}$), 7.54 (d,1H, $J=7.9\text{Hz}$), 7.44 (d,2H, $J=8.6\text{Hz}$), 7.24 (s,1H), 7.09 (d,1H, $J=8.2\text{Hz}$), 6.95 (d,1H, $J=1.4\text{Hz}$); ES-MS m/z 370, 372 ($M - 1$).

Example 28: 6-Bromo-3- $\{[(Z)-[4-(2\text{-hydroxyethyl})\text{anilino}]\text{methylidene}]-1,3\text{-dihydro-2H-indol-2-one}$.

25 Prepared according to Example 1 with 4-(2-hydroxyethyl)aniline in 85% yield. ^1H NMR ($\text{DMSO}-d_6$): δ 10.67 (d,1H, $J=12.8\text{Hz}$), 10.58 (s,1H), 8.61 (d,1H, $J=12.6\text{Hz}$), 7.50 (d,1H, $J=8.2\text{Hz}$), 7.29 (d,2H, $J=8.4\text{Hz}$), 7.19 (d,2H, $J=8.2\text{Hz}$), 7.06 (d,1H, $J=8.1\text{Hz}$), 6.93 (d,1H, $J=1.5\text{Hz}$), 4.60 (t,

1H, J=5.1Hz), 3.55 (m, 2H), 2.66 (t, 2H, J= 7.0Hz); MS-ES *m/z* 357, 359 (M - 1).

Example 29: 1-[(Z)-[4-(4-Morpholinyl)anilino]methylidene]-1,3-dihydro-5 2H-pyrrolo[3,2-f]quinolin-2-one.

Prepared according to Example 13 with 4-(4-morpholino)aniline in 65% yield. ¹H NMR (DMSO-d₆): δ 11.77 (d, 1H, J=12.3Hz), 10.88 (s, 1H), 8.77 (m, 2H), 8.69 (d, 1H, J=4.0Hz), 7.67 (d, 1H, J=8.6Hz), 7.5 - 7.4 (m, 4H), 6.98 (d, 2H, J=8.8Hz), 3.72 (t, 4H, J=4.5Hz), 3.07 (t, 4H, J=4.6Hz); ES-MS 10 *m/z* 373 (M+H).

Example 30: 1-[(Z)-[4-(4-Pyridinylmethyl)anilino]methylidene]-1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one.

Prepared according to Example 13 with 4-(4-pyridylmethyl)aniline in 15 59% yield. ¹H NMR (DMSO-d₆): δ 11.74 (d, 1H, J=12.1Hz), 10.93 (s, 1H), 8.80 - 8.75 (m, 2H), 8.70 (d, 1H, J=3.9Hz), 8.44 (d, 2H, J= 5.3Hz), 7.70 (d, 1H, J=8.6Hz), 7.47 - 7.43 (m, 3H), 7.39 (d, 1H, J=8.8Hz), 7.28 (d, 2H, J=8.1Hz), 7.24 (d, 2H, J=5.1Hz), 3.95 (s, 2H); ES-MS *m/z* 379 (M+H).

20 Example 31: N-(4-[(Z)-[2-Oxo-2,3-dihydro-1H-pyrrolo[3,2-f]quinolin-1-ylidene)methyl]amino}phenyl)acetamide.

Prepared according to Example 13 with N-(4-aminophenyl)acetamide in 72% yield. ¹H NMR (DMSO-d₆): δ 11.77 (d, 1H, J=12.1Hz), 10.92 (s, 1H), 9.95 (s, 1H), 8.8 - 8.77 (m, 2H), 8.70 (d, 1H, J=3.5Hz), 7.69 (d, 1H, J=8.8Hz), 25 7.60 (d, 2H, J=9.0Hz), 7.5 - 7.43 (m, 3H), 7.39 (d, 1H, J=8.6Hz), 2.01 (s, 3H); ES-MS *m/z* 345 (M+H).

Example 32: 1-[(Z)-[4-(2-Hydroxyethyl)anilino]methylidene]-1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one.

Prepared according to Example 13 with 4-(2-hydroxyethyl)aniline in 75% yield. ¹H NMR (DMSO-d₆): δ 11.75 (d,1H, J=12.1Hz), 10.92 (s,1H), 8.82 - 8.79 (m,2H), 8.70 (d,1H, J=3.9 Hz), 7.70 (d,1H, J=8.6Hz), 7.5 - 7.4 (m 4H), 7.23 (d,2H, J=8.2Hz), 4.62 (t,1H, J=5.1Hz), 3.6 - 3.55 (m,2H), 2.70 5 (t,2H, J=7.0Hz); ES-MS *m/z* 332 (M+H).

Example 33: 4-**{[(Z)-(2-Oxo-2,3-dihydro-1H-pyrrolo[3,2-f]quinolin-1-ylidene)methyl]amino}benzamide**.

Prepared according to Example 13 with 4-aminobenzamide in 54% 10 yield. ¹H NMR (DMSO-d₆): δ 11.90 (d,1H, J=11.9Hz), 11.03 (s,1H), 8.87-8.83 (m,2H), 8.76 (d,1H, J=4.0Hz), 7.93-7.89 (m,3H), 7.78 (d,1H, J=8.8Hz), 7.61 (d,2H, J=8.6Hz), 7.52 (dd,1H, J=4.1,8.5Hz), 7.45 (d,1H, J=8.8Hz), 7.3 (s,1H); ES-MS *m/z* 331 (M+H).

15 **Example 34:** 1-**[(Z)-(4-Hydroxyanilino)methylidene]-1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one**.

Prepared according to Example 13 with 4-hydroxyaniline in 84% yield. ¹H NMR (DMSO-d₆): δ 11.76 (d,1H, J=12.3Hz), 10.90 (s,1H), 9.44 (s,1H), 8.77 - 8.68 (m,3H), 7.71 (d, 1H, J=8.8Hz), 7.44 - 7.33 (m,4H), 6.83 (d,2H, 20 J=8.6Hz); ES-MS *m/z* 304 (M+H).

Example 35: 6-(2-Furyl)-3-**[(Z)-[4-(4-morpholinyl)anilino]methylidene]-1,3-dihydro-2H-indol-2-one**.

Prepared from the dimethylaminomethylene derivative (Procedure E) 25 of 6-(2-furyl)oxindole (prepared according to Procedure I) with 4-(4-morpholino)aniline according to Procedure G in 90% yield. ¹H NMR (DMSO-d₆): δ 10.66 (d,1H, J=12.6Hz), 10.51 (s,1H), 8.51 (d,1H, J=12.8Hz), 7.65 (s,1H), 7.56 (d,1H, J=7.9Hz), 7.29 - 7.25 (m,3H), 7.1 (s,1H), 6.95 (d,2H,

J=8.8Hz), 6.77 (d,1H, J=2.9Hz), 6.52 (s,1H), 3.71 (s,4H), 3.05 (s,4H); ES-MS *m/z* 388 (M+H).

Example 36: N-[4-((Z and E)-[6-(2-Furyl)-2-oxo-1,2-dihydro-3H-indol-3-ylidene]methyl)amino]phenyl]acetamide.

Prepared according to Example 35 with N-(4-aminophenyl)acetamide in 29% yield. ¹H NMR (DMSO-d₆): δ 10.68 (d,1H, J=12.6Hz), 10.55 (s,1H), 9.91 (s,1H), 8.53 (d,1H, J=12.6Hz), 7.66 (s,1H), 7.57 - 7.54 (m,3H), 7.32 - 7.27 (m,3H), 7.10 (s,1H), 6.78 (d,1H, J=2.9Hz), 6.53 (s,1H), 2.00 (s,3H); ES-MS *m/z* 358 (M-H).

Example 37: 6-(2-Furyl)-3-((Z)-[4-(2-hydroxyethyl)anilino]methylidene)-1,3-dihydro-2H-indol-2-one.

Prepared according to Example 35 with 4-(2-hydroxyethyl)aniline in 63% yield. ¹H NMR (DMSO-d₆): δ 10.68 (d,1H, J=12.5Hz), 10.56 (s,1H), 8.57 (d,1H, J=12.6Hz), 7.66 (s,1H), 7.58 (d,1H, J=7.9Hz), 7.28 (br s,3H), 7.19 (d,2H, J=8.1Hz), 7.11 (s,1H), 6.78 (s,1H), 6.53 (s,1H), 4.61 (t,1H, J=4.9Hz), 3.58 - 3.55 (m, 2H), 2.67 (t, 2H, J=6.9Hz); ES-MS *m/z* 345 (M-H).

Example 38: 3-((Z)-[4-(4-Morpholinyl)anilino]methylidene)-6-vinyl-1,3-dihydro-2H-indol-2-one.

Prepared according to Procedure G with the dimethylaminomethylene derivative (Procedure E) of 6-vinyloxindole (Procedure I) and 4-(4-morpholino)aniline in 75% yield. ¹H NMR (DMSO-d₆): δ 10.65 (d,1H, J=2.8Hz), 10.43 (s,1H), 8.48 (d,1H, J=12.5Hz), 7.48 (d,1H, J=7.9Hz), 7.27 (d,2H, J=8.9Hz), 7.00 (d,1H, J=7.9Hz), 6.95 (d,2H, J=8.9Hz), 6.89 (s,1H), 6.87 - 6.62 (m,1H), 5.64 (d,1H, J=17.84Hz), 5.08 (d,1H, J=11.1Hz), 3.71 (t,4H, J=4.6Hz), 3.05 (t,4H, J=4.6Hz); ES-MS *m/z* 346 (M-H).

Example 39: 3- $\{(Z)-[4-(4\text{-Pyridinylmethyl})\text{anilino}]\text{methylidene}\}$ -6-vinyl-1,3-dihydro-2H-indol-2-one.

Prepared according to Example 38 with 4-(4-pyridylmethyl)aniline in 51% yield. ^1H NMR (DMSO-d_6): δ 10.66 (d, 1H, $J=12.5\text{Hz}$), 10.49 (s, 1H), 5 8.53 (d, 1H, $J=12.5\text{Hz}$), 8.43 (d, 2H, $J=5\text{Hz}$), 7.50 (d, 1H, $J=7.9\text{Hz}$), 7.32 (d, 2H, $J=8.6\text{Hz}$), 7.24 - 7.21 (m, 4H), 7.01 (d, 1H, $J=7.9\text{Hz}$), 6.89 (s, 1H), 6.70 - 6.63 (m, 1H), 5.65 (d, 1H, $J=17.5\text{Hz}$), 5.10 (d, 1H, $J=11.1\text{Hz}$), 3.91 (s, 2H); ES-MS m/z 352 (M-H).

10 Example 40: N-(4- $\{(Z)-[(2\text{-Oxo-6-vinyl-1,2-dihydro-3H-indol-3-ylidene})\text{methyl}]\text{amino}\}$ phenyl)acetamide.

Prepared according to Example 38 with N-(4-aminophenyl)acetamide in 42% yield. ^1H NMR (DMSO-d_6): δ 10.67 (d, 1H, $J=12.5\text{Hz}$), 10.47 (s, 1H), 9.90 (s, 1H), 8.50 (d, 1H, $J=12.5\text{Hz}$), 7.55 (d, 2H, $J=8.6\text{Hz}$), 7.49 (d, 1H, 15 $J=7.85\text{Hz}$), 7.30 (d, 2H, $J=8.9\text{Hz}$), 7.01 (d, 1H, $J=7.9\text{Hz}$), 6.89 (s, 1H), 6.70 - 6.63 (m, 1H), 5.65 (d, 1H, $J=17.5\text{Hz}$), 5.09 (d, 1H, $J=11.1\text{Hz}$), 2.00 (s, 3H); ES-MS m/z 318 (M-H).

Example 41: 3- $\{(Z)-[4-(2\text{-Hydroxyethyl})\text{anilino}]\text{methylidene}\}$ -6-vinyl-1,3-20 dihydro-2H-indol-2-one.

Prepared according to Example 38 with 4-(2-hydroxyethyl)aniline in 55% yield. ^1H NMR (DMSO-d_6): δ 10.66 (d, 1H, $J=12.5\text{Hz}$), 10.48 (s, 1H), 8.54 (d, 1H, $J=12.5\text{Hz}$), 7.51 (d, 1H, $J=7.9\text{Hz}$), 7.28 (d, 2H, $J=8.2\text{Hz}$), 7.19 (d, 2H, $J=8.2\text{Hz}$), 7.01 (d, 1H, $J=7.9\text{Hz}$), 6.9 (s, 1H), 6.88 - 6.63 (m, 1H), 5.65 25 (d, 1H, $J=17.5\text{Hz}$), 5.09 (d, 1H, $J=11.1\text{Hz}$), 4.60 (t, 1H, $J=5.0\text{Hz}$), 3.58 - 3.53 (m, 2H), 2.66 (t, 2H, $J=7.0\text{Hz}$); ES-MS m/z 305 (M-H).

Example 42: 6-(2-Furyl)-3- $\{(Z)-[4-(4\text{-pyridinylmethyl})\text{anilino}]\text{methylidene}\}$ -1,3-dihydro-2H-indol-2-one.

Prepared according to Example 35 with 4-(4-pyridylmethyl)aniline in 42% yield. ¹H NMR (DMSO-d₆): δ 10.67 (d,1H, J=112.6Hz), 10.56 (s,1H), 8.56 (d,1H, J=12.6Hz), 8.43 (d,2H, J=5.3Hz), 7.66 (s,1H), 7.57 (d,1H, J=8.1Hz), 7.34 - 7.22 (m,7H), 7.10 (s,1H), 6.79 (d,1H, J=3.3Hz), 6.53 (s,1H), 5 3.91 (s,2H); ES-MS *m/z* 391 (M-H).

Example 43: 6-(2-Furyl)-3-[(Z)-(4-hydroxyanilino)methylidene]-1,3-dihydro-2H-indol-2-one.

Prepared according to Example 35 with 4-hydroxyaniline in 44% yield. 10 ¹H NMR (DMSO-d₆): δ 10.63 (d,1H, J=12.8Hz), 10.49 (s,1H), 9.32 (s,1H), 8.46 (d,1H, J=12.8Hz), 7.65 (s,1H), 7.54 (d,1H, J=7.9Hz), 7.26 (d,1H, J=7.9Hz), 7.21 (d,2H, J=8.8Hz), 7.01 (s,1H), 6.77 - 6.74 (m,3H), 6.52 (s,1H); ES-MS *m/z* 317 (M-H).

15 Example 44: 3-[(Z)-[4-(4-Morpholinyl)anilino]methylidene]-6-(2-thienyl)-1,3-dihydro-2H-indol-2-one.

Prepared from the dimethylaminomethylene derivative (Procedure E) of 6-(2-thienyl)oxindole (prepared according to Procedure I) with 4-(4-morpholino)aniline according to Procedure G in 47% yield. 20 ¹H NMR (DMSO-d₆): δ 10.67 (d,1H, J=12.9Hz), 10.65 (s,1H), 8.52 (d,1H, J=12.8Hz), 7.55 (d,1H, J=7.9Hz), 7.42 (d,1H, J=5.1Hz), 7.36 (d,1H, J=3.3Hz), 7.29 (2H, 8.8Hz), 7.22 (d,1H, J=7.9Hz), 7.08 - 7.07 (m,1H), 7.04 (s,1H), 6.95 (d,2H, J=9.0Hz), 3.71 (t,4H, J=4.5Hz), 3.05 (t,4H, J=4.6Hz).

25 Example 45: N-[4-[(Z)-[2-Oxo-6-(2-thienyl)-1,2-dihydro-3H-indol-3-ylidene]methyl]amino)phenyl]acetamide.

Prepared according to Example 44 with N-(4-aminophenyl)acetamide in 33% yield. ¹H NMR (DMSO-d₆): δ 10.68 (d,1H, J=12.6Hz), 10.52 (s,1H), 9.91 (s,1H), 8.53 (d,1H, J=12.6Hz), 7.55 (d,3H, J=8.4Hz), 7.43 (d,1H,

J=5.1Hz), 7.37 (d,1H, J=3.1Hz), 7.32 (d,2H, J=8.8Hz), 7.23 (d,1H, J=9.2Hz), 7.07 (t,1H, J=4.3Hz), 7.04 (s,1H), 2.00 (s,3H); ES-MS *m/z* 374 (M-H).

Example 46: 6-Bromo-3-[(Z)-[3-(hydroxymethyl)anilino]methylidene]-1,3-dihydro-2H-indol-2-one.

Prepared according to Example 1 with 3-(hydroxymethyl)aniline in 54% yield. ¹H NMR (DMSO-d₆): δ 10.74 (d,1H, J=12.8Hz), 10.60 (s,1H), 8.67 (d,1H, J=12.8Hz), 7.53 (d,1H, J=8.2Hz), 7.30 (d,2H, J=7.3Hz), 7.24 (d,1H, J=8.2Hz), 7.07 (d,1H, J=9.3Hz), 7.02 (d,1H, 7.3Hz), 6.94 (s,1H), 5.23 (t,1H, J=5.7Hz), 4.48 (d,2H, J=5.7Hz); ES-MS *m/z* 343, 345 (M - 1).

Example 47: 6-Bromo-3-[(Z and E)-[4-(5-methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene]-1,3-dihydro-2H-indol-2-one.

Prepared according to Example 1 with 2-(4-aminophenyl)-3-methylpyrazoline-5-one in 45% yield as ~9:1 isomer mixture. Major isomer: ¹H NMR (DMSO-d₆): δ 10.73 (d,1H, J=12.6Hz), 10.60(s,1H), 9.82 (s,1H), 8.65 (d,1H, J=12.6Hz), 7.52 (d,1H, J=8.1Hz), 7.47 (d,2H, J=8.8Hz), 7.42 (d,2H, J=8.8Hz), 7.08 (d,1H, J=8.1Hz), 6.94 (s,1H), 5.54 (s, 1H), 2.23 (s,3H); ES-MS *m/z* 411, 413 (M + 1).

20

Example 48: 3-Ethyl-3-(4-[(Z)-[(7-oxo-6,7-dihydro-8H-[1,3]thiazolo[5,4-e]indol-8-ylidene)methyl]amino}phenyl)-2,6-piperidinedione.

A mixture of 8-dimethylamino-methylene-6,8-dihydro-1-thia-3,6-diazas-indacen-7-one (Proceure E, 0.040 g, 0.163 mmol), DL-aminoglutethimide (0.055 g, 0.237 mmol) in absolute ethanol (5 ml) was heated with stirring at 90°C for 16h. The reaction was diluted with ethanol and diethyl ether and the product collected by filtration to yield 0.038 g (54%) of the title compound. ¹H NMR (DMSO-d₆): δ 11.00 (d,1H, J=12.5Hz), 10.88 (s,1H), 10.83 (s,1H), 9.23 (s, H), 8.02 (d,1H, J=12.5Hz), 7.77 (d,1H, J=8.

6Hz), 7.42 (d,2H, J=8.9Hz), 7.31 (d,2H, J=8.9Hz), 7.09 (d,1H, J=8.2Hz), 2.44 - 2.34 (m,2H), 2.15 - 2.13 (m,2H), 1.86 - 1.02 (m,2H), 0.75 (t,3H, J=7.3Hz); ES-MS *m/z* 431 (M-H).

5 Example 49: 8-[(Z)-(4-Phenoxyanilino)methylidene]-6H-[1,3]thiazolo[5,4-e]indol-7-one.

Prepared according to Procedure G with 4-phenoxyaniline in 86% yield. ¹H NMR (DMSO-d₆): δ 11.02 (d,1H, J=12.5Hz), 10.81 (s,1H), 9.22 (s,1H), 7.99 (d,1H, J=12.3Hz), 7.76 (d,1H, J=8.42Hz), 7.46 (d,2H, J=9.0Hz), 7.37 (t,2H, J=7.9Hz), 7.12 - 7.06 (m,4H), 6.98 (d,2H, J=8.1Hz); ES-MS *m/z* 384 (M-H).

Example 50: 8-[(Z)-[4-(Benzyloxy)anilino]methylidene]-6H-[1,3]thiazolo[5,4-e]indol-7-one.

15 Prepared according to Procedure G with 4-(benzyloxy)aniline in 90% yield. ¹H NMR (DMSO-d₆): δ 10.96 (d,1H, J=12.5Hz), 10.77 (s,1H), 9.21 (s,1H), 7.95 (d,1H, J=12.5Hz), 7.74 (d,1H, J=8.4Hz), 7.43 (d,2H, J=7.1Hz), 7.39 - 7.35 (m,4H), 7.32 - 7.30 (m,1H), 7.09 - 7.04 (m,3H), 5.09 (s,2H); ES-MS *m/z* 398 (M-H).

20

Example 51: Methyl 4-(4-[[[(Z)-7-oxo-6,7-dihydro-8H-[1,3]thiazolo[5,4-e]indol-8-ylidene)methyl]amino}phenoxy]benzoate.

Prepared according to Procedure G with methyl 4-(4-aminophenoxy)benzoate in 35% yield. ¹H NMR (DMSO-d₆): δ 11.05 (d,1H, J=12.3Hz), 10.83 (s,1H), 9.22 (s,1H), 8.01 (d,1H, 12.3Hz), 7.95 - 7.93 (m,2H), 7.77 (d,1H, J=8.4Hz), 7.52 (d,2H, J=9.0Hz), 7.19 (d,2H, J=8.8Hz), 7.10 (d,1H, J=8.4Hz), 7.03 (d,2H, J=8.8Hz), 3.80 (s,3H); ES-MS *m/z* 442 (M-H).

Example 52: Methyl 3-(4-((Z)-[(7-oxo-6,7-dihydro-8H-[1,3]thiazolo[5,4-e]indol-8-ylidene)methyl]amino}phenoxy)benzoate.

Prepared according to Procedure G with methyl 3-(4-aminophenoxy)benzoate in 71% yield. ¹H NMR (DMSO-d₆): δ 11.05 (d,1H, J=12.5Hz), 10.82 (s,1H), 9.22 (s,1H), 8.01 (d,1H, J=12.3Hz), 7.77(d,1H, J=8.4Hz), 7.68 (d,1H, J=7.9Hz), 7.55 - 7.49 (m,3H), 7.41 (s,1H), 7.31 (dd,1H, J=2.3,8.2Hz), 7.14 (d,2H, J=8.8Hz), 7.09 (d,1H, J=8.4Hz), 3.08 (s,3H); ES-MS *m/z* 442 (M-H).

10 Example 53: 8-((Z)-[3-(Hydroxymethyl)anilino]methylidene)-6H-[1,3]thiazolo[5,4-e]indol-7-one.

Prepared according to Procedure G with 3-(hydroxymethyl)aniline in 83% yield. ¹H NMR (DMSO-d₆): δ 11.05 (d,1H, J=12.5Hz), 10.84 (s,1H), 9.23 (s,1H), 8.05 (d,1H, 12.3Hz), 7.77 (d, 1H, J=8.4Hz), 7.37 - 7.26 (m,3H),
15 7.10 (d,1H, J=8.4Hz), 7.06 (d,1H, J=7.5Hz), 5.27 (t,1H,J=5.7Hz), 4.52 (d,2H, J=5.9Hz); ES-MS *m/z* 322 (M-H).

Example 54: 3-((Z)-[(7-Oxo-6,7-dihydro-8H-[1,3]thiazolo[5,4-e]indol-8-ylidene)methyl]amino)benzamide.

20 Prepared according to Procedure G with 3-aminobenzamide in 84% yield. ¹H NMR (DMSO-d₆): δ 11.13 (d,1H, J=12.3Hz), 10.87 (s,1H), 9.23 (s,1H), 8.12 - 8.07 (m,2H), 7.84 (s,1H), 7.78 (d,1H, J=8.4Hz), 7.58 (t,2H, J=7.7Hz), 7.47 (t,2H, J=7.8Hz), 7.10 (d,1H, J=8.4Hz); ES-MS *m/z* 335 (M-H).

25

Example 55: 8-((Z)-[4-(5-Methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene)-6H-[1,3]thiazolo[5,4-e]indol-7-one.

A mixture of 8-dimethylamino-methylene-6,8-dihydro-1-thia-3,6-diazas-indacen-7-one (Procedure E, 0.040 g, 0.163 mmol), 2-(4-aminophenyl)-3-

methyl-pyrazoline-5-one (0.047 g, 0.248 mmol) in absolute ethanol (5 ml) was heated with stirring at 90°C for 16h. The reaction was diluted with ethanol and diethyl ether and the product collected by filtration to yield 0.052 (83%) of the title compound. ¹H NMR (DMSO-d₆): δ 11.08 (d,1H, J=12.3Hz), 10.84 (s,1H), 9.85 (s,1H), 9.23 (s,1H), 8.05 (d,1H, J=12.1Hz), 7.78 (d,1H, J=8.2Hz), 7.51 - 7.45 (m,4H), 7.10 (d,1H, J=8.4Hz), 5.55 (s,1H), 2.25 (s,3H); ES-MS *m/z* 388 (M-H).

Example 56: Methyl 4-[(Z)-[(7-oxo-6,7-dihydro-8H-[1,3]thiazolo[5,4-e]indol-8-ylidene)methyl]amino]benzoate.

Prepared according to Procedure G methyl 4-aminobenzoate in 47% yield. ¹H NMR (DMSO-d₆): δ 11.17 (d,1H, J=12.1Hz), 10.91 (s,1H), 9.25 (s,1H), 8.07 (d,1H, J=12.1Hz), 7.95 (d,2H, J=8.6Hz), 7.81 (d,1H, J=8.2Hz), 7.51 (d,2H, J=8.6Hz), 7.10 (d,1H, J=8.4Hz), 3.81 (s,3H); ES-MS *m/z* 350 (M-H).

Example 57: 4-[(Z)-[(7-Oxo-6,7-dihydro-8H-[1,3]thiazolo[5,4-e]indol-8-ylidene)methyl]amino]benzonitrile.

Prepared according to Procedure G with 4-aminobenzonitrile in 71% yield. ¹H NMR (DMSO-d₆): δ 11.14 (d,1H, J=11.9Hz), 10.92 (s,1H), 9.25 (s,1H), 8.04 (d,1H, J=11.9Hz), 7.82 (dd,3H, J=2.4, 8.7Hz), 7.59 (d,2H, J=8.8Hz), 7.10 (d,1H, J=8.4Hz); ES-MS *m/z* 317 (M-H).

Example 58: N-Methyl-N-(4-[(Z)-[(7-oxo-6,7-dihydro-8H-[1,3]thiazolo[5,4-e]indol-8-ylidene)methyl]amino]phenyl)acetamide.

A mixture of 8-dimethylamino-methylene-6,8-dihydro-1-thia-3,6-diazas-indacen-7-one (Procedure E, 0.040 g, 0.163 mmol), 4-amino-N-methylacetanilide (0.040 g, 0.244 mmol) in absolute ethanol (5 ml) was heated with stirring at 90°C for 16h. The reaction was diluted with ethanol

and diethyl ether and the product collected by filtration to yield 0.038 g (64%) of the title compound. ¹H NMR (DMSO-d₆): δ 11.03 (d, 1H, J=12.3Hz), 10.84 (s, 1H), 9.23 (s, 1H), 8.02 (d, 1H, J=12.3Hz), 7.78 (d, 1H, J=8.4Hz), 7.48 (d, 2H, J=8.2Hz), 7.35 (d, 2H, J=8.2Hz), 7.09 (d, 1H, J=8.4Hz), 3.11 (s, 3H), 1.76 (s, 5 3H); ES-MS *m/z* 363 (M-H).

Example 59: 1-[(Z)-(4-Phenoxyanilino)methylidene]-1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one.

Prepared according to Example 13 with 4-phenoxyaniline in 79%
10 yield. ¹H NMR (DMSO-d₆): δ 11.78 (d, 1H, J=12.1Hz), 10.93 (s, 1H), 8.81 - 8.78 (m, 2H), 8.70 (d, 1H, J=4.0Hz), 7.70 (d, 1H, J=8.6Hz), 7.56 (d, 2H, J=8.8Hz), 7.45 (dd, 1H, J=4.1, 8.5Hz), 7.41 - 7.35 (m, 3H), 7.12 - 7.05 (m, 3H), 6.98 (d, 2H, J=8.2Hz); ES-MS *m/z* 378 (M-H).

15 **Example 60: 1-[(Z)-[4-(Benzyloxy)anilino]methylidene]-1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one.**

Prepared according to Example 13 with 4-(benzyloxy)aniline in 59%
yield. ¹H NMR (DMSO-d₆): δ 11.80 (d, 1H, J=12.3Hz), 10.99 (s, 1H), 8.95 (d, 1H, J=8.1Hz), 8.82 - 8.77 (m, 2H), 7.73 (d, 1H, J=8.8Hz), 7.54 (dd, 1H, 20 J=4.1, 8.3Hz), 7.49 - 7.30 (m, 8H), 7.05 (d, 2H, J=9.0Hz), 5.11 (s, 2H), ES-MS *m/z* 394 (M+H).

Example 61: Methyl 4-[4-[(Z)-[(2-oxo-2,3-dihydro-1H-pyrrolo[3,2-f]quinolin-1-ylidene)methyl]amino]phenoxy]benzoate.

25 Prepared according to Example 13 with methyl 4-(4-aminophenoxy)benzoate in 69% yield. ¹H NMR (DMSO-d₆): δ 11.80 (d, 1H, J=12.1Hz), 10.94 (s, 1H), 8.83 - 8.80 (m, 2H), 8.70 (d, 1H, J=3.9Hz), 7.95 (d, 2H, J= 8.8Hz), 7.71 (d, 1H, J=8.8Hz), 7.62 (d, 2H, J=9.0Hz), 7.45 (dd, 1H,

J=4.1, 8.5Hz), 7.41 (d, 1H, J= 8.8Hz), 7.18 (d,2H, J= 8.8Hz), 7.03 (d,2H, J=8.8Hz), 3.80 (s,3H); ES-MS *m/z* 436 (M - H).

Example 62: Methyl 3-(4-((Z)-[(2-oxo-2,3-dihydro-1H-pyrrolo[3,2-f]quinolin-1-ylidene)methyl]amino)phenoxy)benzoate.

Prepared according to Example 13 with methyl 3-(4-aminophenoxy)benzoate in 45% yield. ¹H NMR (DMSO-d₆): δ 11.96 (d,1H, J=12.1Hz), 11.33 (s,1H), 9.44 (d,1H, J=8.2Hz), 9.01 - 8.98 (m,2H), 7.95 (d,1H, J=8.8Hz), 7.85 (dd,1H, J=4.9, 8.4Hz), 7.72 - 7.66 (m,4H), 7.54 (t,1H, J=8.0Hz), 7.42 (s,1H), 7.33 (d,1H, J=8.2Hz), 7.17 (d,2H, J=8.6Hz), 3.80 (s,3H); ES-MS *m/z* 438 (M+H).

Example 63: 3-Ethyl-3-(4-((Z)-[(2-oxo-2,3-dihydro-1H-pyrrolo[3,2-f]quinolin-1-ylidene)methyl]amino)phenyl)-2,6-piperidinedione.

15 Prepared according to Example 13 with DL-aminogluthethimide in 81% yield. ¹H NMR (DMSO-d₆): δ 11.72 (d,1H, J=11.7Hz), 10.93 (s,1H), 10.88 (s,1H), 8.82 - 8.77 (m,2H), 8.70 (d,1H, J=2.9Hz), 7.71 (d,1H, J=8.8Hz), 7.52 (d,2H, J=8.8Hz), 7.45 (dd,1H, J=4.1,8.5Hz), 7.40 (d, 1H, J=8.8Hz), 7.30 (d,2H, J=8.6Hz), 2.36 - 2.16 (m,2H), 2.15 - 2.12 (m,2H), 1.90 - 1.77 (m,2H), 20 0.76 (t,3H, J=7.3Hz); ES-MS *m/z* 425 (M-H).

Example 64: 1-[(Z)-(4-Benzoylanilino)methylidene]-1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one.

Prepared according to Example 13 with 4-benzoylaniline in 76% yield. 25 ¹H NMR (DMSO-d₆): δ 11.92 (d,1H, J=11.7Hz), 11.04 (s,1H), 8.88 (d,1H, J=11.7Hz), 8.84 (d,1H, J=8.6Hz), 8.73 (d,1H, J=3.1Hz), 7.80 - 7.53 (m,10H), 7.49 (dd,1H, J=4.1, 8.5Hz), 7.42 (d,1H, J=8.8Hz); ES-MS *m/z* 390 (M-H).

Example 65: 1-[(Z)-[3-(Hydroxymethyl)anilino]methylidene]-1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one.

Prepared according to Example 13 with 3-(hydroxymethyl)aniline in 78% yield. ¹H NMR (DMSO-d₆): δ 11.84 (d,1H, J=12.1Hz), 10.36 (s,1H),
5 8.86 - 8.80 (m,2H), 8.72 (d,1H, J=4.0Hz), 7.72 (d,1H, J=8.7Hz), 7.48 (dd,1H, J=4.1,8.5Hz), 7.43 - 7.33 (m,4H), 7.07 (d,1H, J=7.5Hz), 5.25 (s,1H), 4.53 (dd,2H, J=4.5Hz); ES-MS *m/z* 316 (M-H).

Example 66: 1-[(Z)-[4-(5-Methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene]-1H-pyrrolo[3,2-f]quinolin-2(3H)-one.

Prepared according to Example 13 with 2-(4-aminophenyl)-3-methylpyrazoline-5-one in 69% yield. ¹H NMR (DMSO-d₆): δ 11.83 (d,1H, J=11.9Hz), 10.05 (s,1H), 9.84 (s,1H), 8.86 - 8.82 (m,2H), 8.71 (d,1H, J=2.9Hz), 7.72 (d,1H, J=8.8Hz), 7.60 (d,2H, J=8.8Hz), 7.48 - 7.44 (m,3H),
15 7.41 (d,1H, J=8.8Hz), 5.55 (s,1H), 2.25 (s,3H); ES-MS *m/z* 382 (M-H).

Example 67: 1-[(E)-[4-[(E)-2-(4-Hydroxyphenyl)ethenyl]anilino]methylidene]-1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one.

Prepared according to Example 13 with 2-(4-hydroxyphenyl)ethenyl]aniline in 90% yield. ¹H NMR (DMSO-d₆): δ 11.86 (d,1H, J=11.9Hz), 10.95 (s,1H), 9.53 (s,1H), 8.86 - 8.83 (m,2H), 8.71 (d,1H, J=3.3Hz), 7.71 (d,1H, J=8.6Hz), 7.58 - 7.45 (m,5H), 7.42 - 7.38 (m,3H), 7.12 - 6.98 (m,2H), 6.74 (d,2H, J=8.6Hz); ES-MS *m/z* 404 (M-H).

Example 68: 3-[(Z)-[(2-Oxo-2,3-dihydro-1H-pyrrolo[3,2-f]quinolin-1-ylidene)methyl]amino}benzamide.

Prepared according to Example 13 with 3-aminobenzamide in 66% yield. ¹H NMR (DMSO-d₆): δ 11.86 (d,1H, J=11.9Hz), 10.98 (s,1H), 8.89 - 8.80 (m,2H), 8.72 (d,1H, J=2.9Hz), 8.06 (s,1H), 7.88 (s,1H) 7.74 - 7.69

(m,2H), 7.59 (d,1H, J=7.7Hz), 7.51 - 7.45 (m,3H), 7.41 (d,1H, J=8.8Hz); ES-MS *m/z* 329 (M-H).

Example 69: 4-[(Z)-[(2-Oxo-2,3-dihydro-1H-pyrrolo[3,2-f]quinolin-1-ylidene)methyl]amino]benzonitrile.

Prepared according to Example 13 with 4-aminobenzonitrile in 82% yield. ¹H NMR (DMSO-d₆): δ 11.84 (d,1H, J=11.7Hz), 11.03 (s,1H), 8.84 - 8.73 (m,2H), 8.72 (dd,1H, J=1.2,4.1Hz), 7.82 (d, 2H, J=8.6Hz), 7.76 (d,1H, J=8.8Hz), 7.69 (d,2H, J=8.8Hz), 7.48 (dd,1H, J=4.1,8.5Hz), 7.41 (d,1H, J=8.6Hz); ES-MS *m/z* 311 (M-H).

Example 70: Methyl 4-[(Z)-[(2-oxo-2,3-dihydro-1H-pyrrolo[3,2-f]quinolin-1-ylidene)methyl]amino]benzoate.

Prepared according to Example 13 with methyl 4-aminobenzoate in 44% yield. ¹H NMR (DMSO-d₆): δ 11.91 (d,1H, J=11.8Hz), 11.03 (s,1H), 8.88 - 8.83 (m,2H), 8.72 (dd,1H, J=1.3,4.1Hz), 7.95 (d,2H, J=8.6Hz), 7.75 (d,1H, J=8.9Hz), 7.63 (d,2H, J=8.6Hz), 7.49 (dd,1H, J=3.9,8.6Hz), 7.41 (d,1H, J=8.6Hz), 3.82 (s,3H); ES-MS *m/z* 344 (M-H).

Example 71: 8-[(Z)-[4-[(2-(Diethylamino)ethyl)sulfonyl]anilino)methylidene]-6H-[1,3]thiazolo[5,4-e]indol-7-one.

Prepared according to Procedure G with 4-[2-(diethylamino)ethylsulfonyl]aniline in 18% yield. ¹H NMR (DMSO-d₆): δ 11.20 (d,1H, J=11.9Hz), 10.93 (s,1H), 9.26 (s,1H), 8.08 (d,1H, J=11.9Hz), 7.87 (d,2H, J=8.6Hz), 7.82 (d,1H, J=8.42Hz), 7.63 (d,2H, J=8.6Hz), 7.11 (d,1H, J=8.24Hz), 3.46 - 3.38 (m,2H), 2.8 - 2.7 (m,2H), 2.3 - 2.2 (m,4H), 0.80 (t,6H, J=7.0Hz); ES-MS *m/z* 455 (M-H).

Example 72: 1-[(Z)-(4-{[2-(Diethylamino)ethyl]sulfonyl}anilino)methylidene]-1H-pyrrolo[3,2-f]quinolin-2(3H)-one.

Prepared according to Example 13 with 4-[2-(diethylamino)ethylsulfonyl]aniline in 64% yield. ¹H NMR (DMSO-d₆): δ 11.91 (d, 1H, J=11.5Hz), 11.04 (s, 1H), 8.88 - 8.84 (m, 2H), 8.73 (d, 1H, J=3.3Hz), 7.87 (d, 2H, J=8.6Hz), 7.78 - 7.72 (m, 3H), 7.50 (dd, 1H, J=4.1, 8.5Hz), 7.41 (d, 1H, J=8.6Hz), 3.45 - 3.38 (m, 2H), 2.71 - 2.67 (m, 2H), 2.32 (q, 4H, J=7.0Hz), 0.80 (t, 6H, J=7.1Hz); ES-MS *m/z* 449 (M-H).

10 Examples 73 - 88 are described in Procedure J (vide infra).

Example 89: 3,6-Dihydro[1,2,3]triazolo[4,5-e]indole-7,8-dione 8-[N-(4-methoxyphenyl)hydrazone].

See Procedure H.

15

Example 90: 3,6-Dihydro[1,2,3]triazolo[4,5-e]indole-7,8-dione 8-{N-[4-(1,3-oxazol-5-yl)phenyl]hydrazone}.

Prepared according to Procedure H with 4-(1,3-oxazol-5-yl)phenylhydrazine hydrochloride in 61% yield as ~2:1 mixture of geometric isomers. ¹H NMR (DMSO-d₆): δ 7.09 and 7.26 (2 d, 1:2 ratio, J= 8.6 Hz, 1H); 7.55 and 7.85 (ABq, J = 8.7 Hz, 4H); 7.65 (s, 1H); 7.85 and 7.98 (2 d, 2:1 ratio, partially obscured by ABq, J = 8.6 Hz, 1H); 8.45 (s, 1H); 10.87 and 11.33 (2 s, ~2:1 ratio, 1H). 12.95 (brd, 1H). APCI-MS *m/z* 344 (M-H).

25 **Example 91: 3,6-Dihydro[1,2,3]triazolo[4,5-e]indole-7,8-dione 8-[N-(4-methylphenyl)hydrazone].**

Prepared according to Procedure H with 4-methylphenylhydrazine hydrochloride in 18% yield as ~1:1 mixture of E and Z isomers. ¹H NMR (DMSO-d₆): δ 2.28 (s, 3H); 7.19 (d, J= 8.4 Hz, 0.5H); 7.15 - 7.35 (m, 3.5H);

7.62 (d, $J = 8.4$ Hz, 1H); 7.76 (d, $J = 8.4$ Hz, 0.5H); 7.93 (d, $J = 8.4$ Hz, 0.5H); 10.7 (s, 0.5H); 11.2 (s, 1H); 12.3 and 12.35 2 s, 1H). APCI-MS m/z 291 (M-H)⁻.

5 Example 92: 3,6-Dihydro[1,2,3]triazolo[4,5-e]indole-7,8-dione 8-[N-(4-[(E)-2-(2-pyridinyl)ethenyl]phenyl)hydrazone].

Prepared according to Procedure H with 4'-hydrazino-2-stilbazole dihydrochloride (TCI, Inc.) in 89% yield as ~1:1 mixture of E and Z isomers as the hydrochloride salt. ¹H NMR (DMSO-d₆): δ 7.10 and 7.26 (2 d, $J = 8.6$ Hz, 1H); 7.37 and 7.41 (2 d, $J = 16$ Hz, 1H); 7.55 (d, $J = 8.6$ Hz, 1H); 7.68 (m, 1H); 7.74 - 7.9 (m, 3.5H); 7.99 (d, $J = 16$ Hz, 1H); 7.99 (d, $J = 8.6$ Hz, 0.5H); 8.16 (t, $J = 8.6$ Hz, 1H); 8.34 (m, 1H); 8.74 (d, $J = 5.5$ Hz, 1H); 10.9 and 11.4 (2 s, 1H); 13.0 (s, 1H). APCI-MS m/z 380 (M-H)⁻.

15 Example 93: 6H-[1,3]Thiazolo[5,4-e]indole-7,8-dione 8-[N-(3-methoxyphenyl)hydrazone].

Prepared according to Procedure H using 6H-[1,3]thiazolo[5,4-e]indole-7,8-dione (Procedure A) and 3-methoxyphenylhydrazine hydrochloride in 44% yield. APCI-MS m/z 323 (M-H)⁻.

20

Example 94: 5-Hydroxy-4,6-dimethyl-1H-indole-2,3-dione 3-[N-(4-methylphenyl)hydrazone].

5-Hydroxy-4,6-dimethyl-1H-indole-2,3-dione was prepared from 3,5-dimethyl-4-hydroxyaniline according to Procedure A: ¹H NMR (DMSO-d₆): δ 2.17 (s, 3H), 2.30 (s, 3H), 6.45 (s, 1H), 8.29 (s, 1H), 10.65 (s, 1H); ESI-MS m/z 190 (M-H)⁻. The isatin was combined with 4-methylphenylhydrazine hydrochloride according to Procedure H to provide the title compound in 41% yield. ¹H NMR (DMSO-d₆): δ 2.15 (s, 3H), 2.24

(s, 3H), 2.43 (s, 3H), 6.46 (s, 1H), 7.16 (m, 4H), 7.88 (s, 1H), 10.64 (s, 1H), 12.85 (s, 1H). APCI-MS m/z 294 (M-H, 45%).

5 Example 95: (Z)-3,6-Dihydro[1,2,3]triazolo[4,5-e]indole-7,8-dione 8-[N-[4-(trifluoromethyl)phenyl]hydrazone].

Prepared according to Procedure H with 4-trifluoromethylphenylhydrazine hydrochloride in 32% yield. ^1H NMR (DMSO- d_6): δ 7.26 (d, J = 8.7 Hz, 1H); 7.59 and 7.83 (ABq, J = 8.4 Hz, 4H); 7.89 (brd d, J = 8.7 Hz, 1H); 10.9 (s, 1H); 13.0 (s, 1H). APCI-MS m/z 345 (M-H).

Example 96: 6H-[1,3]Thiazolo[5,4-e]indole-7,8-dione 8-[N-(3-fluorophenyl)hydrazone] (Z-isomer)

6H-1-Thia-3,6-diaza-as-indacen-7,8-dione (Procedure A, 50 mg, 0.25 mmol) was combined with 3-fluorophenylhydrazine hydrochloride (50 mg, 0.3 mmol) in 2 ml of ethanol and heated at 70 °C for 6 hrs. The product was collected by filtration of the hot solution, washing with ethanol and diethyl ether, to give 47 mg (60%) of the title compound as a brown solid. ^1H NMR (DMSO- d_6): δ 6.82 (t, J = 8.7 Hz, 1H); 7.10 (d, J = 8.6 Hz, 1H); 7.37 (m, 2H); 7.41 (dd, 1H); 7.97 (d, J = 8.6 Hz, 1H); 9.25 (s, 1H); 11.2 (s, 1H). ES-MS m/z 311 (M-1).

Example 97: (Z)-6H-[1,3]Thiazolo[5,4-e]indole-7,8-dione 8-[N-(4-fluorophenyl)hydrazone].

Prepared according to Example 96 with 6H-1-thia-3,6-diaza-as-indacen-7,8-dione (Procedure A) and 4-fluorophenylhydrazine hydrochloride in 49% yield. ^1H NMR (DMSO- d_6): δ 7.11 (d, J = 8.6 Hz, 1H); 7.25 (t, J = 8.8 Hz, 2H); 7.52 (dd, J = 4.8, 8.8 Hz, 2H); 7.94 (d, J = 8.6 Hz, 1H); 9.27 (s, 1H); 11.2 (s, 1H); 12.6 (s, 1H). ES-MS m/z 311 (M-1).

Example 98: (Z)-6H-[1,3]Thiazolo[5,4-e]indole-7,8-dione 8-[N-(4-bromophenyl)hydrazone].

Prepared according to Example 96 with 6H-1-thia-3,6-diaza-as-indacen-7,8-dione (Procedure A) and 4-bromophenylhydrazine hydrochloride in 62% yield. ^1H NMR ($\text{DMSO}-d_6$): δ 7.16 (d, $J = 8.5$ Hz, 1H); 7.50 and 7.62 (Abq, $J = 8.8$ Hz, 4H); 8.02 (d, $J = 8.5$ Hz, 1H); 9.33 (s, 1H); 11.3 (s, 1H); 12.6 (s, 1H). ES-MS m/z 371, 373 ($\text{M}-1$).

10 Procedure J - Method for preparation of a solution phase library containing compounds of the invention.

Set forth below are a selected number of synthesis examples that illustrate the solution library techniques that can be used to obtain the compounds of the invention. It is believed that One of ordinary skill in the art
15 can follow this procedure or modify it accordingly without undue experimentation in order to obtain the substitutions disclosed above. The following examples are illustrative examples of the solution phase synthesis, not intended to limit the scope of the invention in any way.

20 Synthesis of Intermediates:

5-Bromo-7-azaoxindole

a) 3,3,5 Tribromooxindole

A solution of 3,3-dibromo-7-azaoxindole (5.0 g, 13.4 mmol) in tert-
25 BuOH (100 mL) and water (100 mL) is stirred at room temperature and bromine (5.5 g, 34.3 mmol) is added dropwise over 20 min. A saturated aqueous solution of sodium bicarbonate (approx. 15 mL) is added dropwise over 30 min to raise the pH of the solution to 6.5. The yellow solid formed is collected by filtration. The filtrate is condensed to approx. 100 mL and
30 extracted with CHCl_3 (2 x 50 mL). The combined organic extracts are dried

over anhydrous magnesium sulfate and the solvent is evaporated under reduced pressure to leave a yellow solid. The solids are combined and dried under vacuum to give 3,3,5 tribromooxindole as a yellow solid, 6.25 g (98%). ¹H NMR (CDCl₃) δ 9.4 (br s, 1H), 8.28 (d, 1H, J = 2 Hz), 7.95 (d, 1H, J = 2 Hz).

b) 5-Bromo-7-azaoxindole

A solution of 3,3,5-tribromooxindole (5.0 g, 13.4 mmol) in fresh THF (100 mL) is stirred at room temperature and a saturated aqueous solution of ammonium chloride (100 mL) is added. The flask is placed in a water bath and activated zinc dust (15.0 g, 230 mmol) is added. The mixture is stirred for 20 min and the zinc is removed by filtration through a pad of diatomaceous earth. The organic layer is separated and the aqueous layer is extracted with THF (20 mL). The combined organic layers were washed with saturated brine solution, dried over anhydrous magnesium sulfate and the solvent removed under reduced pressure. The brown residue is triturated with water (20 mL) and the tan solid is collected by filtration and dried under vacuum to give 5-bromo-7-azaoxindole as a tan solid, 2.02 g (71%). ¹H NMR (DMSO-d₆) δ 11.13 (s, 1H), 8.15 (s, 1H), 8.76 (s, 1H), 3.57 (s, 2H). MS (AP -ve) 211 (100) (M-H).

5-Phenyl-7-azaoxindole

To a stirred mixture of 5-bromo-7-azaoxindole (213 mg, 1 mmol) and phenylboronic acid (183 mg, 1.5 mmol) in toluene (6 ml) and ethanol (6 ml) were added 1 M sodium carbonate solution (2.5 ml, 2.5 mmol), lithium chloride (127 mg, 3 mmol) and dichlorobis(triphenylphosphine)palladium(II) (35 mg, 0.05 mmol) under N₂ atmosphere. The reaction mixture was heated to reflux at 95 °C for 18 hours. The reaction mixture was diluted with chloroform (50 ml) and washed with brine (20 ml). The aqueous layer was thoroughly extracted with chloroform. The combined organic layers were

dried over anhydrous MgSO_4 , filtered and evaporated under vacuum to give crude product. Trituration of the crude product with diethyl ether yielded 5-phenyl-7-azaoxindole as a yellow solid (108 mg, 51.4%). ^1H NMR ($\text{DMSO}-d_6$): δ 11.04 (s, 1H), 8.32 (s, 1H), 7.83 (s, 1H), 7.60 (d, 2H, $J = 7.4$ Hz), 7.44 (t, 2H, $J = 7.4$ Hz), 7.32 (t, 1H, $J = 7.4$ Hz), 3.58 (s, 2H). MS (-ve APCI): 210 (48, M^+), 209 (100, M-H).

5-(Furan-2-yl)-7-aza-oxindole

5-Bromo-7-azaoxindole (0.75 g, 3.52 mmol), 2-tributyltinfuran (1.26 g, 3.52 mmol), tetraethylammonium chloride hydrate (1.94 g, 10.6 mmol) were combined and dissolved in acetonitrile (10 mL) at room temperature under an atmosphere of nitrogen. Bistriphenylphosphine dichloropalladium (II) (0.25 g, 0.35 mmol) was added and the reaction was warmed to 85 °C for 16 h. The reaction was cooled to room temperature and diluted with aqueous KF (10%, 60 mL). This was stirred for 20 minutes and then diluted with EtOAc (60 mL). The biphasic system was passed through celite, the layers separated, and the volatiles removed in vacuo. The resulting residue was triturated with diethyl ether and the solids were collected by filtration to afford a light yellow solid (0.28 g, 36% yield). ^1H NMR 300 MHz ($\text{DMSO}-d_6$) δ 11.18 (bs, 1H); 8.45 (s, 1H); 7.92 (s, 1H); 7.79 (s, 1H); 6.95 (d, 1H); 6.60 (d, 1H); 3.64 (s, 2H). APCI m/z 201 ($\text{M}+1$).

5-(3-Thienyl)-7-aza-oxindole

5-Bromo-7-azaoxindole (0.20 g, 0.94 mmol), 3-tributyltinthiophene (0.42 g, 1.12 mmol), tetraethylammonium chloride hydrate (0.16 g, 0.94 mmol) were combined and dissolved in acetonitrile (10 mL) at room temperature under nitrogen. Bistriphenylphosphine dichloropalladium (II) (0.033 g, 0.047 mmol) was added and the reaction was warmed to 85 °C for 20 h. Fresh catalyst was added to the reaction mixture, Bistriphenylphosphine dichloropalladium (II) (0.033 g, 0.047 mmol) and the

reaction continued to stir at 85 °C for 24 h. The reaction was cooled to room temperature and diluted with water (20 mL) and EtOAc (20 ml). The biphasic system was passed through celite and the layers were separated. The organic layer was washed with brine (10 mL) and dried over sodium sulfate. The volatiles were removed in vacuo. The resulting residue was triturated with diethyl ether and collected by filtration (0.16 g, 80% yield). ¹H NMR 400 MHz (DMSO-d₆) δ 11.03 (bs, 1H); 8.43 (s, 1H); 7.92 (s, 1H); 7.84 (s, 1H); 7.60 (m, 1H); 7.53 (d, 1H); 3.58 (s, 2H).

10 **5-Carboethoxy-7-azaoxindole**

To a mixture of 5-bromo-7-azaoxindole (213 mg, 1 mmol) in dimethylsulfoxide (1 ml) and ethanol (5 ml) in Parr bomb were added triethylamine (0.31 ml, 2.25 mmol), palladium acetate (33.7 mg, 0.15 mmol), and 1,3-(bis(diphenylphosphino)propane (61.9 mg, 0.15 mmol). Carbon monoxide gas (40 atm) was applied and the reaction mixture was heated at 95 °C for 18 hours with vigorously stirring. The reaction mixture was diluted with diethyl ether (50 ml) and washed with water (10 ml). The aqueous layer was thoroughly extracted with diethyl ether. The combined organic layers were dried over anhydrous MgSO₄, filtered and evaporated under vacuum to give crude product. Trituration of the crude product with methanol yielded 5-(carboethoxy)-7-azaoxindole as a tan solid (53 mg, 25.7%). ¹H NMR (DMSO-d₆): δ 11.39 (s, 1H), 8.62 (s, 1H), 7.95 (s, 1H), 4.27 (q, 2H, J = 7 Hz), 3.59 (s, 2H), 1.28 (t, 3H, J = 7 Hz). MS (-ve APCI): 205 (4, M-H).

6-Chloro-7-azaoxindole**a) 6-Chloro-7-aza-3,3-dibromooxindole**

6-Chloro-7-azaindole was prepared according to the procedure of Minakata et al; Synthesis, 1992, 661-663. To a stirred solution of 1.32 g (8.7 mmol) of 6-chloro-7-azaindole in tert-butanol (80 mL) was added 9.9g (28 mmol) of 90% pyridine hydrobromide perbromide resulting in a thick yellow precipitate forming immediately. The reaction was concentrated and the crude chromatographed on silica gel eluting with hexane to 90% hexane/10% EtOAc gradient to give 2.36 g of the title compound as a white solid, containing about 30% of 5-bromo-6-chloro-7-aza-3,3-dibromooxindole as an inseparable impurity. ^1H NMR (CDCl_3) δ 7.16 (d, J = 8.0 Hz, 1H), 7.81 (d, J = 8.0 Hz, 1H), 9.0 (bs, 1H). (5-Bromo-6-chloro-7-aza-3,3-dibromooxindole, 8.05 (s, 1H), 9.0 (bs, 1H).

15 b) 6-Chloro-7-azaoxindole

A solution of 2.36 g (7.26 mmol) of the mixture of 6-chloro-7-aza-3,3-dibromooxindole and 5-bromo-6-chloro-7-aza-3,3-dibromooxindole in THF (70 mL) and saturated ammonium chloride solution (70 mL) was treated with 6 g (92 mmol) of powdered zinc. After stirring for 2h another 6 g (92 mmol) portion of zinc was added and stirring continued another 2h. The zinc was filtered off and washed with ether. Ether phase separated and the aqueous phase washed twice with a 1:1 mixture of THF/ether. Combined ether fractions were dried over magnesium sulfate, filtered and concentrated. The crude was loaded onto 7.5 g of silica gel and chromatographed on silica gel eluting with 90% hexane/10% ethyl acetate to a 66% hexane/ 33% ethyl acetate gradient to give 0.647 g of the title compound, plus 0.243 g of 5-bromo-6-chloro-7-azaoxindole. ^1H NMR ($\text{DMSO}-d_6$): δ 3.57 (s, 2H), 7.04 (d, J = 7.6 Hz, 1H), 7.61 (d, J = 7.6 Hz, 1H), 11.2 (bs, 1H).

7-Azaoxindole

a) 3,3-Dibromo-7-azaoxindole

A solution of 7-azaindole (4.0g, 34 mmol) in tert-BuOH (200 mL) is stirred at room temperature and pyridinium perbromide (32.5g, 0.1 mol) is added in portions over 30 min. and the reaction mixture is stirred for 3 h. Pyridinium perbromide (10.8 g, 33 mmol) is added and the mixture is stirred for a further 2 h. The tert-BuOH is evaporated under reduced pressure and the residue is partitioned between water (300 mL) and EtOAc (300 mL). The organic layer is separated and the aqueous layer is extracted with EtOAc. The combined organic layers are washed with water (2 x 50 mL), and brine. The organic layer is dried over anhydrous MgSO_4 , filtered and the solvent evaporated. Trituration of the residue with CH_2Cl_2 gives a white solid which is collected by filtration and dried under vacuum to give 3,3-dibromo-7-azaoxindole, 8.35g. $^1\text{H-NMR}$ (DMSO-d_6) δ 11.99 (s, 1H), 8.21 (dd, 1H, J = 5.1, 1.5 Hz), 8.00 (dd, 1H, J = 7.5, 1.5 Hz), 7.17 (dd, 1H, J = 7.5, 5.1 Hz). MS (+ve ES) 293 (28), (M+H), 147 (100).

b) 7-Azaoxindole

A solution of 3,3-dibromo-7-azaoxindole (2.0g, 7.2 mmol) in THF (50 mL) is stirred at room temperature and a saturated aqueous solution of NH_4Cl is added. Activated zinc powder is added and the reaction mixture is stirred for 2 h. The zinc is removed by filtration through a pad of diatomaceous earth and the organic layer is separated. The aqueous layer is extracted with THF (10 mL) and the combined organic layers are dried over anhydrous MgSO_4 , filtered and evaporated. The residue is slurried in 10:1 CHCl_3 :MeOH (15 mL) and filtered through a pad of silica gel and the filtrate is evaporated. The residue is triturated with water and the solid is collected by filtration and dried under vacuum to give 7-azaoxindole, 0.668g (70%). $^1\text{H NMR}$ (DMSO-d_6) δ 10.94 (s, 1H), 8.02 (d, 1H, J = 5.2 Hz), 7.52

(d, 1H, J = 6.8 Hz), 6.90 (dd, 1H, J = 6.8, 5.2 Hz), 3.53 (s, 2H). MS(AP-ve) 133 (100) (M-H)

4-Aza-oxindole

5 a) Diethyl (3-nitropyridin-2-yl)-malonate

Sodium hydride (60% dispersion in oil, 5.57 g, 0.14 mol) was carefully washed with hexanes under nitrogen before the addition of DMSO (115 mL). Diethyl malonate (22.3 g, 0.14 mol) was added dropwise over 20 min and the mixture was stirred for an additional 30 min at room temperature. 2-
10 Chloro-3-nitropyridine (10 g, 0.06 mol) was added to the reaction and the reaction was placed in a pre-heated oil bath set to 100°C for 15 min. The reaction was cooled to room temperature and poured into aqueous ammonium chloride (saturated solution, 150 mL). The aqueous solution was extracted with EtOAc:Hexanes (1:1) four times (200 mL each) and the
15 organic layers were combined. The organics were concentrated to afford a solid that was recrystallized from a minimal amount of EtOAc:Hexanes (1:1) (12.5 g, 70% yield). APCI MS *m/z* 281 (M-1).

b) Ethyl 2-(3-nitro-pyridin-2-yl)-acetate

20 Diethyl (3-nitropyridin-2-yl)-malonate (12.5 g, 0.044 mol) was dissolved in DMSO (150 mL) and water (0.79 mL, 0.044 mol) and lithium chloride (4.65 g, 0.11 mol) were added at room temperature under nitrogen. The reaction was warmed to 100°C 12 h and more lithium chloride (1 g) was added to the reaction. The reaction was heated for another 5 hours and
25 cooled to room temperature. Brine (150 mL) was added to the reaction before extracting with EtOAc (3x, 275 mL each). The organics were combined and dried over sodium sulfate, then concentrated in vacuo. The resulting residue was triturated with diethyl ether and collected by filtration (8.6 g, 92 % yield). 1H NMR 400 MHz (DMSO-d₆) δ 8.83 (m, 1H); 8.53 (m,
30 1H); 7.65 (m, 1H); 4.23 (s, 2H); 4.07 (m, 2H); 1.16 (m, 3H).

c) Ethyl 2-(3-amino-pyridin-2-yl)-acetate

Under an atmosphere of nitrogen, Pd/C (10%, 1.36 g) was charged to a round bottom flask. Ethyl 2-(3-nitro-pyridin-2-yl)-acetate (8.6 g, 0.41 mol) was dissolved in ethanol (200 mL) and added to the reaction vessel. The reaction was placed under an atmosphere of hydrogen and stirred at room temperature for 30 min. The reaction was filtered through celite and the filtrate was concentrated in vacuo to afford the product as a tan solid (6.94g, 94% yield).

10

d) 4-Azaoxindole

Ethyl 2-(3-amino-pyridin-2-yl)-acetate (6.94 g, 0.038 mol) was dissolved in diethyl ether (100 mL) at room temperature. Hydrochloric acid (2M, 35 mL) was added and the reaction was stirred for 30 minutes. The volatiles were removed to afford a brown solid that was re-crystallized from ethanol and diethyl ether (4.0 g, 62% yield). ¹H NMR 400 MHz (DMSO-d₆) δ 12.35 (s, 1H); 8.12 (m, 1H); 7.90 (m, 1H); 7.14 (m, 1H); 5.75 (s, 2H). Electrospray MS *m/z* 135 (M+1).

20 6-(Furan-2-yl)-oxindole

6-Bromo-oxindole (0.40 g, 1.88 mmol), 2-tributyltinfuran (0.71 mL, 2.26 mmol), and tetraethylammonium chloride hydrate (0.31 g, 1.88 mmol) were combined and dissolved in acetonitrile (15 mL). The palladium catalyst, bistrisphenylphosphinedichloropalladium (II) (0.66g, 0.09 mmol) was added and the reaction was warmed to 85 °C under nitrogen for 20 h. The reaction was cooled to room temperature and diluted with water (15 mL) before passing the mixture through celite. The pad of celite was washed with EtOAc and the filtrates were combined and separated. The aqueous layer was washed with EtOAc (2X 20 mL each). The combined organic phases

30

were washed with brine and dried over sodium sulfate. The volatiles were removed in vacuo. The resulting residue was triturated with diethyl ether and the solid was collected by filtration (0.13 g, 34%). ¹H NMR 300 MHz (DMSO-d₆) δ 10.5 (s, 1H); 7.75 (s, 1H); 7.30 (m, 2H); 7.11 (s, 1H); 6.91 (m, 1H); 6.60 (m, 1H); 3.52 (s, 2H).

Synthesis of Monomers:

Monomer 6: 5-Carboethoxy-3-ethoxymethylene-7-azaoxindole

10 (Procedure J - 1)

5-Carboethoxy-7-azaoxindole (0.040 g, 0.19 mmol) and diethoxymethylacetate (0.16 mL, 0.97 mmol) were combined and dissolved in acetic acid (1 mL). The reaction was warmed to 110 °C and stirred at this temperature for 1 h. The reaction was cooled to room temperature and 15 diethyl ether was added to precipitate a beige solid that was collected by filtration (35 mg, 69% yield). ¹H NMR 400 MHz (DMSO-d₆) δ 11.30 (s, 1H); 8.58 (s, 1H); 8.05 (s, 1H); 7.93 (s, 1H); 4.44 (m, 2H); 4.28 (m, 2H); 1.35 (m, 3H); 1.28 (m, 3H).

20 Monomer 7: 3-Dimethylaminomethylene-6-phenyloxindole

(Procedure J - 2)

6-Phenyloxindole (0.053 mg, 0.25 mmol) and dimethylformamide di-t-butylacetal (0.08 mL, 0.32 mmol) were combined and dissolved in DMF. The reaction mixture was stirred at room temperature for 4 hours. The 25 volatiles were removed in vacuo and the resulting residue was triturated with diethyl ether. The solids were collected by filtration (50 mg, 75% yield). ¹H NMR 400 MHz (DMSO-d₆) mixture of E and Z isomers: δ 10.09 (s, 1H); 10.02 (s, 1H); 7.59-7.51 (m, 4H); 7.45-7.22 (m, 10H); 7.10-7.05 (m, 2H); 6.98 (s, 1H); 6.91 (s, 1H); 3.29 (s, 12H).

Monomer 1: 3-Ethoxymethylene-5-phenyl-7-azaoxindole

Synthesized according to Procedure J -1. ¹H NMR 400 MHz (DMSO-d₆) δ 10.93 (s, 1H); 8.24 (s, 1H); 7.86-7.82 (m, 2H); 7.62-7.56 (m, 2H); 7.44 (m, 2H); 7.34 (m, 1H); 4.40 (m, 2H); 1.35 (m, 3H).

5

Monomer 2: 3-Ethoxymethylene-5-(2-furanyl)-7-azaoxindole

Synthesized according to Procedure J -1. ¹H NMR 400 MHz (DMSO-d₆) δ 10.96 (s, 1H); 8.36 (s, 1H); 7.87-7.84 (m, 2H); 7.72 (s, 1H); 6.87 (d, 1H); 6.56 (m, 1H); 4.42 (m, 2H); 1.36 (m, 3H).

10

Monomer 3: 3-Ethoxymethylene-5-(3-thienyl)-7-azaoxindole

Synthesized according to Procedure J -1. ¹H NMR 400 MHz (DMSO-d₆) δ 10.89 (s, 1H); 8.33 (s, 1H); 7.88 (s, 1H); 7.84 (s, 1H); 7.79 (s, 1H); 7.62 (m, 1H); 7.49 (d, 1H); 4.40 (m, 2H); 1.36 (m, 3H).

15

Monomer 4: 5-Bromo-3-ethoxymethylene-7-azaoxindole

Synthesized according to Procedure J -1. ¹H NMR 400 MHz (DMSO-d₆) δ 11.02 (s, 1H); 8.07 (s, 1H); 7.88 (s, 1H); 7.71 (s, 1H); 4.40 (m, 2H); 1.34 (m, 3H).

20

Monomer 5: 6-Chloro-3-ethoxymethylene-7-azaoxindole

Synthesized according to Procedure J -1. ¹H NMR 400 MHz (DMSO-d₆) δ 11.06 (s, 1H); 7.84 (s, 1H); 7.63 (d, 1H); 6.98 (d, 1H); 4.39 (m, 2H); 1.32 (m, 3H).

25

Monomer 8: 3-Dimethylaminomethylene-6-(2-furanyl)oxindole

Synthesized according to Procedure J -2. ¹H NMR 400 MHz (DMSO-d₆) mixture of E and Z isomers: δ 10.10 (s, 1H); 10.03 (s, 1H); 7.64 (s, 1H); 7.61 (s, 1H); 7.56 (s, 1H); 7.38 (s, 1H); 7.37 (d, 1H); 7.26 (d, 1H); 7.14 (s,

1H); 7.12 (s, 1H); 7.03 (s, 1H); 6.96 (s, 1H); 6.72 (d, 1H); 6.67 (d, 1H); 6.52 (s, 1H); 6.49 (s, 1H); 3.29 (s, 12H)




Monomer 9: 3-Dimethylaminomethylene-7-azaoxindole








- 5 This monomer was generated in situ (during library synthesis) from 7-aza-oxindole and dimethylformamide di-t-butylacetal in DMF.







Monomer 10: 3-Dimethylaminomethylene-4-azaoxindole

- 10 This monomer was generated in situ (during library synthesis) from 4-aza-oxindole and dimethylformamide di-t-butylacetal in DMF.

Representative Amine Monomers are shown in Table 6.

Monomer Code	Structure
A	
B	
C	

Monomer Code	Structure
D	
E	
F	
G	
H	
I	
J	

Monomer Code	Structure
K	
L	
M	
N	
O	
P	

Solution Phase Library Synthesis:

- 5 Stock solutions (0.037M) were prepared for each set of monomers. For example, 35 mg of Monomer 1 was dissolved in 3.5 mL of ethanol. Each monomer was prepared in a separate solution and carefully labeled 1

through 8. For the amine set, a slight excess of stock solution was prepared. For example, 20.4 mg was dissolved in 4.0 mL of methanol. Each amine was prepared in a separate solution and carefully labeled A through P.

Monomer 9 and Monomer 10 were generated *in situ* by preparing
 5 stock solutions of the corresponding aza-oxindole. For example, 20.1 mg of 4-aza-oxindole was dissolved in 4.0 mL of ethanol. Both of the precursors for monomers 9 and 10 were transferred (0.20 mL/well) to a 96-well dry heating block (vwrBRAND Dry Block Heater, cat #13259-066) according to the map below where M represents the monomer and A represents the
 10 amine. The ethanol was evaporated off at 50 °C until it was clear that there was no solvent remaining. DMF (0.20 mL) was added followed by the addition of dimethylformamide di-*t*-butylacetal (0.003 mL) and this remained at room temperature for 1 h.

Monomers 1 through 8 (0.20 mL/well) were transferred to the
 15 appropriate wells in the dry block heater according to the plate map below. After the *in situ* conversion of monomer 9 and 10 was complete, the aniline set (0.20 mL/well) was transferred to the appropriate wells according to the plate map below. The plates were heated to 70 °C for 4 h and then the reaction was cooled to 40 °C and heating was continued for another 16 h.
 20 Ethanol was added as necessary to keep a constant reaction volume in the wells.

Plate 1

M1-AA	M1-AI	M2-AA	M2-AI	M3-AA	M3-AI	M4-AA	M4-AI	M5-AA	M5-AI
M1-AB	M1-AJ	M2-AB	M2-AJ	M3-AB	M3-AJ	M4-AB	M4-AJ	M5-AB	M5-AJ
M1-AC	M1-AK	M2-AC	M2-AK	M3-AC	M3-AK	M4-AC	M4-AK	M5-AC	M5-AK
M1-AD	M1-AL	M2-AD	M2-AL	M3-AD	M3-AL	M4-AD	M4-AL	M5-AD	M5-AL
M1-AE	M1-AM	M2-AE	M2-AM	M3-AE	M3-AM	M4-AE	M4-AM	M5-AE	M5-AM
M1-AF	M1-AN	M2-AF	M2-AN	M3-AF	M3-AN	M4-AF	M4-AN	M5-AF	M5-AN
M1-AG	M1-AO	M2-AG	M2-AO	M3-AG	M3-AO	M4-AG	M4-AO	M5-AG	M5-AO
M1-AH	M1-AP	M2-AH	M2-AP	M3-AH	M3-AP	M4-AH	M4-AP	M5-AH	M5-AP

Plate 2

M6-AA	M6-AI	M7-AA	M7-AI	M8-AA	M8-AI	M9-AA	M9-AI	M10-AA	M10-AI
M6-AB	M6-AJ	M7-AB	M7-AJ	M8-AB	M8-AJ	M9-AB	M9-AJ	M10-AB	M10-AJ
M6-AC	M6-AK	M7-AC	M7-AK	M8-AC	M8-AK	M9-AC	M9-AK	M10-AC	M10-AK
M6-AD	M6-AL	M7-AD	M7-AL	M8-AD	M8-AL	M9-AD	M9-AL	M10-AD	M10-AL
M6-AE	M6-AM	M7-AE	M7-AM	M8-AE	M8-AM	M9-AE	M9-AM	M10-AE	M10-AM
M6-AF	M6-AN	M7-AF	M7-AN	M8-AF	M8-AN	M9-AF	M9-AN	M10-AF	M10-AN
M6-AG	M6-AO	M7-AG	M7-AO	M8-AG	M8-AO	M9-AG	M9-AO	M10-AG	M10-AO
M6-AH	M6-AP	M7-AH	M7-AP	M8-AH	M8-AP	M9-AH	M9-AP	M10-AH	M10-AP

Upon completion of the reaction, methanol (1.0 mL) was added to each well. Using a multi-pipettor, the contents of the reaction wells were transferred to the appropriate wells of a 96-well (Beckmann) plate. The volatiles were removed using a nitrogen flow to substantially reduce the volume of solvent, followed by placing the plates in a vacuum drying oven at 70 °C under 15 mmHg of pressure. The average weight of product determined for plate 1 was 1.91 mg/well (70% conversion). The average weight of product determined for plate 2 was 1.78 mg/mL (70% conversion). All of the wells were analysed by LC-MS.

A Micromass Platform II mass spectrometer equipped with an electrospray ion source was used to acquire low resolution LC-MS data for the samples. The system software runs on a PC computer with the Microsoft operating system, and consists of Masslynx v3.1 and Openlynx v3.1 software packages. The mass spectrometer inlet system was comprised of a Hewlett Packard 1100 HPLC Chromatograph, a Gilson 215 autosampler, and a Hewlett Packard 1100 photo-diode array detector. A Supelco ABZ+ 5cm column was used to provide separations prior to electrospray ionization. The HPLC was programmed as follows:

	Time	%A	%B	Flow Rate
	0.0 min		85	15
	3.0 min		25	75
	4.0 min		0	100
	5.0 min		0	100

The data were processed automatically using standard peak detection parameters provided by the Openlynx software.

A Micromass LCT bench-top mass spectrometer equipped with an electrospray ionization source was used to obtain accurate mass (high resolution) data. The LCT utilizes two hexapole RF lenses to transfer ions from the source to an orthogonal acceleration time-of-flight (TOF) analyser. The ions emerging from the analyser are detected using a dual microchannel plate detector and ion counting system. The system software runs on a PC computer with the Microsoft operating system, and consists of Masslynx v3.2 and Openlynx v3.2 software packages. The mass spectrometer inlet system is comprised of a Waters Alliance 2690 Separations Module, Waters 2700 autosampler, Waters 996 photo-diode array detector and Valco column switching device. A mobile phase flow rate of 1 ml/min exits the Alliance 2690 and is reduced to a mass spectrometer flow rate of 20 μ l/min using an accurate flow splitter. A lock mass solution at a flow rate of 4 μ l/min is added to the spectrometer flow via a Harvard syringe pump and a tee piece placed immediately before the electrospray probe. The instrument resolution was determined by acquiring a spectrum and measuring the full peak width at half peak height (FWHH). The instrument was tuned to provide a resolution of 4600 to 5000 (FWHH). The instrument was calibrated using the ions of polyethylene glycol (PEG) as reference standards. The lock mass used [3,5-Dil-Tyr, Ala,N-Me-Phe, Gly-OI] Enkephalin (MH^+ $C_{26}H_{34}I_2N_5O_6$ = 766.0599) at a concentration of 5 ng/ μ l.

Following are examples of the invention prepared in the solution phase library with accompanying high resolution or low resolution mass spectral data.

Example 73: 3-[[[(Z and E)-(2-Oxo-6-phenyl-1,2-dihydro-3H-indol-3-ylidene)methyl]amino}benzamide. Accurate mass $M + H = 356.1388$ (mmu error 1.0).

Example 74: Diethyl 4-(((Z and E)- (2-oxo-6-phenyl-1,2-dihydro-3H-indol-3-ylidene)methyl)amino)benzylphosphonate. Accurate mass $M + H = 463.1952$ (mmu error 3.4).

Example 75: 3-(((Z and E)-(2-Oxo-6-phenyl-1,2-dihydro-3H-indol-3-ylidene)methyl)amino)benzonitrile. Accurate mass $M + H = 338.1289$ (mmu error 0.3).

Example 76: 3-((Z and E)-[2-(2-Hydroxyethyl)anilino]methylidene)-6-phenyl-1,3-dihydro-2H-indol-2-one. Accurate mass $M + H = 357.1595$ (mmu error 0.7).

Example 77: 3-((Z and E)-[3-(Hydroxymethyl)anilino]methylidene)-6-phenyl-1,3-dihydro-2H-indol-2-one. Accurate mass $M + H = 343.1438$ (mmu error 0.7).

Example 78: 3-((Z and E)-(2-Methoxyanilino)methylidene)-6-phenyl-1,3-dihydro-2H-indol-2-one. Accurate mass $M + H = 343.1442$ (mmu error 0.3).

Example 79: 3-((Z and E)-[4-(5-Methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene)-6-phenyl-1H-indol-2-one. Accurate mass $M + H = 409.1645$ (mmu error 1.9).

Example 80: 3-((Z and E)-(4-Iodoanilino)methylidene)-6-phenyl-1H-indol-2-one. Accurate mass $M + H = 439.0246$ (mmu error 1.9).

Example 81: 3-(((Z and E)-[6-(2-Furyl)-2-oxo-1,2-dihydro-3H-indol-3-ylidene]methyl)amino)benzamide. Accurate mass $M + H = 346.1188$ (mmu error 0.2).

Example 82: Diethyl 4-(((Z and E)-[6-(2-furyl)-2-oxo-1,2-dihydro-3H-indol-3-ylidene]methyl)amino)benzylphosphonate. Accurate mass $M + H = 453.1755$ (mmu error 2.3).

Example 83: 3-(((Z and E)-[6-(2-Furyl)-2-oxo-1,2-dihydro-3H-indol-3-ylidene]methyl)amino)benzonitrile. LC-ES-MS ($M+H$) = 328.

Example 84: 6-(2-Furyl)-3-((Z and E)-[2-(2-hydroxyethyl)anilino]methylidene)-1,3-dihydro-2H-indol-2-one. Accurate mass $M + H = 347.1377$ (mmu error 1.7).

Example 85: 6-(2-Furyl)-3-((Z and E)-[3-(hydroxymethyl)anilino]methylidene)-1,3-dihydro-2H-indol-2-one. Accurate mass $M + H = 333.1227$ (mmu error 1.0).

Example 86: 6-(2-Furyl)-3-((Z and E)-(2-methoxyanilino)methylidene)-1,3-dihydro-2H-indol-2-one. Accurate mass $M + H = 333.1234$ (mmu error 0.4).

Example 87: 6-(2-Furyl)-3-((Z and E)-[4-(5-methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene)-1H-indol-2-one. Accurate mass $M + H = 399.1444$ (mmu error 1.1).

Example 88: 6-(2-Furyl)-3-((Z and E)-(4-iodoanilino)methylidene)-1H-indol-2-one. Accurate mass $M + H = 429.0031$ (mmu error 2.7).

Example 99: 3-(((2-Oxo-5-phenyl-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene)methyl)amino)benzamide. Accurate mass $M + H = 357.1339$ (mmu error 1.2).

Example 100: Diethyl 4-(((2-oxo-5-phenyl-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene)methyl)amino)benzylphosphonate. Accurate mass $M + H = 464.1918$ (mmu error 2.0).

Example 101: 3-(((2-Oxo-5-phenyl-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene)methyl)amino)benzonitrile. Accurate mass $M + H = 339.1237$ (mmu error 0.8).

Example 102: 3-([2-(2-Hydroxyethyl)anilino]methylidene)-5-phenyl-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 358.1534$ (mmu error 2.1).

Example 103: 3-([3-(Hydroxymethyl)anilino]methylidene)-5-phenyl-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 344.1377$ (mmu error 2.1).

Example 104: 3-((2-Methoxyanilino)methylidene)-5-phenyl-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 344.1386$ (mmu error 1.2).

Example 105: 3-[[4-(5-Methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene]-5-phenyl-1H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 410.1597$ (mmu error 1.9).

Example 106: 3-[(4-Iodoanilino)methylidene]-5-phenyl-1H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 440.0218$ (mmu error 0.0).

Example 107: 3-([5-(2-Furyl)-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene]methyl)amino)benzamide. Accurate mass $M + H = 347.1135$ (mmu error 0.8).

Example 108: Diethyl 4-([5-(2-furyl)-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene]methyl)amino)benzylphosphonate. Accurate mass $M + H = 454.1724$ (mmu error 0.7).

Example 109: 3-([5-(2-Furyl)-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene]methyl)amino)benzonitrile. Accurate mass $M + H = 329.1017$ (mmu error 2.1).

Example 110: 5-(2-Furyl)-3-[[2-(2-hydroxyethyl)anilino]methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 348.1333$ (mmu error 1.4).

Example 111: 5-(2-Furyl)-3-[[3-(hydroxymethyl)anilino]methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 334.1168$ (mmu error 2.3).

Example 112: 5-(2-Furyl)-3-[(2-methoxyanilino)methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 334.1177$ (mmu error 1.4).

Example 113: 5-(2-Furyl)-3-[[4-(5-methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene]-1H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 400.1394$ (mmu error 1.5).

Example 114: 5-(2-Furyl)-3-[(4-iodoanilino)methylidene]-1H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 430.0008$ (mmu error 0.3).

Example 115: 3-({[2-Oxo-5-(3-thienyl)-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene]methyl}amino)benzamide. Accurate mass $M + H = 363.0903$ (mmu error 1.2).

Example 116: Diethyl 4-({[2-oxo-5-(3-thienyl)-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene]methyl}amino)benzylphosphonate. Accurate mass $M + H = 470.1497$ (mmu error 0.6).

Example 117: 3-({[2-Oxo-5-(3-thienyl)-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene]methyl}amino)benzonitrile. Accurate mass $M + H = 345.0800$ (mmu error 0.9).

Example 118: 3-([2-(2-Hydroxyethyl)anilino]methylidene)-5-(3-thienyl)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 364.1098$ (mmu error 2.1).

Example 119: 3-([3-(Hydroxymethyl)anilino]methylidene)-5-(3-thienyl)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 350.0959$ (mmu error 0.4).

Example 120: 3-([2-Methoxyanilino]methylidene)-5-(3-thienyl)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 350.0946$ (mmu error 1.7).

Example 121: 3-([4-(5-Methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene)-5-(3-thienyl)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 416.1172$ (mmu error 0.8).

Example 122: 3-([4-Iodoanilino]methylidene)-5-(3-thienyl)-1H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 445.9771$ (mmu error 1.2).

Example 123: 3-({[5-Bromo-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene]methyl}amino)benzamide. Accurate mass $M + H = 359.0134$ (mmu error 0.9).

Example 124: Diethyl 4-({[5-Bromo-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene]methyl}amino)benzylphosphonate. Accurate mass $M + H = 466.0730$ (mmu error 0.0).

Example 125: 3-(((5-Bromo-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene)methyl)amino)benzonitrile. Accurate mass $M + H = 341.0033$ (mmu error 0.4).

Example 126: 5-Bromo-3-([2-(2-hydroxyethyl)anilino]methylidene)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 360.0337$ (mmu error 1.0).

Example 127: 5-Bromo-3-([3-(hydroxymethyl)anilino]methylidene)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 346.0176$ (mmu error 1.4).

Example 128: 5-Bromo-3-([2-methoxyanilino]methylidene)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 346.0169$ (mmu error 2.1).

Example 129: 5-Bromo-3-([4-(5-methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 412.0388$ (mmu error 2.0).

Example 130: 5-Bromo-3-([4-iodoanilino]methylidene)-1H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 441.9002$ (mmu error 0.8).

Example 131: 3-(((6-Chloro-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene)methyl)amino)benzamide. Accurate mass $M + H = 315.0633$ (mmu error 1.5).

Example 132: Diethyl 4-([6-chloro-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene)methyl)amino)benzylphosphonate. Accurate mass $M + H = 422.1234$ (mmu error 0.2).

Example 133: 3-(((6-Chloro-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene)methyl)amino)benzonitrile. Accurate mass $M + H = 297.0532$ (mmu error 1.0).

Example 134: 6-Chloro-3-([2-(2-hydroxyethyl)anilino]methylidene)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 316.0836$ (mmu error 1.6).

Example 135: 6-Chloro-3-[[3-(hydroxymethyl)anilino]methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 302.0683$ (mmu error 1.3).

Example 136: 6-Chloro-3-[(2-methoxyanilino)methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 302.0667$ (mmu error 2.9).

Example 137: 6-Chloro-3-[[4-(5-methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 368.0894$ (mmu error 1.9).

Example 138: 6-Chloro-3-[(4-iodoanilino)methylidene]-1H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 397.9504$ (mmu error 1.2).

Example 139: Ethyl 3-[[3-(aminocarbonyl)anilino]methylidene]-2-oxo-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine-5-carboxylate. Accurate mass $M + H = 353.1242$ (mmu error 0.7).

Example 140: Ethyl 3-[(4-[(diethoxyphosphoryl)methyl]anilino)methylidene]-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridine-5-carboxylate. Accurate mass $M + H = 460.1826$ (mmu error 1.0).

Example 141: Ethyl 3-[(3-cyanoanilino)methylidene]-2-oxo-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine-5-carboxylate. Accurate mass $M + H = 335.1127$ (mmu error 1.6).

Example 142: Ethyl 3-[[2-(2-hydroxyethyl)anilino]methylidene]-2-oxo-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine-5-carboxylate. Accurate mass $M + H = 354.1434$ (mmu error 1.9).

Example 143: Ethyl 3-[[3-(hydroxymethyl)anilino]methylidene]-2-oxo-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine-5-carboxylate. Accurate mass $M + H = 340.1287$ (mmu error 0.9).

Example 144: Ethyl 3-[(2-methoxyanilino)methylidene]-2-oxo-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine-5-carboxylate. Accurate mass $M + H = 340.1285$ (mmu error 1.1).

Example 145: Ethyl 3-[[4-(5-methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene]-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridine-5-carboxylate. Accurate mass $M + H = 406.1499$ (mmu error 1.5).

Example 146: Ethyl 3-[[4-iodoanilino]methylidene]-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridine-5-carboxylate. Accurate mass $M + H = 436.0103$ (mmu error 1.3).

Example 147: 3-[[[2-Oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene)methyl]amino]benzamide. Accurate mass $M + H = 281.1024$ (mmu error 1.4).

Example 148: Diethyl 4-[[[2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene)methyl]amino]benzylphosphonate. Accurate mass $M + H = 388.1607$ (mmu error 1.8).

Example 149: 3-[[[2-Oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene)methyl]amino]benzonitrile. Accurate mass $M + H = 263.0923$ (mmu error 0.9).

Example 150: 3-[[2-(2-Hydroxyethyl)anilino]methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 282.1229$ (mmu error 1.2).

Example 151: 3-[[3-(Hydroxymethyl)anilino]methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 268.1082$ (mmu error 0.3).

Example 152: 3-[[2-Methoxyanilino]methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 268.1082$ (mmu error 0.3).

Example 153: 3-[[4-(5-Methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene]-1H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 334.1301$ (mmu error 0.2).

Example 154: 3-[[4-iodoanilino]methylidene]-1H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 363.9898$ (mmu error 0.7).

Example 155: 3-[[[2-Oxo-1,2-dihydro-3H-pyrrolo[3,2-b]pyridin-3-ylidene)methyl]amino]benzamide. Accurate mass $M + H = 281.1035$ (mmu error 0.3).

Example 156: Diethyl 4-[[[2-oxo-1,2-dihydro-3H-pyrrolo[3,2-b]pyridin-3-ylidene)methyl]amino]benzylphosphonate. LC-ES-MS ($M+H$) = 388.

Example 157: 3-[[[2-Oxo-1,2-dihydro-3H-pyrrolo[3,2-b]pyridin-3-ylidene)methyl]amino]benzonitrile. Accurate mass $M + H = 263.0912$ (mmu error 2.0).

Example 158: 3-[[2-(2-Hydroxyethyl)anilino]methylidene]-1,3-dihydro-2H-pyrrolo[3,2-b]pyridin-2-one. Accurate mass $M + H = 282.1241$ (mmu error 0.0).

Example 159: 3-[[3-(Hydroxymethyl)anilino]methylidene]-1,3-dihydro-2H-pyrrolo[3,2-b]pyridin-2-one. Accurate mass $M + H = 268.1079$ (mmu error 0.6).

Example 160: 3-[[2-Methoxyanilino]methylidene]-1,3-dihydro-2H-pyrrolo[3,2-b]pyridin-2-one. Accurate mass $M + H = 268.1083$ (mmu error 0.2).

Example 161: 3-[[4-(5-Methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene]-1H-pyrrolo[3,2-b]pyridin-2-one. Accurate mass $M + H = 334.1301$ (mmu error 0.2).

Example 162: 3-[[4-Iodoanilino]methylidene]-1H-pyrrolo[3,2-b]pyridin-2-one. Accurate mass $M + H = 363.9896$ (mmu error 0.9).

Example 163: 3-[(Z and E)-[(1-methyl-3-phenyl-1H-pyrazol-5-yl)amino]methylidene]-5-phenyl-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 394.1645$ (mmu error 2.2).

Example 164: 3-[(Z and E)-[(4-Hydroxy-1-naphthyl)amino]methylidene]-5-phenyl-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 380.1388$ (mmu error 1.0).

Example 165: 3-[(Z and E)-[(1-Ethyl-1H-pyrazol-5-yl)amino]methylidene]-5-phenyl-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 332.1508$ (mmu error 0.3).

- Example 166:** 3-[(Z and E)-(1H-Benzimidazol-2-ylamino)methylidene]-5-phenyl-1H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 354.1336$ (mmu error 1.8).
- Example 167:** 3-[(Z and E)-{[6-(4-Morpholinyl)-3-pyridinyl]amino}methylidene]-5-phenyl-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 400.1754$ (mmu error 1.9).
- Example 168:** 5-(2-Furyl)-3-[(Z and E)-{(1-methyl-3-phenyl-1H-pyrazol-5-yl)amino}methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 384.1455$ (mmu error 0.5).
- Example 169:** 5-(2-Furyl)-3-[(Z and E)-{(4-hydroxy-1-naphthyl)amino}methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 370.1189$ (mmu error 0.2).
- Example 170:** 3-[(Z and E)-{(1-Ethyl-1H-pyrazol-5-yl)amino}methylidene]-5-(2-furyl)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 322.1297$ (mmu error 0.6).
- Example 171:** 3-[(Z and E)-(1H-Benzimidazol-2-ylamino)methylidene]-5-(2-furyl)-1H-pyrrolo[2,3-b]pyridin-2-one. LC-ES-MS ($M+H$) = 344.
- Example 172:** 5-(2-Furyl)-3-[(Z and E)-{[6-(4-morpholinyl)-3-pyridinyl]amino}methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 390.1556$ (mmu error 0.9).
- Example 173:** 3-[(Z and E)-{(1-Methyl-3-phenyl-1H-pyrazol-5-yl)amino}methylidene]-5-(3-thienyl)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 400.1231$ (mmu error 0.0).
- Example 174:** 3-[(Z and E)-{(4-Hydroxy-1-naphthyl)amino}methylidene]-5-(3-thienyl)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 386.0960$ (mmu error 0.3).
- Example 175:** 3-[(Z and E)-{(1-Ethyl-1H-pyrazol-5-yl)amino}methylidene]-5-(3-thienyl)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 338.1060$ (mmu error 1.5).

Example 176: 3-[(Z and E)-(1H-Benzimidazol-2-ylamino)methylidene]-5-(3-thienyl)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass M + H = 360.0915 (mmu error 0.3).

Example 177: 3-[(Z and E)-{[6-(4-Morpholinyl)-3-pyridinyl]amino}methylidene]-5-(3-thienyl)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass M + H = 406.1321 (mmu error 1.6).

Example 178: 5-Bromo-3-[(Z and E)-[(1-methyl-3-phenyl-1H-pyrazol-5-yl)amino]methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass M + H = 396.0442 (mmu error 1.7).

Example 179: 5-Bromo-3-[(Z and E)-[(4-hydroxy-1-naphthyl)amino]methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass M + H = 382.0187 (mmu error 0.3).

Example 180: 5-Bromo-3-[(Z and E)-[(1-ethyl-1H-pyrazol-5-yl)amino]methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass M + H = 334.0297 (mmu error 0.5).

Example 181: 3-[(Z and E)-(1H-Benzimidazol-2-ylamino)methylidene]-5-bromo-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass M + H = 356.0134 (mmu error 1.2).

Example 182: 5-Bromo-3-[(Z and E)-{[6-(4-morpholinyl)-3-pyridinyl]amino}methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass M + H = 402.0556 (mmu error 0.8).

Example 183: 6-Chloro-3-[(Z and E)-[(1-methyl-3-phenyl-1H-pyrazol-5-yl)amino]methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass M + H = 352.0957 (mmu error 0.7).

Example 184: 6-Chloro-3-[(Z and E)-[(4-hydroxy-1-naphthyl)amino]methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass M + H = 338.0669 (mmu error 2.7).

Example 185: 6-Chloro-3-[(Z and E)-[(1-ethyl-1H-pyrazol-5-yl)amino]methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass M + H = 290.0808 (mmu error 0.0).

Example 186: 3-[(Z and E)-(1H-Benzimidazol-2-ylamino)methylidene]-6-chloro-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 312.0633$ (mmu error 1.8).

Example 187: 6-Chloro-3-[(Z and E)-[6-(4-morpholinyl)-3-pyridinyl]amino]methylidene)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 358.1053$ (mmu error 1.7).

Example 188: Ethyl 3-[(Z and E)-[(1-methyl-3-phenyl-1H-pyrazol-5-yl)amino]methylidene]-2-oxo-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine-5-carboxylate. Accurate mass $M + H = 390.1556$ (mmu error 0.9).

Example 189: Ethyl 3-[(Z and E)-[(4-hydroxy-1-naphthyl)amino]methylidene]-2-oxo-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine-5-carboxylate. Accurate mass $M + H = 376.1289$ (mmu error 0.7).

Example 190: Ethyl 3-[(Z and E)-[(1-ethyl-1H-pyrazol-5-yl)amino]methylidene]-2-oxo-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine-5-carboxylate. Accurate mass $M + H = 328.1403$ (mmu error 0.5).

Example 191: Ethyl 3-[(Z and E)-(1H-benzimidazol-2-ylamino)methylidene]-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridine-5-carboxylate. Accurate mass $M + H = 350.1241$ (mmu error 1.1).

Example 192: Ethyl 3-[(Z and E)-[6-(4-morpholinyl)-3-pyridinyl]amino]methylidene)-2-oxo-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine-5-carboxylate. Accurate mass $M + H = 396.1655$ (mmu error 1.6).

Example 193: 3-[(Z and E)-[(1-Methyl-3-phenyl-1H-pyrazol-5-yl)amino]methylidene]-6-phenyl-1,3-dihydro-2H-indol-2-one. Accurate mass $M + H = 393.1691$ (mmu error 2.3).

Example 194: 3-[(Z and E)-[(4-Hydroxy-1-naphthyl)amino]methylidene]-6-phenyl-1,3-dihydro-2H-indol-2-one. Accurate mass $M + H = 379.1430$ (mmu error 1.5).

Example 195: 3-[(Z and E)-[(1-Ethyl-1H-pyrazol-5-yl)amino]methylidene]-6-phenyl-1,3-dihydro-2H-indol-2-one. Accurate mass $M + H = 331.1552$ (mmu error 0.6).

Example 196: 3-[(Z and E)-(1H-Benzimidazol-2-ylamino)methylidene]-6-phenyl-1H-indol-2-one. Accurate mass $M + H = 353.1388$ (mmu error 1.4).

Example 197: 3-[(Z and E)-{[6-(4-Morpholinyl)-3-pyridinyl]amino}methylidene]-6-phenyl-1,3-dihydro-2H-indol-2-one. Accurate mass $M + H =$
5 399.1809 (mmu error 1.1).

Example 198: 6-(2-Furyl)-3-[(Z and E)-{(1-methyl-3-phenyl-1H-pyrazol-5-yl)amino}methylidene]-1,3-dihydro-2H-indol-2-one. Accurate mass $M + H =$
383.1498 (mmu error 0.9).

Example 199: 6-(2-Furyl)-3-[(Z and E)-{(4-hydroxy-1-naphthyl)amino}methylidene]-1,3-dihydro-2H-indol-2-one.
10

Example 200: 3-[(Z and E)-{(1-Ethyl-1H-pyrazol-5-yl)amino}methylidene]-6-(2-furyl)-1,3-dihydro-2H-indol-2-one. Accurate mass $M + H = 321.1342$ (mmu error 0.8).

Example 201: 3-[(Z and E)-(1H-Benzimidazol-2-ylamino)methylidene]-6-(2-furyl)-1H-indol-2-one. Accurate mass $M + H = 343.1192$ (mmu error 0.2).
15

Example 202: 6-(2-Furyl)-3-[(Z and E)-{[6-(4-morpholinyl)-3-pyridinyl]amino}methylidene]-1,3-dihydro-2H-indol-2-one. Accurate mass $M + H = 389.1604$ (mmu error 0.8).

Example 203: 3-[(Z and E)-{(1-Methyl-3-phenyl-1H-pyrazol-5-yl)amino}methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M +$
20 $H = 318.1347$ (mmu error 0.6).

Example 204: 3-[(Z and E)-{(4-Hydroxy-1-naphthyl)amino}methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one.

Example 205: 3-[(Z and E)-{(1-Ethyl-1H-pyrazol-5-yl)amino}methylidene]-
25 1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass $M + H = 256.1195$ (mmu error 0.2).

Example 206: 3-[(Z and E)-(1H-Benzimidazol-2-ylamino)methylidene]-1H-pyrrolo[2,3-b]pyridin-2-one.

Example 207: 3-((Z and E)-{[6-(4-Morpholinyl)-3-pyridinyl]amino} methylidene)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one. Accurate mass M + H = 324.1453 (mmu error 0.7).

Example 208: 3-((Z and E)-[(1-Methyl-3-phenyl-1H-pyrazol-5-yl)amino] methylidene)-1,3-dihydro-2H-pyrrolo[3,2-b]pyridin-2-one. Accurate mass M + H = 318.1347 (mmu error 0.7).

Example 209: 3-((Z and E)-[(4-Hydroxy-1-naphthyl)amino]methylidene)-1,3-dihydro-2H-pyrrolo[3,2-b]pyridin-2-one.

Example 210: 3-((Z and E)-[(1-Ethyl-1H-pyrazol-5-yl)amino]methylidene)-1,3-dihydro-2H-pyrrolo[3,2-b]pyridin-2-one. Accurate mass M + H = 256.1178 (mmu error 1.9).

Example 211: 3-((Z and E)-(1H-Benzimidazol-2-ylamino)methylidene)-1H-pyrrolo[3,2-b]pyridin-2-one.

Example 212: 3-((Z and E)-{[6-(4-Morpholinyl)-3-pyridinyl]amino} methylidene)-1,3-dihydro-2H-pyrrolo[3,2-b]pyridin-2-one. Accurate mass M + H = 324.1449 (mmu error 1.0).

PHARMACEUTICAL FORMULATION AND DOSES

The compounds of the present invention can be administered in such oral (including buccal and sublingual) dosage forms as tablets, capsules (each including timed release and sustained release formulations), pills, powders, granules, elixirs, tinctures, suspensions, syrups and emulsions. Likewise, they may also be administered in nasal, ophthalmic, otic, rectal, topical, intravenous (both bolus and infusion), intraperitoneal, intraarticular, subcutaneous or intramuscular inhalation or insufflation form, all using forms well known to those of ordinary skill in the pharmaceutical arts.

The dosage regimen utilizing the compounds of the present invention is selected in accordance with a variety of factors including type, species, age, weight, sex and medical condition of the patient; the severity of the

condition to be treated; the route of administration; the renal and hepatic function of the patient; and the particular compound or salt thereof employed. An ordinarily skilled physician or veterinarian can readily determine and prescribe the effective amount of the drug required to
5 prevent, counter or arrest the progress of the condition.

Oral dosages of the present invention, when used for the indicated effects, will range between about 0.1 to about 100 mg/kg of body weight per day, and particularly about 1 to 10 mg/kg of body weight per day. Oral dosage units will generally be administered in the range of from 1 to about
10 250 mg and more preferably from about 25 to about 250 mg. The daily dosage for a 70 kg mammal will generally be in the range of about 70 mg to 7 grams of a compound of formula I or II.

The dosage to be administered is based on the usual conditions such as the physical condition of the patient, age, body weight, past medical
15 history, route of administrations, severity of the conditions and the like. Oral administration is generally preferred for administration to a human. In some cases, a relatively lower dose is sufficient and, in some cases, a relatively higher dose or increased number of doses may be necessary. Topical application similarly may be once or more than once per day depending
20 upon the usual medical considerations. Advantageously, compounds of the present invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three or four times daily. The compounds of the invention can be prepared in a range of concentrations for topical use of about 0.5 to about 5mg/ml of suitable
25 solvent. A preferred volume for application to the scalp is about 2 ml, resulting in an effective dosage delivered to the patient of about 1 to about 10mg.

For treatment of chemotherapy-induced alopecia, administration 1 to
2 times prior to chemotherapy administration is preferred, with additional
30 applications administered as needed. A similar regimen can be pursued for

treatment of alopecia induced by radiation therapy. Furthermore, preferred compounds for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal routes, using those forms of transdermal skin patches well known to those of
5 ordinary skill in that art. To be administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

In the methods of the present invention, the compounds herein described in detail can form the active ingredient and are typically
10 administered in admixture with suitable pharmaceutical diluents, excipients or carriers (collectively referred to herein as "carrier" materials) suitably selected with respect to the intended form of administration, that is, oral tablets, capsules, elixirs, syrups and the like, and consistent with conventional pharmaceutical practices.

15 For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic pharmaceutically acceptable inert carrier such as ethanol, glycerol, water and the like. Powders are prepared by comminuting the compound to a suitable fine size and mixing with a similarly comminuted pharmaceutical
20 carrier such as an edible carbohydrate, as, for example, starch or mannitol. Flavoring, preservative, dispersing and coloring agent can also be present.

Capsules are made by preparing a powder mixture as described above, and filling formed gelatin sheaths. Glidants and lubricants such as colloidal silica, talc, magnesium stearate, calcium stearate or solid
25 polyethylene glycol can be added to the powder mixture before the filling operation. A disintegrating or solubilizing agent such as agar-agar, calcium carbonate or sodium carbonate can also be added to improve the availability of the medicament when the capsule is ingested.

Moreover, when desired or necessary, suitable binders, lubricants,
30 disintegrating agents and coloring agents can also be incorporated into the

mixture. Suitable binders include starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium alginate, carboxymethylcellulose, polyethylene glycol, waxes and the like. Lubricants used in these dosage forms include sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum and the like. Tablets are formulated, for example, by preparing a powder mixture, granulating or slugging, adding a lubricant and disintegrant and pressing into tablets. A powder mixture is prepared by mixing the compound, suitably comminuted, with a diluent or base as described above, and optionally, with a binder such as carboxymethylcellulose, an aliginate, gelatin, or polyvinyl pyrrolidone, a solution retardant such as paraffin, a resorption accelerator such as a quaternary salt and/or an absorption agent such as bentonite, kaolin or dicalcium phosphate. The powder mixture can be granulated by wetting with a binder such as syrup, starch paste, acadia mucilage or solutions of cellulosic or polymeric materials and forcing through a screen. As an alternative to granulating, the powder mixture can be run through the tablet machine and the result is imperfectly formed slugs broken into granules. The granules can be lubricated to prevent sticking to the tablet forming dies by means of the addition of stearic acid, a stearate salt, talc or mineral oil. The lubricated mixture is then compressed into tablets. The compounds of the present invention can also be combined with free flowing inert carrier and compressed into tablets directly without going through the granulating or slugging steps. A clear or opaque protective coating consisting of a sealing coat of shellac, a coating of sugar or polymeric material and a polish coating of wax can be provided. Dyestuffs can be added to these coatings to distinguish different unit dosages.

Oral fluids such as solution, syrups and elixirs can be prepared in dosage unit form so that a given quantity contains a predetermined amount

of the compound. Syrups can be prepared by dissolving the compound in a suitably flavored aqueous solution, while elixirs are prepared through the use of a non-toxic alcoholic vehicle. Suspensions can be formulated by dispersing the compound in a non-toxic vehicle. Solubilizers and emulsifiers
5 such as ethoxylated isostearyl alcohols and polyoxy ethylene sorbitol ethers, preservatives, flavor additive such as peppermint oil or saccharin, and the like can also be added.

Where appropriate, dosage unit formulations for oral administration can be microencapsulated. The formulation can also be prepared to prolong
10 or sustain the release as for Example by coating or embedding particulate material in polymers, wax or the like.

The compounds of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be
15 formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines.

Compounds of the present invention may also be delivered by the use of monoclonal antibodies as individual carriers to which the compound molecules are coupled. The compounds of the present invention may also
20 be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamide-phenol, polyhydroxyethylaspartamidephenol, or polyethyleneoxidepolylysine substituted with palmitoyl residues. Furthermore, the compounds of the
25 present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid, polepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacrylates and cross-linked or amphipathic block copolymers of hydrogels.

The present invention includes pharmaceutical compositions containing about 0.01 to about 99.5%, more particularly, about 0.5 to about 90% of a compound of the formula (II) in combination with a pharmaceutically acceptable carrier.

- 5 Parenteral administration can be effected by utilizing liquid dosage unit forms such as sterile solutions and suspensions intended for subcutaneous, intramuscular or intravenous injection. These are prepared by suspending or dissolving a measured amount of the compound in a non-toxic liquid vehicle suitable for injection such as aqueous oleaginous medium
10 and sterilizing the suspension or solution.

Alternatively, a measured amount of the compound is placed in a vial and the vial and its contents are sterilized and sealed. An accompanying vial or vehicle can be provided for mixing prior to administration. Non-toxic salts and salt solutions can be added to render the injection isotonic.
15 Stabilizers, preservations and emulsifiers can also be added.

Rectal administration can be effected utilizing suppositories in which the compound is admixed with low-melting water-soluble or insoluble solids such as polyethylene glycol, cocoa butter, higher ester as for Example flavored aqueous solution, while elixirs are prepared through
20 myristyl palmitate or mixtures thereof.

Topical formulations of the present invention may be presented as, for instance, ointments, creams or lotions, eye ointments and eye or ear drops, impregnated dressings and aerosols, and may contain appropriate conventional additives such as preservatives, solvents to assist drug
25 penetration and emollients in ointments and creams. The formulations may also contain compatible conventional carriers, such as cream or ointment bases and ethanol or oleyl alcohol for lotions. Such carriers may be present as from about 1% up to about 98% of the formulation. More usually they will form up to about 80% of the formulation.

For administration by inhalation the compounds according to the invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant, e.g. dichlorodifluoromethane, trichlorofluoromethane, 5 dichlorotetrafluoroethane, tetrafluoroethane, heptafluoropropane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of e.g. gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of a compound of the 10 invention and a suitable powder base such as lactose or starch.

The preferred pharmaceutical compositions are those in a form suitable for oral administration, such as tablets and liquids and the like and topical formulations.

15

BIOLOGICAL DATA

The compounds of the present invention have valuable pharmacologic properties. Different compounds from this class are particularly effective at inhibiting the VEGF-R2 receptor kinase enzyme at concentrations which range from 0.0001 to 1 μ M and additionally show 20 selectivity relative to other kinases. Substrate phosphorylation assays were carried out as follows:

VEGFR-2

The peptide substrate used in the VEGFR-2 assay was biotin- 25 aminohexyl-EEEEYFELVAKKKK-NH₂. The kinase domain of the enzyme was purified to homogeneity from a baculovirus expression system. The enzyme was preactivated on ice for 15 min in the presence of 100 μ M ATP and 20 mM MgCl₂, and stored at -80°C until needed for assay. The activated enzyme was diluted to 0.4 nM into a 60 μ l reaction containing 30 mM HEPES, pH 7.5, 5 μ M ATP, 10 mM MgCl₂, 5 μ M peptide, 0.1 mM DTT,

0.05 mg/ml BSA, and an inhibitor at varying concentrations. The controls were reactions in the presence (negative controls) or absence (positive controls) of 50 mM EDTA. Reactions were incubated for 30 min at room temperature, and then quenched by the addition of EDTA to 60 mM in 210
5 μ l. The quenched samples (190 μ l) were transferred to a neutravidin-coated plate (Pierce) and incubated at room temperature for 40 min to allow biotinylated peptide to bind to the neutravidin. The unbound components of the reaction were removed by washing with a plate washer, then 200 μ l HRP-PY20 anti-phosphotyrosine antibody conjugate was added to each
10 well. After incubation for 40 min, the plate was washed to remove any unbound antibody. A HRP substrate, K-blue (Neogen) was added and the reaction was quenched with Red Stop (Neogen) after 20 min. The absorbance of the wells was read at A_{650} in a plate reader. IC_{50} values were obtained by fitting raw data to $A_{650} = V_{MAX} * (1 - [I]/IC_{50} + [I])) + b$, where b is
15 background.

CDK1 and CDK2

Cyclin dependent protein kinase assays utilized the peptides Biotin-aminohexyl-AAKAKKTPKKAKK and Biotin-aminohexyl-ARRPMSPKKKA-
20 NH_2 as phosphoryl group acceptors. CDK1 and CDK2 were both expressed utilizing a baculovirus expression system and were partially purified to comprise 20-80% of total protein, with no detectable competing reactions present. Typically, assays were performed by incubating either enzyme (0.2-10 nM), with and without inhibitor, one of the two peptide substrates (1-
25 10 nM), $[\gamma-^{32}P]ATP$ (1-20 nM), and 10-20 mM Mg^{2+} for periods of time generally within the range 10-120 min. Reactions were terminated with 0.2-2 volumes of either 20% acetic acid or 50-100 mM EDTA buffered to pH 7 (substrate consumption < 20%). The buffer employed in enzyme assays was either 30 mM HEPES 7.4 containing 0.15 M NaCl and 5% DMSO, the
30 buffer 50 mM MOPS 7.0 containing 0.15 M NaCl and 5% DMSO, or the

buffer 100 mM HEPES pH 7.5 containing 0.1 mg/mL BSA and 5% DMSO. Inhibitors were diluted in 100% DMSO prior to addition into the assay. Detection of peptide phosphorylation was accomplished by scintillation counting following either collection of peptide onto phosphocellulose filters
5 (for reactions stopped with acetic acid), collection of peptide in wells of 96 well plates coated with Streptavidin (Pierce) (reactions were stopped with EDTA), or addition of Avidin coated Scintillant impregnated beads (Scintillation Proximity Assays from Amersham, reactions were stopped with EDTA). Counts detected by any of these methodologies minus the
10 appropriate background (assays with additional 40mM EDTA or lacking peptide substrate) were assumed to be proportional to the reaction initial rates, and IC50s were determined by a least squares fit to the equation $CPM = V_{max} * (1 - ([I]/(K + [I]))) + nsb$, or pIC50s were determined by a fit to the equation $CPM = nsb + (V_{max} - nsb) / (1 + (x/10^x - pIC50))$, where nsb are the
15 background counts.

Tie-2

The peptide substrate used in the Tie-2 assay was biotin-aminohexyl-LEAREYRWLGKKKamide. The kinase domain of the enzyme was purified
20 to homogeneity from a baculovirus expression system. The enzyme was diluted to 10 nM into a 60 µl reaction containing 100 mM HEPES, pH 7.5, 500 µM ATP, 10 mM MgCl₂, 2 µM peptide, 1 mM DTT, 0.05 mg/ml BSA, and an inhibitor at varying concentrations. The controls were reactions in the presence (negative controls) or absence (positive controls) of 50 mM EDTA.
25 Reactions were incubated for 30 min at room temperature, and then quenched by stopped by 80 µl of 0.15 M EDTA. The quenched samples (125 µl) were transferred to a Neutravidin plates # 15128 and incubated at room temperature for 30-60 minutes, allowing the biotinylated peptide to bind to the neutravidin on the plates. The neutravidin plates were then washed
30 with water for 5 times. Europium conjugated anti-phosphotyrosine antibody,

(EG & G Wallac, # CR04-100) (1 mg/ml) was diluted 1:10,000 in 1%BSA-0.05% Tween 20-TBS, and 150 μ l of the diluted antibody was added to each well of the neutravidin plate, so the phosphorylated peptide was bound with the Europium labelled antibody. After another 30-60 min incubation at room temperature, the plates were washed again with water for 5 times. 150 μ l of Enhancemant solution was then added to each well, dissociating Eu^{3+} from solid phase bound antibodies to form a homogeneous and highly fluorescent $\text{Eu}-(2\text{-NTA})_3(\text{TOPO})_{2-3}$ micellar chelate solution. The plates were incubated for 10 minutes at room temperature to allow the above process, and fluorescent signal for each well was determined in a Wallac 1420 Victor Multilabel Counter with "Europium" protocol.

The kinase activity of all wells was calculated as %S, the percentage of the fluorescent counts vs. positive controls after subtraction of negative controls, as in eq. 1.

15

$$\% S = 100 * \frac{\text{Counts}_{\text{sample}} - \text{Counts}_{\text{negative}}}{\text{Counts}_{\text{positive}} - \text{Counts}_{\text{negative}}} \quad (1)$$

Plots of compound concentration versus %S were constructed. IC50s (K, expressed in units of molarity), the compound concentration at which the enzyme activity was inhibited by 50%, were determined from nonlinear least squares fits of the data to the simple competitive binding model of eq. 2.

$$\%S = \%S_{\text{max}} * (1 - (X / (K + X))) + Y_2 \quad (2)$$

Where %S is the experimentally observed count rate at sample compound concentration X, $\%S_{\text{max}}$ is the best fit value for the maximum amplitude of the concentration-response curve, Y_2 is the count rate observed at infinitely high inhibitor concentration.

c-Fms

c-fms protein kinase assays utilized the peptide substrate, biotin-EAIYAAPFAKKK-NH₂, as the phosphoryl group acceptor. The c-fms intracellular domain was expressed from a baculovirus expression system, as an amino-terminal GST fusion protein, and purified to homogeneity using Glutathione agarose from Sigma Chemical Co.. Maximum activation of the enzyme was achieved by preactivation at room temperature for 120 min in the presence of 100 μ M ATP and 15 mM MgCl₂. This enzyme stock was diluted to 150 nM prior to using in the assay. Typically assays were performed in white, opaque, 96-well plates in a 45ul assay volume including 15ul 6% DMSO, with or without compounds, 15ul of the preactivated, diluted enzyme, and 15ul of a substrate mixture. Reactions contained 50 mM HEPES, pH 7.5, 1.7 μ M ATP, 15 mM MgCl₂, 3 μ M peptide, 2.5 mM DTT, 50mM NaCl and 0.15uCi/assay [³²P]ATP. The controls were reactions in the presence (negative controls) or absence (positive controls) of 50 mM EDTA. Reactions were allowed to proceed for 90 min at room temperature. The reaction products were quantified using Scintillation Proximity technology. The reactions were quenched by the addition of 200ul of a solution containing 0.3mg streptavidin SPA beads from Amersham, 50 mM EDTA, 0.1% TX-100, 50uM ATP, in PBS, pH7.2 (phosphate buffered solution). Plates were sealed and counted in a Packard Topcount scintillation counter. IC50 values were obtained by fitting raw data to the equation $y=V_{max}*(1-(x/(k+x)))$.

The results shown in Table 7 summarise representative data: Table 7 illustrates the inhibitory activity of compounds of the present invention against several different kinases (VEGFR2, CDK2, Tie-2, and c-fms).

Table 7. Kinase inhibition data of representative compounds

Compound	VEGFR2	CDK2	Tie-2	c-fms
Example 24	+++	++	+	+
Example 29	+++	+	+	+
Example 30	+++	+++	+	+

Compound	VEGFR2	CDK2	Tie-2	c-fms
Example 32	+++	++	+	+
Example 48	++++	+++	+	+
Example 55	++++	+++	+	+
Example 58	+++	+++	+	
Example 63	++++		+	
Example 66	++++		+	
Example 71	+++		+	
Example 72	++++		+	

Key (IC₅₀, nM)

1-10: +++++

11-50: +++

5 51-100: ++

>100: +

VEGF-R2 receptor kinase mediates intracellular signal transduction produced by extracellular receptor binding of Vascular Endothelial Growth Factor (VEGF) in the endothelial cells involved in angiogenesis. An inhibitor of VEGF-R2 receptor kinase can therefore block mitogenesis in endothelial cells, and will show selectivity (greater inhibition) compared to effects of other growth factors. An assay to measure this effect involves measurement of the inhibition of bromo-deoxy uridine (BrdU) incorporation (evidence of DNA synthesis) in human umbilical vein endothelial cells (HUVECs). Selectivity is demonstrated by comparison to the inhibition of BrdU incorporation prompted by basic Fibroblast Growth Factor (bFGF). These assays are described below.

20 HUVEC BrdU incorporation

Materials

HUVEC cells and EGM (Endothelial cell growth medium) were purchased from Clonetics (San Diego, CA). VEGF and bFGF were purchased from Genzyme. M199 medium (Gibco BRL) with BSA (Sigma) was used for cell culture in Type I collagen coated plates (Becton Dickinson).

The degree of BrdU incorporation was measured using a commercial colorimetric ELISA assay from Boehringer Mannheim.

Method

HUVECs were plated at a density of 2500 cells per well in M199 medium containing 5% FBS (Hyclone) in a Type-1 Collagen coated plate. The plate was incubated at 37°C overnight. The medium was removed by aspiration, and test compounds were added to each well in a volume of 0.1 ml per well in serum-free M199 medium. Compound concentrations ranged from 0.15 nM to 3.0 micromolar. The plate was incubated for 30 min at 37 °C. Another 0.1 ml of serum-free M199 medium containing BSA and VEGF (or bFGF) was added to give a final concentration of 0.1% BSA and 20 ng/ml VEGF (0.3 ng/ml bFGF). The plate was incubated at 37°C for 72 hrs. BrdU was added to each well after the first 48 hrs to give a concentration of 10 micromolar. The colorimetric ELISA assay was performed according to manufacturer's instructions, with detection by absorbance reading at 450 nm. Results were plotted as concentration of test compound vs. absorbance to give an IC₅₀ value for inhibition of BrdU incorporation.

Values of IC₅₀ inhibition for VEGF and bFGF by representative compounds are given in Table 8.

20

Table 8. Inhibition of BrdU incorporation in HUVEC cells		
Compound	VEGF stimulation	bFGF stimulation
Example 24	++++	+
Example 29	++++	+
Example 30	++++	++
Example 32	++++	++
Example 48	++++	+++
Example 55	++	+
Example 58	++++	++
Example 63	++++	+++
Example 66	+++	+
Example 71	++++	++++
Example 72	++++	++++

5	Key (IC ₅₀ , μ M)	0.01-0.5: +++++
		0.6-1.0: +++
		1.1-5.0: ++
		>5.0: +

UTILITY OF INVENTION

Inhibitors of VEGF-R2 kinase have utility as agents in the treatment of a wide variety of disorders which have a proliferative component dependent upon angiogenesis. These include, for example, cancers, arthritis, diabetic retinopathy, macular degeneration, and psoriasis.

The tumour inhibitory activity of the compounds of the present invention can be demonstrated in vivo, using Swiss Nu/Nu female mice in which mouse or human tumor cell lines have been implanted subcutaneously. Typical tumor types are the murine Lewis lung tumor and the human HT-29 colon carcinoma. In this assay, the compounds induce a marked reduction in the average tumour volume compared to vehicle treated controls.

The present invention demonstrates methodologies by which the process of angiogenesis induced by abnormal release of Vascular Endothelial Growth Factor (VEGF) may be inhibited by treatment with inhibitors of VEGF-R2 receptor tyrosine kinase. Proliferative disorders and other disorders involving abnormal angiogenesis can therefore be treated.

When the compounds of the present invention are used in conjunction with chemotherapeutic agents or radiation therapy for cancer treatment, they provide a secondary means of suppressing tumor growth either when administered simultaneously with chemotherapeutic agents, or in an alternating regimen to suppress tumor growth between chemotherapeutic or radiation treatments.

While the invention has been described and illustrated with reference to certain preferred embodiments thereof, those skilled in the art will appreciate that various changes, modifications and substitutions can be


made therein without departing from the spirit and scope of the invention. For example, effective dosages other than the preferred dosages as set forth herein above may be applicable as a consequence of variations in the responsiveness of the mammal being treated for cancer conditions, or for
5 other indications for the compounds of the invention as indicated above. Likewise, the specific pharmacologic responses observed may vary according to and depending upon the particular active compound selected or whether there are present certain pharmaceutical carriers, as well as the type of formulation and mode of administration employed, and such
10 expected variations or differences in the results are contemplated in accordance with the objects and practices of the present invention. It is intended, therefore, that the invention be limited only by the scope of the claims which follow and that such claims be interpreted as broadly as is reasonable.

15

The application of which this description and claim(s) forms part may be used as a basis for priority in respect of any subsequent application. The claims of such subsequent application may be directed to any feature or combination of features described herein. They may take the form of
20 product, formulation, process or use claims and may include, by way of example and without limitation, one or more of the following claim(s):

Claims

1. A compound of the formula:



5

wherein:

- Y, Z, A, and D are independently selected from the group consisting of: carbon and nitrogen, with the provisos that: (1) Z and D may be nitrogen, but otherwise no more than one of Y, Z, A, and D may be nitrogen, and (2)
- 10 when Y, Z, or A are nitrogen, substituent R¹, R², or R³ designated for the respective nitrogen atom is non-existent;

X is selected from the group consisting of: N, CH, CCF₃, and C(C₁₋₁₂ aliphatic);

- R¹ is selected from the group consisting of: hydrogen, C₁₋₁₂ aliphatic, thiol, hydroxy, hydroxy-C₁₋₁₂ aliphatic, Aryl, Aryl-C₁₋₁₂ aliphatic, R⁹-Aryl-C₁₋₁₂ aliphatic, Cyc, Cyc-C₁₋₆ aliphatic, Het, Het-C₁₋₁₂ aliphatic, C₁₋₁₂ alkoxy, Aryloxy, amino, C₁₋₁₂ aliphatic amino, di-C₁₋₁₂ aliphatic amino, di-C₁₋₁₂ aliphatic aminocarbonyl, di-C₁₋₁₂ aliphatic aminosulfonyl, C₁₋₁₂ alkoxy carbonyl, fluoro, bromo, iodo, cyano, sulfonamide, or nitro, where R⁹,
- 15
- 20 Aryl, Cyc and Het are as defined below;

R² is selected from the group consisting of: hydrogen, C₁₋₁₂ aliphatic, N-hydroxyimino-C₁₋₁₂ aliphatic, C₁₋₁₂ alkoxy, hydroxy-C₁₋₁₂ aliphatic, C₁₋₁₂ alkoxy carbonyl, carboxyl C₁₋₁₂ aliphatic, Aryl, R⁹-Aryl-oxycarbonyl, R⁹-

- oxycarbonyl-Aryl, Het, aminocarbonyl, C₁₋₁₂ aliphatic-aminocarbonyl, Aryl-C₁₋₁₂ aliphatic-aminocarbonyl, R⁹-Aryl-C₁₋₁₂ aliphatic-aminocarbonyl, Het-C₁₋₁₂ aliphatic-aminocarbonyl, hydroxy-C₁₋₁₂ aliphatic-aminocarbonyl, C₁₋₁₂-alkoxy-C₁₋₁₂ aliphatic-aminocarbonyl, C₁₋₁₂ alkoxy-C₁₋₁₂ aliphatic-amino, di-
- 5 C₁₋₁₂ aliphatic amino, di-C₁₋₁₂ aliphatic aminocarbonyl, di-C₁₋₁₂ aliphatic aminosulfonyl, halogen, hydroxy, C₁₋₁₂ aliphatic-sulfonyl, aminosulfonyl, and C₁₋₁₂ aliphatic-aminosulfonyl, where R⁹, Aryl and Het are as defined below, with the proviso that where X is nitrogen, R² is not chloro or 3,6-dihydro-6-methyl-2-oxo-2H-1,3,4-thiadiazin-5-yl;
- 10 R¹ and R² are optionally joined to form a fused ring selected from the group as defined for Het below, and said fused ring is optionally substituted by a substituent selected from the group consisting of: C₁₋₁₂ aliphatic, halogen, nitro, cyano, C₁₋₁₂ alkoxy, amino, hydroxyl, (R¹⁰, R¹¹)-amino, and oxo; R³ is selected from the group consisting of: hydrogen, C₁₋₁₂ aliphatic,
- 15 hydroxy, hydroxy C₁₋₁₂ aliphatic, di-C₁₋₁₂ aliphatic amino, di-C₁₋₁₂ aliphatic aminocarbonyl, di-C₁₋₁₂ aliphatic aminosulfonyl, C₁₋₁₂ alkoxy, Aryl, Aryloxy, hydroxy-Aryl, Het, hydroxy-Het, Het-oxy, or halogen, where Aryl and Het are as defined below, with the proviso that where X is nitrogen R³ is not fluoro; R² and R³ are optionally joined to form a fused ring selected from the group
- 20 as defined for Het below, and said fused ring is optionally substituted by C₁₋₆ aliphatic or C₁₋₆ aliphatic-carbonyl; with the proviso that R¹, R², and R³ cannot simultaneously be hydrogen; R⁴, R⁵ and R⁶ may be the same or different and are independently selected from the group consisting of: hydrogen, C₁₋₁₂ aliphatic, thiol, C₁₋₆aliphatic-
- 25 thio, di(trifluoromethyl)hydroxymethyl, carboxamide, mono-C₁₋₁₂aliphatic aminocarbonyl, hydroxy, hydroxy-C₁₋₁₂ aliphatic, Aryl, Aryl-C₁₋₁₂ aliphatic, R⁹-Aryl-C₁₋₁₂ aliphatic, Cyc, Cyc-C₁₋₆ aliphatic, Het, Het-C₁₋₁₂ aliphatic, C₁₋₁₂ alkoxy, Aryloxy, Het-oxy, amino, (R¹⁰,R¹¹)-amino-C₁₋₁₂ aliphatic aminocarbonyl, (R¹⁰,R¹¹)-amino-C₁₋₁₂ aliphatic alkoxycarbonyl, (R¹⁰,R¹¹)-
- 30 amino-C₁₋₁₂ aliphatic aminocarbonylamino, (R¹⁰,R¹¹)-amino-C₁₋₆ aliphatic

- alkoxycarbonylamino, (R¹⁰,R¹¹)-amino-C₁₋₆ aliphaticsulfonyl, Het-C₁₋₆ aliphatic aminocarbonyl, Het-C₁₋₆ aliphatic aminocarbonylamino, Het-C₁₋₆ alkoxycarbonylamino, Het-C₁₋₆ aliphatic carbonyl, Het-C₁₋₆ alkoxycarbonyl, C₁₋₆ aliphaticsulfonyl-C₁₋₆ aliphatic aminoalkyl, C₁₋₆ aliphaticsulfonyl-C₁₋₆
- 5 aliphatic aminoalkyl-Het-, C₁₋₆ alkoxycarbonyl, C₁₋₆ aliphatic carbonylamino, (C₁₋₆ aliphatic carbonyl)(C₁₋₆ aliphatic)amino, (R¹⁰,R¹¹)-amino-C₁₋₆ aliphatic carbonylamino, [(R¹⁰,R¹¹)-amino-C₁₋₆ aliphatic carbonyl][C₁₋₆ aliphatic]amino, (R¹⁰,R¹¹)-amino-C₁₋₆ aliphatic sulfonylamino, [(R¹⁰,R¹¹)-amino-C₁₋₆ aliphaticsulfonyl][C₁₋₆ aliphatic]amino, halogen, cyano,
- 10 diethoxyphosphorylmethyl, nitro, trifluoromethyl, and trifluoromethoxy, where R⁹, R¹⁰, R¹¹, Aryl, Cyc and Het are as defined below, with the proviso that where X is nitrogen, R⁴, R⁵ and R⁶ is not nitro; R⁷ and R⁸ may be the same or different and are independently selected from the group consisting of: hydrogen, halogen, C₁₋₂ alkoxy, hydroxy, C₁₋₃-
- 15 aliphatic, and C₁₋₃ aliphatic; with the proviso that R⁴, R⁵, R⁶, R⁷, and R⁸ cannot simultaneously be hydrogen; wherein R⁷ may additionally be optionally fused to R⁵ so as to form a fused benzo ring from the R⁵ to the R⁷ positions;
- 20 R⁹ is selected from the group consisting of: C₁₋₁₂ aliphatic, hydroxy, C₁₋₁₂ alkoxy and halogen; R¹⁰ and R¹¹ may be the same or different and are independently selected from the group consisting of: hydrogen, C₁₋₆ aliphatic and Het; Aryl is selected from the group consisting of: phenyl, naphthyl, phenanthryl
- 25 and anthracenyl; Cyc is selected from the group consisting of: cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl, and optionally has one or more degrees of unsaturation; Het is a saturated or unsaturated heteroatom ring system selected from the
- 30 group consisting of: benzimidazole, dihydrothiophene, dioxin, dioxane,

dioxolane, dithiane, dithiazine, dithiazole, dithiolane, furan, imidazole, isoquinoline, morpholine, oxazole, oxadiazole, oxathiazole, oxathiazolidine, oxazine, oxadiazine, piperazine, piperidine, pyran, pyrazine, pyrazole, pyridine, pyrimidine, pyrrole, pyrrolidine, quinoline, tetrahydrofuran, tetrazine,
5 thiadiazine, thiadiazole, thiatriazole, thiazine, thiazole, thiomorpholine, thiophene, thiopyran, triazine and triazole, where any of said heterocyclic rings may be optionally substituted by a substituent selected from the group consisting of: C₁₋₁₂ aliphatic, hydroxy, C₁₋₁₂ alkoxy, (R¹⁰,R¹¹)-amino, (R¹⁰,R¹¹)-amino-C₁₋₁₂ aliphatic, (R¹⁰,R¹¹)-amino-C₁₋₁₂ aliphatic amino, oxo
10 and dioxo;
and the pharmaceutically acceptable salts, polymorphs, esters, amides, carbamates, solvates, hydrates, affinity reagents and prodrugs thereof in either crystalline or amorphous form.

15 2. The compound of claim 1 wherein Y is carbon.

3. The compound of claim 1 wherein Z is carbon.

4. The compound of claim 1 wherein A is carbon.

20

5. The compound of claim 1 wherein D is carbon.

6. The compound of claim 1 wherein X is carbon.

25 7. The compound of claim 1 wherein Y, Z, A, D, and X are each carbon.

8. The compound of claim 1 wherein X is nitrogen.

9. The compound of claim 1 wherein Y, Z, A, and D are each carbon, and
30 X is nitrogen.

10. The compound of claim 1 where Y is nitrogen.
11. The compound of claim 1 where Z is nitrogen.
- 5 12. The compound of claim 1 where A is nitrogen.
13. The compound of claim 1 where D is nitrogen.
- 10 14. The compound of claim 1 wherein Z and D are nitrogen.
15. The compound of claim 1 wherein R¹ is selected from the group consisting of: hydrogen, fluoro, bromo, iodo, lower alkyl, cyano and nitro.
- 15 16. The compound of claim 1 wherein R¹ is selected from the group consisting of: hydrogen and methyl.
17. The compound of claim 1 wherein R² is selected from the group consisting of: hydroxyl, hydroxy and lower alkyl.
- 20 18. The compound of claim 1 wherein R² is selected from the group consisting of: 2-furanyl, 3-thienyl, Br, CO₂Et, H, and Ph.
19. The compound of claim 1 wherein R² is selected from the group
- 25 consisting of: hydrogen, lower alkyl, lower alkoxy, hydroxy lower alkyl, C₁₋₁₂ alkoxy carbonyl, Aryl, Het, aminocarbonyl, lower alkyl aminocarbonyl, halogen and hydroxy.

20. The compound of claim 1 wherein R¹ is joined with R² to form a fused ring structure selected from the group consisting of: thiazole, imidazole, triazole and pyridine.
- 5 21. The compound of claim 20 wherein the ring is selected from the group consisting of: fused thiazole and fused pyridine.
22. The compound of claim 20 wherein the ring is substituted by one or more substituents selected from the group consisting of: halogen, amino,
10 lower alkyl substituted amino, lower alkyl and lower alkyl carbonyl.
23. The compound of claim 20 wherein the ring is pyridine substituted by halogen or methyl.
- 15 24. The compound of claim 1 wherein R¹ and R² are joined to form a fused ring structure and are together selected from the group consisting of: -NH-C=N-, -NH-N=N-, and -S-CH=N-, and -CH=CH-CH=N-.
25. The compound of claim 1 wherein R¹ and R² are independently selected
20 from the group consisting of: CH₃ and OH.
26. The compound of claim 1 wherein R³ is selected from the group consisting of: hydrogen, lower alkyl, lower alkenyl, halogen, phenyl, Het and alkoxy.
- 25
27. The compound of claim 1 wherein R³ is selected from the group consisting of: hydrogen, halogen, ethenyl and methyl.
28. The compound of claim 1 wherein R³ is selected from the group
30 consisting of: Cl, CH₃, 2-furanyl, 2-thienyl, Br, CH=CH₂, H, and Ph.

29. The compound of claim 1 wherein R^4 is selected from the group consisting of: hydrogen, lower alkyl, hydroxy, hydroxy-lower alkyl, carboxamide, mono- lower alkyl aminocarbonyl, substituted Aryl-lower alkyl, 5 Het, Het-lower alkyl, lower alkoxy, Aryloxy, Het-oxy, amino, mono- or di-lower alkyl-amino lower alkyl aminocarbonyl, mono- or di- lower alkyl-amino lower alkoxy carbonyl, mono- or di-lower alkyl-amino lower alkyl aminocarbonylamino, mono- or di- lower alkyl-amino lower alkoxy carbonylamino, lower alkyl carbonylamino, (lower alkyl carbonyl)(lower 10 alkyl)amino, mono- or di-lower alkyl-amino lower alkyl carbonylamino, [mono- or di-lower alkyl-amino lower alkyl carbonyl][lower alkyl]amino, mono- or di-lower alkyl-amino lower alkyl sulfonylamino, [mono- or di-lower alkyl-amino lower alkyl sulfonyl][lower alkyl]amino, mono- or di- lower alkyl-amino lower alkyl sulfonyl, Het lower alkyl aminocarbonyl, Het lower alkyl 15 aminocarbonylamino, Het lower alkoxy carbonylamino, Het lower alkyl carbonyl, Het lower alkoxy carbonyl, lower alkyl sulfonyl lower alkyl aminoalkyl, lower alkyl sulfonyl-lower alkyl-aminoalkyl-Het-, lower alkoxy carbonyl, halogen, cyano, diethoxyphosphorylmethyl, trifluoromethyl and trifluoromethoxy.

20

30. The compound of claim 1 wherein R^4 is selected from the group consisting of: lower alkyl, hydroxy, hydroxy-lower alkyl, carboxamide, mono-lower alkyl aminocarbonyl, substituted Aryl-lower alkyl, Het, Het-lower alkyl, Het-oxy, mono- or di- lower alkyl-amino lower alkyl aminocarbonyl, mono- or 25 di- lower alkyl-amino lower alkoxy carbonyl, mono- or di-lower alkyl-amino lower alkyl aminocarbonylamino, mono- or di- lower alkyl-amino lower alkoxy carbonylamino, lower alkyl carbonylamino, (lower alkyl carbonyl)(lower alkyl)amino, mono- or di-lower alkyl-amino lower alkyl carbonylamino, [mono- or di-lower alkyl-amino lower alkyl carbonyl][lower alkyl]amino, mono- 30 or di-lower alkyl-amino lower alkyl sulfonylamino, [mono- or di-lower alkyl-

amino lower alkyl sulfonyl][lower alkyl]amino, mono- or di- lower alkyl-amino lower alkyl sulfonyl, Het lower alkyl aminocarbonyl, Het lower alkyl carbonyl, lower alkyl sulfonyl lower alkyl aminoalkyl, lower alkyl sulfonyl-lower alkyl-aminoalkyl-Het-, halogen, cyano and trifluoromethyl.

5

31. The compound of claim 1 wherein R^4 is selected from the group consisting of: hydroxymethyl, hydroxyethyl, 4-pyridylmethyl, 4-morpholino, acetamido, N-methylacetamido, carboxamide, diethylaminoethylsulfonyl, 5-methyl-3-pyrazolon-1-yl and 3-ethyl-piperidine-2,6-dion-3-yl.

10

32. The compound of claim 1 wherein R^4 is selected from the group consisting of: 2-(2-pyridyl)ethenyl, 2-(4-hydroxyphenyl)ethenyl, 3-(methoxycarbonyl)phenoxy, 3-ethyl-piperidine-2,6-dion-3-yl, 4-(methoxycarbonyl)phenoxy, 4-morpholino, 4-pyridylmethyl, 5-methyl-3-
15 pyrazolon-1-yl, 5-oxazolyl, benzoyl, Br, $C(OH)(CF_3)_2$, CF_3 , CH_2CH_2OH , $-CH_2PO(OEt)_2$, CH_3 , CN, CO_2Me , $CONH_2$, F, H, I, NHAc, NMeAc, NO_2 , OCH_2Ph , OH, OMe, OMe, OPh, SMe, and $SO_2CH_2CH_2NEt_2$.

33. The compound of claim 1 wherein R^5 is selected from the group
20 consisting of: hydrogen, lower alkyl, hydroxy, hydroxy-lower alkyl, carboxamide, mono- lower alkyl aminocarbonyl, substituted Aryl-lower alkyl, Het, Het-lower alkyl, lower alkoxy, Aryloxy, Het-oxy, amino, mono- or di-lower alkyl-amino lower alkyl aminocarbonyl, mono- or di- lower alkyl-amino lower alkoxy carbonyl, mono- or di-lower alkyl-amino lower alkyl
25 aminocarbonylamino, mono- or di- lower alkyl-amino lower alkoxy carbonylamino, lower alkyl carbonylamino, (lower alkyl carbonyl)(lower alkyl)amino, mono- or di-lower alkyl-amino lower alkyl carbonylamino, [mono- or di-lower alkyl-amino lower alkyl carbonyl][lower alkyl]amino, mono- or di-lower alkyl-amino lower alkyl sulfonylamino, [mono- or di-lower alkyl-
30 amino lower alkyl sulfonyl][lower alkyl]amino, mono- or di- lower alkyl-amino

lower alkyl sulfonyl, Het lower alkyl aminocarbonyl, Het lower alkyl aminocarbonylamino, Het lower alkoxy carbonylamino, Het lower alkyl carbonyl, Het lower alkoxy carbonyl, lower alkyl sulfonyl lower alkyl aminoalkyl, lower alkyl sulfonyl-lower alkyl-aminoalkyl-Het-, lower
5 alkoxy carbonyl, halogen, cyano, diethoxyphosphorylmethyl, trifluoromethyl and trifluoromethoxy.

34. The compound of claim 1 wherein R^5 is selected from the group consisting of: lower alkyl, hydroxy, hydroxy-lower alkyl, carboxamide, mono-
10 lower alkyl aminocarbonyl, substituted Aryl-lower alkyl, Het, Het-lower alkyl, Het-oxy, mono- or di- lower alkyl-amino lower alkyl aminocarbonyl, mono- or di- lower alkyl-amino lower alkoxy carbonyl, mono- or di-lower alkyl-amino lower alkyl aminocarbonylamino, mono- or di- lower alkyl-amino lower alkoxy carbonylamino, lower alkyl carbonylamino, (lower alkyl carbonyl)(lower
15 alkyl) amino, mono- or di-lower alkyl-amino lower alkyl carbonylamino, [mono- or di-lower alkyl-amino lower alkyl carbonyl][lower alkyl]amino, mono- or di-lower alkyl-amino lower alkyl sulfonylamino, [mono- or di-lower alkyl-amino lower alkyl sulfonyl][lower alkyl]amino, mono- or di- lower alkyl-amino lower alkyl sulfonyl, Het lower alkyl aminocarbonyl, Het lower alkyl carbonyl,
20 lower alkyl sulfonyl lower alkyl aminoalkyl, lower alkyl sulfonyl-lower alkyl-aminoalkyl-Het-, halogen, cyano and trifluoromethyl.

35. The compound of claim 1 wherein R^5 is selected from the group consisting of: hydroxymethyl, hydroxyethyl, 4-pyridylmethyl, 4-morpholino,
25 acetamido, N-methylacetamido, carboxamide, diethylaminoethylsulfonyl, 5-methyl-3-pyrazolon-1-yl and 3-ethyl-piperidine-2,6-dion-3-yl.

36. The compound of claim 1 wherein R^5 is selected from the group consisting of: CF_3 , CH_2OH , CH_3 , Cl, CN, $CONH_2$, F, H, and OMe.

37. The compound of claim 1 wherein R^6 is selected from the group consisting of: CH_3 , H, and OMe.


38. The compound of claim 1 wherein R^6 is hydrogen.

5

39. The compound of claim 1 wherein R^7 and R^8 are independently selected from the group consisting of: hydrogen, halogen and methyl.

40. The compound of claim 1 wherein R^7 is selected from the group
10 consisting of: $-CH_2CH_2OH$, H, and OMe.

41. The compound of claim 1 wherein R^8 is H.



15 42. A compound of formula (I) as claimed in claim 1 wherein

Y, Z, A, and D are independently selected from the group consisting of: carbon and nitrogen, with the provisos that: (1) Z and D may be nitrogen, but otherwise no more than one of Y, Z, A, and D may be nitrogen, and (2)
20 when Y, Z, or A are nitrogen, substituent R^1 , R^2 , or R^3 designated for the respective nitrogen atom is non-existent;

X is selected from the group consisting of: N, CH, CCF_3 , and $C(C_{1-12} \text{ aliphatic})$;

R¹ is selected from the group consisting of: hydrogen, fluoro, bromo, iodo, lower alkyl, cyano and nitro;

R² is selected from the group consisting of: hydrogen, lower alkyl, lower alkoxy, hydroxy lower alkyl, C₁₋₁₂ alkoxy carbonyl, Aryl, Het, aminocarbonyl,

5 lower alkyl aminocarbonyl, halogen and hydroxy;

R³ is selected from the group consisting of: hydrogen, lower alkyl, lower alkenyl, halogen, phenyl, Het and alkoxy;

R⁴ is selected from the group consisting of: hydrogen, lower alkyl, hydroxy, hydroxy-lower alkyl, carboxamide, mono- lower alkyl aminocarbonyl, 10 substituted Aryl-lower alkyl, Het, Het-lower alkyl, lower alkoxy, Aryloxy, Het-oxy, amino, mono- or di- lower alkyl-amino lower alkyl aminocarbonyl, mono- or di- lower alkyl-amino lower alkoxy carbonyl, mono- or di-lower alkyl-amino lower alkyl aminocarbonylamino, mono- or di- lower alkyl-amino lower alkoxy carbonylamino, lower alkyl carbonylamino, (lower alkyl carbonyl)(lower 15 alkyl)amino, mono- or di-lower alkyl-amino lower alkyl carbonylamino, [mono- or di-lower alkyl-amino lower alkyl carbonyl][lower alkyl]amino, mono- or di-lower alkyl-amino lower alkyl sulfonylamino, [mono- or di-lower alkyl-amino lower alkyl sulfonyl][lower alkyl]amino, mono- or di- lower alkyl-amino lower alkyl sulfonyl, Het lower alkyl aminocarbonyl, Het lower alkyl 20 aminocarbonylamino, Het lower alkoxy carbonylamino, Het lower alkyl carbonyl, Het lower alkoxy carbonyl, lower alkyl sulfonyl lower alkyl aminoalkyl, lower alkyl sulfonyl-lower alkyl-aminoalkyl-Het-, lower alkoxy carbonyl, halogen, cyano, diethoxyphosphorylmethyl, trifluoromethyl and trifluoromethoxy;

25 R⁶ is hydrogen;

R⁷ and R⁸ are independently selected from the group consisting of: hydrogen, halogen and methyl;

Aryl is selected from the group consisting of: phenyl, naphthyl, phenanthryl and anthracenyl;

Cyc is selected from the group consisting of: cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl, and optionally has one or more degrees of unsaturation;

Het is a saturated or unsaturated heteroatom ring system selected from the
5 group consisting of: benzimidazole, dihydrothiophene, dioxin, dioxane, dioxolane, dithiane, dithiazine, dithiazole, dithiolane, furan, imidazole, isoquinoline, morpholine, oxazole, oxadiazole, oxathiazole, oxathiazolidine, oxazine, oxadiazine, piperazine, piperidine, pyran, pyrazine, pyrazole, pyridine, pyrimidine, pyrrole, pyrrolidine, quinoline, tetrahydrofuran, tetrazine,
10 thiadiazine, thiadiazole, thiatriazole, thiazine, thiazole, thiomorpholine, thiophene, thiopyran, triazine and triazole, where any of said heterocyclic rings may be optionally substituted by a substituent selected from the group consisting of: C₁₋₁₂ aliphatic, hydroxy, C₁₋₁₂ alkoxy, (R¹⁰,R¹¹)-amino, (R¹⁰,R¹¹)-amino-C₁₋₁₂ aliphatic, (R¹⁰,R¹¹)-amino-C₁₋₁₂ aliphatic amino, oxo
15 and dioxo;
and the pharmaceutically acceptable salts, polymorphs, esters, amides, carbamates, solvates, hydrates, affinity reagents and prodrugs thereof.

43. A compound of the formula (I) as claimed in claim 1 wherein
20

Y, Z, A, and D are independently selected from the group consisting of: carbon and nitrogen, with the provisos that: (1) Z and D may be nitrogen, but otherwise no more than one of Y, Z, A, and D may be nitrogen, and (2) when Y, Z, or A are nitrogen, substituent R¹, R², or R³ designated for the
25 respective nitrogen atom is non-existent;

X is selected from the group consisting of: N, CH, CCF₃, and C(C₁₋₁₂ aliphatic);

R¹ is joined with R² to form a fused ring structure selected from the group consisting of: thiazole, imidazole, triazole and pyridine;

R³ is selected from the group consisting of: hydrogen, halogen, ethenyl and methyl;

- R⁴ is selected from the group consisting of: hydrogen, lower alkyl, hydroxy, hydroxy-lower alkyl, carboxamide, mono- lower alkyl aminocarbonyl, substituted Aryl-lower alkyl, Het, Het-lower alkyl, lower alkoxy, Aryloxy, Het-oxy, amino, mono- or di- lower alkyl-amino lower alkyl aminocarbonyl, mono- or di- lower alkyl-amino lower alkoxycarbonyl, mono- or di-lower alkyl-amino lower alkyl aminocarbonylamino, mono- or di- lower alkyl-amino lower alkoxycarbonylamino, lower alkyl carbonylamino, (lower alkyl carbonyl)(lower alkyl)amino, mono- or di-lower alkyl-amino lower alkyl carbonylamino, [mono- or di-lower alkyl-amino lower alkyl carbonyl][lower alkyl]amino, mono- or di-lower alkyl-amino lower alkyl sulfonylamino, [mono- or di-lower alkyl-amino lower alkyl sulfonyl][lower alkyl]amino, mono- or di- lower alkyl-amino lower alkyl sulfonyl, Het lower alkyl aminocarbonyl, Het lower alkyl aminocarbonylamino, Het lower alkoxycarbonylamino, Het lower alkyl carbonyl, Het lower alkoxycarbonyl, lower alkyl sulfonyl lower alkyl aminoalkyl, lower alkyl sulfonyl-lower alkyl-aminoalkyl-Het-, lower alkoxycarbonyl, halogen, cyano, diethoxyphosphorylmethyl, trifluoromethyl and trifluoromethoxy;
- R⁵ is selected from the group consisting of: lower alkyl, hydroxy, hydroxy-lower alkyl, carboxamide, mono- lower alkyl aminocarbonyl, substituted Aryl-lower alkyl, Het, Het-lower alkyl, Het-oxy, mono- or di- lower alkyl-amino lower alkyl aminocarbonyl, mono- or di- lower alkyl-amino lower alkoxycarbonyl, mono- or di-lower alkyl-amino lower alkyl aminocarbonylamino, mono- or di- lower alkyl-amino lower alkoxycarbonylamino, lower alkyl carbonylamino, (lower alkyl carbonyl)(lower alkyl) amino, mono- or di-lower alkyl-amino lower alkyl carbonylamino, [mono- or di-lower alkyl-amino lower alkyl carbonyl][lower alkyl]amino, mono- or di-lower alkyl-amino lower alkyl sulfonylamino, [mono- or di-lower alkyl-amino lower alkyl sulfonyl][lower alkyl]amino, mono- or di- lower alkyl-amino

lower alkyl sulfonyl, Het lower alkyl aminocarbonyl, Het lower alkyl carbonyl, lower alkyl sulfonyl lower alkyl aminoalkyl, lower alkyl sulfonyl-lower alkyl-aminoalkyl-Het-, halogen, cyano and trifluoromethyl;

R⁶ is hydrogen;

- 5 R⁷ and R⁸ are independently selected from the group consisting of: hydrogen, halogen and methyl;

Aryl is selected from the group consisting of: phenyl, naphthyl, phenanthryl and anthracenyl;

- Cyc is selected from the group consisting of: cyclopropyl, cyclobutyl, 10 cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl, and optionally has one or more degrees of unsaturation;

Het is a saturated or unsaturated heteroatom ring system selected from the group consisting of: benzimidazole, dihydrothiophene, dioxin, dioxane, dioxolane, dithiane, dithiazine, dithiazole, dithiolane, furan, imidazole,

- 15 isoquinoline, morpholine, oxazole, oxadiazole, oxathiazole, oxathiazolidine, oxazine, oxadiazine, piperazine, piperidine, pyran, pyrazine, pyrazole, pyridine, pyrimidine, pyrrole, pyrrolidine, quinoline, tetrahydrofuran, tetrazine, thiadiazine, thiadiazole, thiatriazole, thiazine, thiazole, thiomorpholine, thiophene, thiopyran, triazine and triazole, where any of said heterocyclic 20 rings may be optionally substituted by a substituent selected from the group consisting of: C₁₋₁₂ aliphatic, hydroxy, C₁₋₁₂ alkoxy, (R¹⁰, R¹¹)-amino, (R¹⁰, R¹¹)-amino-C₁₋₁₂ aliphatic, (R¹⁰, R¹¹)-amino-C₁₋₁₂ aliphatic amino, oxo and dioxo;

- and the pharmaceutically acceptable salts, polymorphs, esters, amides, 25 carbamates, solvates, hydrates, affinity reagents and prodrugs thereof.

44. A compound of the formula (I) as claimed in claim 1 wherein

- Y, Z, A, and D are independently selected from the group consisting of: 30 carbon and nitrogen, with the provisos that: (1) Z and D may be nitrogen,

but otherwise no more than one of Y, Z, A, and D may be nitrogen, and (2) when Y, Z, or A are nitrogen, substituent R¹, R², or R³ designated for the respective nitrogen atom is non-existent

X is selected from the group consisting of: N, CH, CCF₃, and C(C₁₋₁₂

5 aliphatic)

R¹ is selected from the group consisting of: hydrogen and methyl;

R² is selected from the group consisting of: hydroxyl, hydroxy and lower alkyl;

R³ is selected from the group consisting of: hydrogen, halogen, ethenyl and

10 methyl;

R⁴ is selected from the group consisting of: hydrogen, lower alkyl, hydroxy, hydroxy-lower alkyl, carboxamide, mono- lower alkyl aminocarbonyl, substituted Aryl-lower alkyl, Het, Het-lower alkyl, lower alkoxy, Aryloxy, Het-oxy, amino, mono- or di- lower alkyl-amino lower alkyl aminocarbonyl, mono-

15 or di- lower alkyl-amino lower alkoxycarbonyl, mono- or di-lower alkyl-amino lower alkyl aminocarbonylamino, mono- or di- lower alkyl-amino lower alkoxycarbonylamino, lower alkyl carbonylamino, (lower alkyl carbonyl)(lower alkyl)amino, mono- or di-lower alkyl-amino lower alkyl carbonylamino, [mono- or di-lower alkyl-amino lower alkyl carbonyl][lower alkyl]amino, mono-

20 or di-lower alkyl-amino lower alkyl sulfonylamino, [mono- or di-lower alkyl-amino lower alkyl sulfonyl][lower alkyl]amino, mono- or di- lower alkyl-amino lower alkyl sulfonyl, Het lower alkyl aminocarbonyl, Het lower alkyl aminocarbonylamino, Het lower alkoxycarbonylamino, Het lower alkyl carbonyl, Het lower alkoxycarbonyl, lower alkyl sulfonyl lower alkyl

25 aminoalkyl, lower alkyl sulfonyl-lower alkyl-aminoalkyl-Het-, lower alkoxycarbonyl, halogen, cyano, diethoxyphosphorylmethyl, trifluoromethyl and trifluoromethoxy;

R⁵ is selected from the group consisting of: hydroxymethyl, hydroxyethyl, 4-pyridylmethyl, 4-morpholino, acetamido, N-methylacetamido, carboxamide,

diethylaminoethylsulfonyl, 5-methyl-3-pyrazolon-1-yl and 3-ethyl-piperidine-2,6-dion-3-yl;

R⁶ is hydrogen;

R⁷ and R⁸ are independently selected from the group consisting of: hydrogen,

5 halogen and methyl;

Aryl is selected from the group consisting of: phenyl, naphthyl, phenanthryl and anthracenyl;

Cyc is selected from the group consisting of: cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl, and optionally has one or

10 more degrees of unsaturation;


Het is a saturated or unsaturated heteroatom ring system selected from the group consisting of: benzimidazole, dihydrothiophene, dioxin, dioxane, dioxolane, dithiane, dithiazine, dithiazole, dithiolane, furan, imidazole, isoquinoline, morpholine, oxazole, oxadiazole, oxathiazole, oxathiazolidine,

15 oxazine, oxadiazine, piperazine, piperidine, pyran, pyrazine, pyrazole, pyridine, pyrimidine, pyrrole, pyrrolidine, quinoline, tetrahydrofuran, tetrazine, thiadiazine, thiadiazole, thiatriazole, thiazine, thiazole, thiomorpholine, thiophene, thiopyran, triazine and triazole, where any of said heterocyclic rings may be optionally substituted by a substituent selected from the group
20 consisting of: C₁₋₁₂ aliphatic, hydroxy, C₁₋₁₂ alkoxy, (R¹⁰, R¹¹)-amino, (R¹⁰, R¹¹)-amino-C₁₋₁₂ aliphatic, (R¹⁰, R¹¹)-amino-C₁₋₁₂ aliphatic amino, oxo and dioxo;

and the pharmaceutically acceptable salts, polymorphs, esters, amides, carbamates, solvates, hydrates, affinity reagents and prodrugs thereof.

25

45. A compound of the formula (II):



wherein:

R¹ is selected from the group consisting of: hydrogen, C₁₋₁₂ aliphatic, thiol, hydroxy, hydroxy-C₁₋₁₂ aliphatic, Aryl, Aryl-C₁₋₁₂ aliphatic, R⁹-Aryl-C₁₋₁₂ aliphatic, Cyc, Cyc-C₁₋₆ aliphatic, Het, Het-C₁₋₁₂ aliphatic, C₁₋₁₂ alkoxy, Aryloxy, amino, C₁₋₁₂ aliphatic amino, di-C₁₋₁₂ aliphatic amino, di-C₁₋₁₂ aliphatic aminocarbonyl, di-C₁₋₁₂ aliphatic aminosulfonyl, C₁₋₁₂ alkoxycarbonyl, fluoro, bromo, iodo, cyano, sulfonamide, or nitro, where R⁹, Aryl, Cyc and Het are as defined below;

10 R² is selected from the group consisting of: hydrogen, C₁₋₁₂ aliphatic, N-hydroxyimino-C₁₋₁₂ aliphatic, C₁₋₁₂ alkoxy, hydroxy-C₁₋₁₂ aliphatic, C₁₋₁₂ alkoxycarbonyl, carboxyl C₁₋₁₂ aliphatic, Aryl, R⁹-Aryl-oxycarbonyl, R⁹-oxycarbonyl-Aryl, Het, aminocarbonyl, C₁₋₁₂ aliphatic-aminocarbonyl, Aryl-C₁₋₁₂ aliphatic-aminocarbonyl, R⁹-Aryl-C₁₋₁₂ aliphatic-aminocarbonyl, Het-C₁₋₁₂ aliphatic-aminocarbonyl, hydroxy-C₁₋₁₂ aliphatic-aminocarbonyl, C₁₋₁₂-alkoxy-C₁₋₁₂ aliphatic-aminocarbonyl, C₁₋₁₂ alkoxy-C₁₋₁₂ aliphatic-amino, di-C₁₋₁₂ aliphatic amino, di-C₁₋₁₂ aliphatic aminocarbonyl, di-C₁₋₁₂ aliphatic aminosulfonyl, halogen, hydroxy, C₁₋₁₂ aliphatic-sulfonyl, aminosulfonyl, or one or more substituents selected from the group consisting of: C₁₋₁₂ aliphatic-aminosulfonyl, where R⁹, Aryl and Het are as defined below;

R¹ and R² are optionally joined to form a fused ring selected from the group as defined for Het below, and said fused ring is optionally substituted by C₁₋₁₂ aliphatic, halogen, nitro, cyano, C₁₋₁₂ alkoxy, amino, hydroxyl, (R¹⁰, R¹¹)-amino, or oxo;

25 R³ is selected from the group consisting of: hydrogen, C₁₋₁₂ aliphatic, hydroxy, hydroxy C₁₋₁₂ aliphatic, di-C₁₋₁₂ aliphatic amino, di-C₁₋₁₂ aliphatic aminocarbonyl, di-C₁₋₁₂ aliphatic aminosulfonyl, C₁₋₁₂ alkoxy, Aryl, Aryloxy, hydroxy-Aryl, Het, hydroxy-Het, Het-oxy, or halogen, where Aryl and Het are as defined below;

R² and R³ are optionally joined to form a fused ring selected from the group as defined for Het below, and said fused ring is optionally substituted by C₁₋₆ aliphatic or C₁₋₆ aliphatic-carbonyl;

with the proviso that R¹, R² and R³ cannot simultaneously be hydrogen;

- 5 R⁴, R⁵ and R⁶ may be the same or different and are independently selected from the group consisting of: hydrogen, C₁₋₁₂ aliphatic, thiol, C₁₋₆aliphatic-thio, di(trifluoromethyl)hydroxymethyl, carboxamide, mono-C₁₋₁₂aliphatic aminocarbonyl, hydroxy, hydroxy-C₁₋₁₂ aliphatic, Aryl, Aryl-C₁₋₁₂ aliphatic, R⁹-Aryl-C₁₋₁₂ aliphatic, Cyc, Cyc-C₁₋₆ aliphatic, Het, Het-C₁₋₁₂ aliphatic, C₁₋₁₂
- 10 alkoxy, Aryloxy, Het-oxy, amino, (R¹⁰,R¹¹)-amino-C₁₋₁₂ aliphatic aminocarbonyl, (R¹⁰,R¹¹)-amino-C₁₋₁₂ aliphatic alkoxycarbonyl, (R¹⁰,R¹¹)-amino-C₁₋₁₂ aliphatic aminocarbonylamino, (R¹⁰,R¹¹)-amino-C₁₋₆ aliphatic alkoxycarbonylamino, (R¹⁰,R¹¹)-amino-C₁₋₆ aliphatic sulfonyl, Het-C₁₋₆ aliphatic aminocarbonyl, Het-C₁₋₆ aliphatic aminocarbonylamino, Het-C₁₋₆
- 15 alkoxycarbonylamino, Het-C₁₋₆ aliphatic carbonyl, Het-C₁₋₆ alkoxycarbonyl, C₁₋₆ aliphatic sulfonyl-C₁₋₆ aliphatic aminoalkyl, C₁₋₆ aliphatic sulfonyl-C₁₋₆ aliphatic aminoalkyl-Het-, C₁₋₆ alkoxycarbonyl, C₁₋₆ aliphatic carbonylamino, (C₁₋₆ aliphatic carbonyl)(C₁₋₆ aliphatic)amino, (R¹⁰,R¹¹)-amino-C₁₋₆ aliphatic carbonylamino, [(R¹⁰,R¹¹)-amino-C₁₋₆ aliphatic carbonyl][C₁₋₆ aliphatic]amino,
- 20 (R¹⁰,R¹¹)-amino-C₁₋₆ aliphatic sulfonylamino, [(R¹⁰,R¹¹)-amino-C₁₋₆ aliphatic sulfonyl][C₁₋₆ aliphatic]amino, halogen, cyano, diethoxyphosphorylmethyl, nitro, trifluoromethyl, or trifluoromethoxy, where R⁹, R¹⁰, R¹¹, Aryl, Cyc and Het are as defined below;

R⁷ and R⁸ may be the same or different and are independently selected from

- 25 the group consisting of: hydrogen, halogen, C₁₋₂ alkoxy, hydroxy, C₁₋₃-aliphatic and C₁₋₃ aliphatic;

with the proviso that R⁴, R⁵, R⁶, R⁷ and R⁸ cannot simultaneously be hydrogen;

wherein R⁷ may additionally be optionally fused to R⁵ so as to form a fused

- 30 benzo ring from the R⁵ to the R⁷ positions;

R⁹ is selected from the group consisting of: C₁₋₁₂ aliphatic, hydroxy, C₁₋₁₂ alkoxy, or halogen;

R¹⁰ and R¹¹ may be the same or different and are independently selected from the group consisting of: hydrogen, C₁₋₆ aliphatic and Het;


5 Aryl is selected from the group consisting of: phenyl, naphthyl, phenanthryl or anthracenyl;

Cyc is selected from the group consisting of: cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl, and optionally has one or more degrees of unsaturation;

10 Het is a saturated or unsaturated heteroatom ring system selected from the group consisting of: benzimidazole, dihydrothiophene, dioxin, dioxane, dioxolane, dithiane, dithiazine, dithiazole, dithiolane, furan, imidazole, isoquinoline, morpholine, oxazole, oxadiazole, oxathiazole, oxathiazolidine, oxazine, oxadiazine, piperazine, piperidine, pyran, pyrazine, pyrazole,
15 pyridine, pyrimidine, pyrrole, pyrrolidine, quinoline, tetrahydrofuran, tetrazine, thiadiazine, thiadiazole, thiatriazole, thiazine, thiazole, thiomorpholine, thiophene, thiopyran, triazine and triazole, where any of said heterocyclic rings may be optionally substituted by a substituent selected from the group consisting of: C₁₋₁₂ aliphatic, hydroxy, C₁₋₁₂ alkoxy, (R¹⁰,R¹¹)-amino,
20 (R¹⁰,R¹¹)-amino-C₁₋₁₂ aliphatic, (R¹⁰,R¹¹)-amino-C₁₋₁₂ aliphatic amino, oxo or dioxo;

and the pharmaceutically acceptable salts, esters, amides, carbamates, solvates, hydrates, affinity reagents and prodrugs thereof.

25 46. A compound of the formula (III):



- R¹ is selected from the group consisting of: hydrogen, C₁₋₁₂ aliphatic, thiol, hydroxy, hydroxy-C₁₋₁₂ aliphatic, Aryl, Aryl-C₁₋₁₂ aliphatic, R⁹-Aryl-C₁₋₁₂ aliphatic, Cyc, Cyc-C₁₋₆ aliphatic, Het, Het-C₁₋₁₂ aliphatic, C₁₋₁₂ alkoxy, Aryloxy, amino, C₁₋₁₂ aliphatic amino, di-C₁₋₁₂ aliphatic amino, di-C₁₋₁₂ aliphatic aminocarbonyl, di-C₁₋₁₂ aliphatic aminosulfonyl, C₁₋₁₂ alkoxy-carbonyl, fluoro, bromo, iodo, cyano, sulfonamide, or nitro, where R⁹, Aryl, Cyc and Het are as defined below;
- 10 R² is selected from the group consisting of: hydrogen, C₁₋₁₂ aliphatic, N-hydroxyimino-C₁₋₁₂ aliphatic, C₁₋₁₂ alkoxy, hydroxy-C₁₋₁₂ aliphatic, C₁₋₁₂ alkoxy-carbonyl, carboxyl C₁₋₁₂ aliphatic, Aryl, R⁹-Aryl-oxycarbonyl, R⁹-oxycarbonyl-Aryl, Het, aminocarbonyl, C₁₋₁₂ aliphatic-aminocarbonyl, Aryl-C₁₋₁₂ aliphatic-aminocarbonyl, R⁹-Aryl-C₁₋₁₂ aliphatic-aminocarbonyl, Het-
- 15 C₁₋₁₂ aliphatic-aminocarbonyl, hydroxy-C₁₋₁₂ aliphatic-aminocarbonyl, C₁₋₁₂-alkoxy-C₁₋₁₂ aliphatic-aminocarbonyl, C₁₋₁₂ alkoxy-C₁₋₁₂ aliphatic-amino, di-C₁₋₁₂ aliphatic amino, di-C₁₋₁₂ aliphatic aminocarbonyl, di-C₁₋₁₂ aliphatic aminosulfonyl, halogen, hydroxy, C₁₋₁₂ aliphatic-sulfonyl, aminosulfonyl, or one or more substituents selected from the group consisting of: C₁₋₁₂
- 20 aliphatic-aminosulfonyl, where R⁹, Aryl and Het are as defined below, with the proviso that R² is not chloro or 3,6-dihydro-6-methyl-2-oxo-2H-1,3,4-thiadiazin-5-yl;

R¹ and R² are optionally joined to form a fused ring selected from the group as defined for Het below, and said fused ring is optionally substituted by C₁₋₁₂ aliphatic, halogen, nitro, cyano, C₁₋₁₂ alkoxy, amino, hydroxyl, (R¹⁰, R¹¹)-amino, or oxo;

- 5 R³ is selected from the group consisting of: hydrogen, C₁₋₁₂ aliphatic, hydroxy, hydroxy C₁₋₁₂ aliphatic, di-C₁₋₁₂ aliphatic amino, di-C₁₋₁₂ aliphatic aminocarbonyl, di-C₁₋₁₂ aliphatic aminosulfonyl, C₁₋₁₂ alkoxy, Aryl, Aryloxy, hydroxy-Aryl, Het, hydroxy-Het, Het-oxy, or halogen, where Aryl and Het are as defined below, with the proviso R³ is not fluoro;
- 10 R² and R³ are optionally joined to form a fused ring selected from the group as defined for Het below, and said fused ring is optionally substituted by C₁₋₆ aliphatic or C₁₋₆ aliphatic-carbonyl;
with the proviso that R¹, R² and R³ cannot simultaneously be hydrogen;
R⁴, R⁵ and R⁶ may be the same or different and are independently selected
- 15 from the group consisting of: hydrogen, C₁₋₁₂ aliphatic, thiol, C₁₋₆aliphatic-thio, di(trifluoromethyl)hydroxymethyl, carboxamide, mono-C₁₋₁₂aliphatic aminocarbonyl, hydroxy, hydroxy-C₁₋₁₂ aliphatic, Aryl, Aryl-C₁₋₁₂ aliphatic, R⁹-Aryl-C₁₋₁₂ aliphatic, Cyc, Cyc-C₁₋₆ aliphatic, Het, Het-C₁₋₁₂ aliphatic, C₁₋₁₂ alkoxy, Aryloxy, Het-oxy, amino, (R¹⁰,R¹¹)-amino-C₁₋₁₂ aliphatic
- 20 aminocarbonyl, (R¹⁰,R¹¹)-amino-C₁₋₁₂ aliphatic alkoxycarbonyl, (R¹⁰,R¹¹)-amino-C₁₋₁₂ aliphatic aminocarbonylamino, (R¹⁰,R¹¹)-amino-C₁₋₆ aliphatic alkoxycarbonylamino, (R¹⁰,R¹¹)-amino-C₁₋₆ aliphatic sulfonyl, Het-C₁₋₆ aliphatic aminocarbonyl, Het-C₁₋₆ aliphatic aminocarbonylamino, Het-C₁₋₆ alkoxycarbonylamino, Het-C₁₋₆ aliphatic carbonyl, Het-C₁₋₆ alkoxycarbonyl,
- 25 C₁₋₆ aliphatic sulfonyl-C₁₋₆ aliphatic aminoalkyl, C₁₋₆ aliphatic sulfonyl-C₁₋₆ aliphatic aminoalkyl-Het-, C₁₋₆ alkoxycarbonyl, C₁₋₆ aliphatic carbonylamino, (C₁₋₆ aliphatic carbonyl)(C₁₋₆ aliphatic)amino, (R¹⁰,R¹¹)-amino-C₁₋₆ aliphatic carbonylamino, [(R¹⁰,R¹¹)-amino-C₁₋₆ aliphatic carbonyl][C₁₋₆ aliphatic]amino, (R¹⁰,R¹¹)-amino-C₁₋₆ aliphatic sulfonylamino, [(R¹⁰,R¹¹)-amino-C₁₋₆
- 30 aliphatic sulfonyl][C₁₋₆ aliphatic]amino, halogen, cyano,

diethoxyphosphorylmethyl, nitro, trifluoromethyl, or trifluoromethoxy, where R^9 , R^{10} , R^{11} , Aryl, Cyc and Het are as defined below, with the proviso that R^4 , R^5 and R^6 is not nitro;

R^7 and R^8 may be the same or different and are independently selected from
5 the group consisting of: hydrogen, halogen, C_{1-2} alkoxy, hydroxy, C_{1-3} -aliphatic and C_{1-3} aliphatic;
with the proviso that R^4 , R^5 , R^6 , R^7 and R^8 cannot simultaneously be hydrogen;

R^9 is selected from the group consisting of: C_{1-12} aliphatic, hydroxy, C_{1-12}
10 alkoxy, or halogen;

R^{10} and R^{11} may be the same or different and are independently selected from the group consisting of: hydrogen, C_{1-6} aliphatic and Het;

Aryl is selected from the group consisting of: phenyl, naphthyl, phenanthryl or anthracenyl;


15 Cyc is selected from the group consisting of: cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or cyclooctyl, any one of which may have one or more degrees of unsaturation;

Het is a saturated or unsaturated heteroatom ring system selected from the group consisting of: benzimidazole, dihydrothiophene, dioxin, dioxane,
20 dioxolane, dithiane, dithiazine, dithiazole, dithiolane, furan, imidazole, isoquinoline, morpholine, oxazole, oxadiazole, oxathiazole, oxathiazolidine, oxazine, oxadiazine, piperazine, piperidine, pyran, pyrazine, pyrazole, pyridine, pyrimidine, pyrrole, pyrrolidine, quinoline, tetrahydrofuran, tetrazine, thiadiazine, thiadiazole, thiatriazole, thiazine, thiazole, thiomorpholine,
25 thiophene, thiopyran, triazine and triazole, where any of said heterocyclic rings may be optionally substituted by a substituent selected from the group consisting of: C_{1-12} aliphatic, hydroxy, C_{1-12} alkoxy, (R^{10}, R^{11}) -amino, (R^{10}, R^{11}) -amino- C_{1-12} aliphatic, (R^{10}, R^{11}) -amino- C_{1-12} aliphatic amino, oxo or dioxo;

and the pharmaceutically acceptable salts, esters, amides, carbamates, solvates, hydrates, affinity reagents and prodrugs thereof.

47. A compound of the formula (IV):

5



Y, Z, A, and D are independently selected from the group consisting of: carbon and nitrogen, with the provisos that: (1) Z and D may be nitrogen, but otherwise no more than one of Y, Z, A, and D may be nitrogen, and (2)
 10 when Y, Z, or A are nitrogen, substituent R¹, R², or R³ designated for the respective nitrogen atom is non-existent;

R¹ is selected from the group consisting of: hydrogen, C₁₋₁₂ aliphatic, thiol, hydroxy, hydroxy-C₁₋₁₂ aliphatic, Aryl, Aryl-C₁₋₁₂ aliphatic, R⁹-Aryl-C₁₋₁₂ aliphatic, Cyc, Cyc-C₁₋₆ aliphatic, Het, Het-C₁₋₁₂ aliphatic, C₁₋₁₂ alkoxy,
 15 Aryloxy, amino, C₁₋₁₂ aliphatic amino, di-C₁₋₁₂ aliphatic amino, di-C₁₋₁₂ aliphatic aminocarbonyl, di-C₁₋₁₂ aliphatic aminosulfonyl, C₁₋₁₂ alkoxy carbonyl, fluoro, bromo, iodo, cyano, sulfonamide, or nitro, where R⁹, Aryl, Cyc and Het are as defined below;

R² is selected from the group consisting of: hydrogen, C₁₋₁₂ aliphatic, N-
 20 hydroxyimino-C₁₋₁₂ aliphatic, C₁₋₁₂ alkoxy, hydroxy-C₁₋₁₂ aliphatic, C₁₋₁₂ alkoxy carbonyl, carboxyl C₁₋₁₂ aliphatic, Aryl, R⁹-Aryl-oxycarbonyl, R⁹-oxycarbonyl-Aryl, Het, aminocarbonyl, C₁₋₁₂ aliphatic-aminocarbonyl, Aryl-C₁₋₁₂ aliphatic-aminocarbonyl, R⁹-Aryl-C₁₋₁₂ aliphatic-aminocarbonyl, Het-

C₁₋₁₂ aliphatic-aminocarbonyl, hydroxy-C₁₋₁₂ aliphatic-aminocarbonyl, C₁₋₁₂-alkoxy-C₁₋₁₂ aliphatic-aminocarbonyl, C₁₋₁₂ alkoxy-C₁₋₁₂ aliphatic-amino, di-C₁₋₁₂ aliphatic amino, di-C₁₋₁₂ aliphatic aminocarbonyl, di-C₁₋₁₂ aliphatic aminosulfonyl, halogen, hydroxy, C₁₋₁₂ aliphatic-sulfonyl, aminosulfonyl, or
5 one or more substituents selected from the group consisting of: C₁₋₁₂ aliphatic-aminosulfonyl, where R⁹, Aryl and Het are as defined below;

R¹ and R² are optionally joined to form a fused ring selected from the group as defined for Het below, and said fused ring is optionally substituted by C₁₋₁₂ aliphatic, halogen, nitro, cyano, C₁₋₁₂ alkoxy, amino, hydroxyl, (R¹⁰,
10 R¹¹)-amino, or oxo;

R³ is selected from the group consisting of: hydrogen, C₁₋₁₂ aliphatic, hydroxy, hydroxy C₁₋₁₂ aliphatic, di-C₁₋₁₂ aliphatic amino, di-C₁₋₁₂ aliphatic aminocarbonyl, di-C₁₋₁₂ aliphatic aminosulfonyl, C₁₋₁₂ alkoxy, Aryl, Aryloxy, hydroxy-Aryl, Het, hydroxy-Het, Het-oxy, or halogen, where Aryl and Het are
15 as defined below;

R² and R³ are optionally joined to form a fused ring selected from the group as defined for Het below, and said fused ring is optionally substituted by C₁₋₆ aliphatic or C₁₋₆ aliphatic-carbonyl;

with the proviso that R¹, R² and R³ cannot simultaneously be hydrogen;

20 R⁴, R⁵ and R⁶ may be the same or different and are independently selected from the group consisting of: hydrogen, C₁₋₁₂ aliphatic, thiol, C₁₋₆aliphatic-thio, di(trifluoromethyl)hydroxymethyl, carboxamide, mono-C₁₋₁₂aliphatic aminocarbonyl, hydroxy, hydroxy-C₁₋₁₂ aliphatic, Aryl, Aryl-C₁₋₁₂ aliphatic, R⁹-Aryl-C₁₋₁₂ aliphatic, Cyc, Cyc-C₁₋₆ aliphatic, Het, Het-C₁₋₁₂ aliphatic, C₁₋₁₂
25 alkoxy, Aryloxy, Het-oxy, amino, (R¹⁰,R¹¹)-amino-C₁₋₁₂ aliphatic aminocarbonyl, (R¹⁰,R¹¹)-amino-C₁₋₁₂ aliphatic alkoxycarbonyl, (R¹⁰,R¹¹)-amino-C₁₋₁₂ aliphatic aminocarbonylamino, (R¹⁰,R¹¹)-amino-C₁₋₆ aliphatic alkoxycarbonylamino, (R¹⁰,R¹¹)-amino-C₁₋₆ aliphatic sulfonyl, Het-C₁₋₆
30 aliphatic aminocarbonyl, Het-C₁₋₆ aliphatic aminocarbonylamino, Het-C₁₋₆ alkoxycarbonylamino, Het-C₁₋₆ aliphatic carbonyl, Het-C₁₋₆ alkoxycarbonyl,

- C₁₋₆ aliphaticsulfonyl-C₁₋₆ aliphatic aminoalkyl, C₁₋₆ aliphaticsulfonyl-C₁₋₆ aliphatic aminoalkyl-Het-, C₁₋₆ alkoxycarbonyl, C₁₋₆ aliphatic carbonylamino, (C₁₋₆ aliphatic carbonyl)(C₁₋₆ aliphatic)amino, (R¹⁰, R¹¹)-amino-C₁₋₆ aliphatic carbonylamino, [(R¹⁰, R¹¹)-amino-C₁₋₆ aliphatic carbonyl][C₁₋₆ aliphatic]amino,
- 5 (R¹⁰, R¹¹)-amino-C₁₋₆ aliphatic sulfonylamino, [(R¹⁰, R¹¹)-amino-C₁₋₆ aliphaticsulfonyl][C₁₋₆ aliphatic]amino, halogen, cyano, diethoxyphosphorylmethyl, nitro, trifluoromethyl, or trifluoromethoxy, where R⁹, R¹⁰, R¹¹, Aryl, Cyc and Het are as defined below;
- R⁷ and R⁸ may be the same or different and are independently selected from
- 10 the group consisting of: hydrogen, halogen, C₁₋₂ alkoxy, hydroxy, C₁₋₃-aliphatic and C₁₋₃ aliphatic;
- with the proviso that R⁴, R⁵, R⁶, R⁷ and R⁸ cannot simultaneously be hydrogen;
- wherein R⁷ may additionally be optionally fused to R⁵ so as to form a fused
- 15 benzo ring from the R⁵ to the R⁷ positions;
- R⁹ is selected from the group consisting of: C₁₋₁₂ aliphatic, hydroxy, C₁₋₁₂ alkoxy, or halogen;
- R¹⁰ and R¹¹ may be the same or different and are independently selected from the group consisting of: hydrogen, C₁₋₆ aliphatic and Het;
- 20 Aryl is selected from the group consisting of: phenyl, naphthyl, phenanthryl or anthracenyl;
- Cyc is selected from the group consisting of: cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or cyclooctyl, and optionally has one or more degrees of unsaturation;
- 25 Het is a saturated or unsaturated heteroatom ring system selected from the group consisting of: benzimidazole, dihydrothiophene, dioxin, dioxane, dioxolane, dithiane, dithiazine, dithiazole, dithiolane, furan, imidazole, isoquinoline, morpholine, oxazole, oxadiazole, oxathiazole, oxathiazolidine, oxazine, oxadiazine, piperazine, piperidine, pyran, pyrazine, pyrazole,
- 30 pyridine, pyrimidine, pyrrole, pyrrolidine, quinoline, tetrahydrofuran, tetrazine,

thiadiazine, thiadiazole, thiatriazole, thiazine, thiazole, thiomorpholine, thiophene, thiopyran, triazine and triazole, where any of said heterocyclic rings may be optionally substituted by a substituent selected from the group consisting of: C₁₋₁₂ aliphatic, hydroxy, C₁₋₁₂ alkoxy, (R¹⁰,R¹¹)-amino, 5 (R¹⁰,R¹¹)-amino-C₁₋₁₂ aliphatic, (R¹⁰,R¹¹)-amino-C₁₋₁₂ aliphatic amino, oxo or dioxo;

and the pharmaceutically acceptable salts, esters, amides, carbamates, solvates, hydrates, affinity reagents and prodrugs thereof.


10 48. A compound as claimed in any one of claims 1 to 47 in the form of a substantially pure E geometric isomer.


49. A compound as claimed in any one of claims 1 to 47 in the form of a substantially pure Z geometric isomer.

15

50. A compound as claimed in any one of claims 1 to 47 in the form of a mixture of E geometric isomer and Z geometric isomer in any proportions of said geometric isomers.


51. A compound as claimed in claim 1 selected from the group consisting of:









; and the pharmaceutically acceptable salts, polymorphs, esters, amides, carbamates, solvates, hydrates, affinity reagents and prodrugs thereof in 5 either crystalline or amorphous form.

52. A compound selected from the group consisting of:





; and the pharmaceutically acceptable salts, polymorphs, esters, amides, carbamates, solvates, hydrates, affinity reagents and prodrugs thereof in
5 either crystalline or amorphous form.

53. A compound as claimed in claim 1, in Z, E or Z and E form, selected from the group consisting of:

6-Bromo-3-[[4-(4-morpholinyl)anilino]methylidene]-1,3-dihydro-2H-indol-2-one;

6-Bromo-3-[[4-(4-pyridinylmethyl)anilino]methylidene]-1,3-dihydro-2H-indol-2-one;

8-[4-Toluidinomethylidene]-6,8-dihydroimidazo[4,5-e]indol-7(3H)-one;

8-[(3-Methyl-4-nitroanilino)methylidene]-3,6-dihydro[1,2,3]triazolo[4,5-e]indol-7-one;

8-[[4-Nitro-3-(trifluoromethyl)anilino]methylidene]-3,6-dihydro[1,2,3]triazolo[4,5-e]indol-7-one;

8-[(3-Chloro-4-nitroanilino)methylidene]-1,6-dihydro[1,2,3]triazolo[4,5-e]indol-7-one;

8-[(3,5-Dimethyl-4-nitroanilino)methylidene]-1,6-dihydro[1,2,3]triazolo[4,5-e]indol-7-one;

8-[(4-[2,2,2-Trifluoro-1-hydroxy-1-(trifluoromethyl)ethyl]anilino)methylidene]-1,6-dihydro[1,2,3]triazolo[4,5-e]indol-7-one;

8-[(4-(2-Hydroxyethyl)anilino)methylidene]-1,6-dihydro[1,2,3]triazolo[4,5-e]indol-7-one;

8-[(4-(Methylsulfanyl)anilino)methylidene]-6H-[1,3]thiazolo[5,4-e]indol-7-one;

8-[(3,5-Dimethoxyanilino)methylidene]-6H-[1,3]thiazolo[5,4-e]indol-7-one;

8-[(4-Hydroxyanilino)methylidene]-6H-[1,3]thiazolo[5,4-e]indol-7-one;

1-[(3-Methoxyanilino)methylidene]-1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one;

3-[(2-Oxo-2,3-dihydro-1H-pyrrolo[3,2-f]quinolin-1-ylidene)methyl]amino)benzonitrile;

1-[4-Toluidinomethylidene]-1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one;

1-[(4-Methoxyanilino)methylidene]-1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one;

3-[(7-Oxo-6,7-dihydro[1,2,3]triazolo[4,5-e]indol-8(1H)-ylidene)methyl]amino)benzonitrile;

8-[4-Toluidinomethylidene]-1,6-dihydro[1,2,3]triazolo[4,5-e]indol-7-one;

8-[(4-Methoxyanilino)methylidene]-1,6-dihydro[1,2,3]triazolo[4,5-e]indol-7-one;

8-[(4-(4-Morpholinyl)anilino)methylidene]-6H-[1,3]thiazolo[5,4-e]indol-7-one;

N-(4-[(7-Oxo-6,7-dihydro-8H-[1,3]thiazolo[5,4-e]indol-8-ylidene)methyl]amino)phenyl)acetamide;

8-[(4-(2-Hydroxyethyl)anilino)methylidene]-6H-[1,3]thiazolo[5,4-e]indol-7-one;

8-[(4-(4-Pyridinylmethyl)anilino)methylidene]-6H-[1,3]thiazolo[5,4-e]indol-7-one;

4-[(7-Oxo-6,7-dihydro-8H-[1,3]thiazolo[5,4-e]indol-8-ylidene)methyl]amino)benzamide;

6-Bromo-3-[[4-(4-morpholinyl)anilino]methylidene]-1,3-dihydro-2H-indol-2-one;
4-[[[(6-Bromo-2-oxo-1,2-dihydro-3H-indol-3-ylidene)methyl]amino]benzamide;
N-(4-[[[(6-Bromo-2-oxo-1,2-dihydro-3H-indol-3-ylidene)methyl]amino]phenyl]acetamide;
6-Bromo-3-[[4-(2-hydroxyethyl)anilino]methylidene]-1,3-dihydro-2H-indol-2-one;
1-[[4-(4-Morpholinyl)anilino]methylidene]-1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one;
1-[[4-(4-Pyridinylmethyl)anilino]methylidene]-1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one;
N-(4-[[[(2-Oxo-2,3-dihydro-1H-pyrrolo[3,2-f]quinolin-1-ylidene)methyl]amino]phenyl]acetamide;
1-[[4-(2-Hydroxyethyl)anilino]methylidene]-1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one;
4-[[[(2-Oxo-2,3-dihydro-1H-pyrrolo[3,2-f]quinolin-1-ylidene)methyl]amino]benzamide;
1-[[4-(4-Hydroxyanilino)methylidene]-1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one;
6-(2-Furyl)-3-[[4-(4-morpholinyl)anilino]methylidene]-1,3-dihydro-2H-indol-2-one;
N-[4-[[[(6-(2-Furyl)-2-oxo-1,2-dihydro-3H-indol-3-ylidene)methyl]amino]phenyl]acetamide;
6-(2-Furyl)-3-[[4-(2-hydroxyethyl)anilino]methylidene]-1,3-dihydro-2H-indol-2-one;
3-[[4-(4-Morpholinyl)anilino]methylidene]-6-vinyl-1,3-dihydro-2H-indol-2-one;
3-[[4-(4-Pyridinylmethyl)anilino]methylidene]-6-vinyl-1,3-dihydro-2H-indol-2-one;
N-(4-[[[(2-Oxo-6-vinyl-1,2-dihydro-3H-indol-3-ylidene)methyl]amino]phenyl]acetamide;
3-[[4-(2-Hydroxyethyl)anilino]methylidene]-6-vinyl-1,3-dihydro-2H-indol-2-one;
6-(2-Furyl)-3-[[4-(4-pyridinylmethyl)anilino]methylidene]-1,3-dihydro-2H-indol-2-one;
6-(2-Furyl)-3-[[4-(4-hydroxyanilino)methylidene]-1,3-dihydro-2H-indol-2-one;

3-[[4-(4-Morpholinyl)anilino]methylidene]-6-(2-thienyl)-1,3-dihydro-2H-indol-2-one;

N-[4-[[[2-Oxo-6-(2-thienyl)-1,2-dihydro-3H-indol-3-ylidene]methyl]amino]phenyl]acetamide;

6-Bromo-3-[[3-(hydroxymethyl)anilino]methylidene]-1,3-dihydro-2H-indol-2-one;

6-Bromo-3-[[4-(5-methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene]-1,3-dihydro-2H-indol-2-one;

3-Ethyl-3-(4-[[[7-oxo-6,7-dihydro-8H-[1,3]thiazolo[5,4-e]indol-8-ylidene)methyl]amino]phenyl)-2,6-piperidinedione;

8-[[4-Phenoxyanilino]methylidene]-6H-[1,3]thiazolo[5,4-e]indol-7-one;

8-[[4-(Benzyloxy)anilino]methylidene]-6H-[1,3]thiazolo[5,4-e]indol-7-one;

Methyl 4-(4-[[[7-oxo-6,7-dihydro-8H-[1,3]thiazolo[5,4-e]indol-8-ylidene)methyl]amino]phenoxy)benzoate;

Methyl 3-(4-[[[7-oxo-6,7-dihydro-8H-[1,3]thiazolo[5,4-e]indol-8-ylidene)methyl]amino]phenoxy)benzoate;

8-[[3-(Hydroxymethyl)anilino]methylidene]-6H-[1,3]thiazolo[5,4-e]indol-7-one;

3-[[[7-Oxo-6,7-dihydro-8H-[1,3]thiazolo[5,4-e]indol-8-ylidene)methyl]amino]benzamide;

8-[[4-(5-Methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene]-6H-[1,3]thiazolo[5,4-e]indol-7-one;

Methyl 4-[[[7-Oxo-6,7-dihydro-8H-[1,3]thiazolo[5,4-e]indol-8-ylidene)methyl]amino]benzoate;

4-[[[7-Oxo-6,7-dihydro-8H-[1,3]thiazolo[5,4-e]indol-8-ylidene)methyl]amino]benzonitrile;

N-Methyl-N-(4-[[[7-oxo-6,7-dihydro-8H-[1,3]thiazolo[5,4-e]indol-8-ylidene)methyl]amino]phenyl)acetamide;

1-[[4-Phenoxyanilino]methylidene]-1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one;

1-[[4-(Benzyloxy)anilino]methylidene]-1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one;

Methyl 4-(4-[[[2-Oxo-2,3-dihydro-1H-pyrrolo[3,2-f]quinolin-1-ylidene)methyl]amino]phenoxy)benzoate;

Methyl 3-(4-{{(2-Oxo-2,3-dihydro-1H-pyrrolo[3,2-f]quinolin-1-ylidene)methyl}amino}phenoxy)benzoate;

3-Ethyl-3-(4-{{(2-oxo-2,3-dihydro-1H-pyrrolo[3,2-f]quinolin-1-ylidene)methyl}amino}phenyl)-2,6-piperidinedione;

1-[(4-Benzoylanilino)methylidene]-1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one;

1-[[3-(Hydroxymethyl)anilino]methylidene]-1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one;

1-[[4-(5-Methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene]-1H-pyrrolo[3,2-f]quinolin-2(3H)-one

1-((E)-{4-[(E)-2-(4-Hydroxyphenyl)ethenyl]anilino}methylidene)-1,3-dihydro-2H-pyrrolo[3,2-f]quinolin-2-one;

3-[[{(2-Oxo-2,3-dihydro-1H-pyrrolo[3,2-f]quinolin-1-ylidene)methyl}amino]benzamide;

4-[[{(2-Oxo-2,3-dihydro-1H-pyrrolo[3,2-f]quinolin-1-ylidene)methyl}amino]benzonitrile;

Methyl 4-[[{(2-oxo-2,3-dihydro-1H-pyrrolo[3,2-f]quinolin-1-ylidene)methyl}amino]benzoate;

8-[(4-{{2-(Diethylamino)ethyl}sulfonyl}anilino)methylidene]-6H-[1,3]thiazolo[5,4-e]indol-7-one;

1-[(4-{{2-(Diethylamino)ethyl}sulfonyl}anilino)methylidene]-1H-pyrrolo[3,2-f]quinolin-2(3H)-one;

3-[[{(2-Oxo-6-phenyl-1,2-dihydro-3H-indol-3-ylidene)methyl}amino]benzamide;

Diethyl 4-[[{(Z and E)-(2-oxo-6-phenyl-1,2-dihydro-3H-indol-3-ylidene)methyl}amino]benzylphosphonate;

3-[[{(2-Oxo-6-phenyl-1,2-dihydro-3H-indol-3-ylidene)methyl}amino]benzonitrile;

3-[[2-(2-Hydroxyethyl)anilino]methylidene]-6-phenyl-1,3-dihydro-2H-indol-2-one;

3-[[3-(Hydroxymethyl)anilino]methylidene]-6-phenyl-1,3-dihydro-2H-indol-2-one;

3-[(2-Methoxyanilino)methylidene]-6-phenyl-1,3-dihydro-2H-indol-2-one;

3-[[4-(5-Methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene]-6-phenyl-1H-indol-2-one;

3-[(4-Iodoanilino)methylidene]-6-phenyl-1H-indol-2-one;

3-[[[6-(2-Furyl)-2-oxo-1,2-dihydro-3H-indol-3-ylidene]methyl]amino]benzamide;

Diethyl 4-[[[6-(2-furyl)-2-oxo-1,2-dihydro-3H-indol-3-ylidene]methyl]amino]benzylphosphonate;

3-[[[6-(2-Furyl)-2-oxo-1,2-dihydro-3H-indol-3-ylidene]methyl]amino]benzonitrile;

6-(2-Furyl)-3-[[2-(2-hydroxyethyl)anilino]methylidene]-1,3-dihydro-2H-indol-2-one;

6-(2-Furyl)-3-[[3-(hydroxymethyl)anilino]methylidene]-1,3-dihydro-2H-indol-2-one;

6-(2-Furyl)-3-[(2-methoxyanilino)methylidene]-1,3-dihydro-2H-indol-2-one;

6-(2-Furyl)-3-[[4-(5-methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene]-1H-indol-2-one;

6-(2-Furyl)-3-[(4-iodoanilino)methylidene]-1H-indol-2-one;

3,6-Dihydro[1,2,3]triazolo[4,5-e]indole-7,8-dione-8-[N-(4-methoxyphenyl)hydrazone];

3,6-Dihydro[1,2,3]triazolo[4,5-e]indole-7,8-dione-8-[N-[4-(1,3-oxazol-5-yl)phenyl]hydrazone];

3,6-Dihydro[1,2,3]triazolo[4,5-e]indole-7,8-dione-8-[N-(4-methylphenyl)hydrazone];

3,6-Dihydro[1,2,3]triazolo[4,5-e]indole-7,8-dione-8-(N-{4-[(E)-2-(2-pyridinyl)ethenyl]phenyl}hydrazone);

6H-[1,3]Thiazolo[5,4-e]indole-7,8-dione-8-[N-(3-methoxyphenyl)hydrazone];

5-Hydroxy-4,6-dimethyl-1H-indole-2,3-dione-3-[N-(4-methylphenyl)hydrazone];

3,6-Dihydro[1,2,3]triazolo[4,5-e]indole-7,8-dione-8-[N-[4-(trifluoromethyl)phenyl]hydrazone];

6H-[1,3]Thiazolo[5,4-e]indole-7,8-dione-8-[N-(3-fluorophenyl)hydrazone];

6H-[1,3]Thiazolo[5,4-e]indole-7,8-dione-8-[N-(4-fluorophenyl)hydrazone];

6H-[1,3]Thiazolo[5,4-e]indole-7,8-dione-8-[N-(4-bromophenyl)hydrazone];

3-[[2-Oxo-5-phenyl-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene)methyl]amino}benzamide;

Diethyl 4-[[2-oxo-5-phenyl-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene)methyl]amino}benzylphosphonate;

3-[[2-Oxo-5-phenyl-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene)methyl]amino}benzonitrile;

3-[[2-(2-Hydroxyethyl)anilino]methylidene]-5-phenyl-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

3-[[3-(Hydroxymethyl)anilino]methylidene]-5-phenyl-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

3-[[2-Methoxyanilino]methylidene]-5-phenyl-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

3-[[4-(5-Methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene]-5-phenyl-1H-pyrrolo[2,3-b]pyridin-2-one;

3-[[4-Iodoanilino]methylidene]-5-phenyl-1H-pyrrolo[2,3-b]pyridin-2-one;

3-[[5-(2-Furyl)-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene)methyl]amino}benzamide;

Diethyl 4-[[5-(2-furyl)-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene)methyl]amino}benzylphosphonate;

3-[[5-(2-Furyl)-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene)methyl]amino}benzonitrile;

5-(2-Furyl)-3-[[2-(2-hydroxyethyl)anilino]methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

5-(2-Furyl)-3-[[3-(hydroxymethyl)anilino]methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

5-(2-Furyl)-3-[[2-methoxyanilino]methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

5-(2-Furyl)-3-[[4-(5-methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene]-1H-pyrrolo[2,3-b]pyridin-2-one;

5-(2-Furyl)-3-[[4-iodoanilino]methylidene]-1H-pyrrolo[2,3-b]pyridin-2-one;

3-({[2-Oxo-5-(3-thienyl)-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene]methyl}amino)benzamide;

Diethyl 4-({[2-oxo-5-(3-thienyl)-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene]methyl}amino)benzylphosphonate;

3-({[2-Oxo-5-(3-thienyl)-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene]methyl}amino)benzonitrile;

3-([2-(2-Hydroxyethyl)anilino]methylidene)-5-(3-thienyl)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

3-([3-(Hydroxymethyl)anilino]methylidene)-5-(3-thienyl)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

3-([2-Methoxyanilino]methylidene)-5-(3-thienyl)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

3-([4-(5-Methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene)-5-(3-thienyl)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

3-([4-Iodoanilino]methylidene)-5-(3-thienyl)-1H-pyrrolo[2,3-b]pyridin-2-one;

3-({[5-Bromo-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene]methyl}amino)benzamide;

Diethyl 4-({[5-Bromo-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene]methyl}amino)benzylphosphonate;

3-({[5-Bromo-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene]methyl}amino)benzonitrile;

5-Bromo-3-([2-(2-hydroxyethyl)anilino]methylidene)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

5-Bromo-3-([3-(hydroxymethyl)anilino]methylidene)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

5-Bromo-3-([2-methoxyanilino]methylidene)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

5-Bromo-3-([4-(5-methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

5-Bromo-3-([4-iodoanilino]methylidene)-1H-pyrrolo[2,3-b]pyridin-2-one;

3-[[[6-Chloro-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene)methyl]amino]benzamide;

Diethyl 4-[[[6-chloro-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene)methyl]amino]benzylphosphonate;

3-[[[6-Chloro-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene)methyl]amino]benzonitrile;

6-Chloro-3-[[2-(2-hydroxyethyl)anilino]methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

6-Chloro-3-[[3-(hydroxymethyl)anilino]methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

6-Chloro-3-[(2-methoxyanilino)methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

6-Chloro-3-[[4-(5-methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

6-Chloro-3-[(4-iodoanilino)methylidene]-1H-pyrrolo[2,3-b]pyridin-2-one;

Ethyl 3-[[3-(aminocarbonyl)anilino]methylidene]-2-oxo-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine-5-carboxylate;

Ethyl 3-[[4-[(diethoxyphosphoryl)methyl]anilino]methylidene]-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridine-5-carboxylate;

Ethyl 3-[(3-cyanoanilino)methylidene]-2-oxo-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine-5-carboxylate;

Ethyl 3-[[2-(2-hydroxyethyl)anilino]methylidene]-2-oxo-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine-5-carboxylate;

Ethyl 3-[[3-(hydroxymethyl)anilino]methylidene]-2-oxo-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine-5-carboxylate;

Ethyl 3-[(2-methoxyanilino)methylidene]-2-oxo-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine-5-carboxylate;

Ethyl 3-[[4-(5-methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino]methylidene]-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridine-5-carboxylate;

Ethyl 3-[(4-iodoanilino)methylidene]-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridine-5-carboxylate;

3-[(2-Oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene)methyl]amino}benzamide;

Diethyl 4-[(2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene)methyl]amino}benzylphosphonate;

3-[(2-Oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridin-3-ylidene)methyl]amino}benzonitrile;

3-[(2-(2-Hydroxyethyl)anilino)methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

3-[(3-(Hydroxymethyl)anilino)methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

3-[(2-Methoxyanilino)methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

3-[(4-(5-Methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino)methylidene]-1H-pyrrolo[2,3-b]pyridin-2-one;

3-[(4-Iodoanilino)methylidene]-1H-pyrrolo[2,3-b]pyridin-2-one;

3-[(2-Oxo-1,2-dihydro-3H-pyrrolo[3,2-b]pyridin-3-ylidene)methyl]amino}benzamide;

Diethyl 4-[(2-oxo-1,2-dihydro-3H-pyrrolo[3,2-b]pyridin-3-ylidene)methyl]amino}benzylphosphonate;

3-[(2-Oxo-1,2-dihydro-3H-pyrrolo[3,2-b]pyridin-3-ylidene)methyl]amino}benzonitrile;

3-[(2-(2-Hydroxyethyl)anilino)methylidene]-1,3-dihydro-2H-pyrrolo[3,2-b]pyridin-2-one;

3-[(3-(Hydroxymethyl)anilino)methylidene]-1,3-dihydro-2H-pyrrolo[3,2-b]pyridin-2-one;

3-[(2-Methoxyanilino)methylidene]-1,3-dihydro-2H-pyrrolo[3,2-b]pyridin-2-one;

3-[(4-(5-Methyl-3-oxo-2,3-dihydro-1H-pyrazol-1-yl)anilino)methylidene]-1H-pyrrolo[3,2-b]pyridin-2-one; and

3-[(4-Iodoanilino)methylidene]-1H-pyrrolo[3,2-b]pyridin-2-one;

and pharmaceutically active salts, solvates, hydrates, esters, amides, carbamates, affinity reagents and/or prodrugs thereof.

54. A compound as claimed in claim 1, in Z, E or Z and E form, selected from the group consisting of:

3-(((1-methyl-3-phenyl-1H-pyrazol-5-yl)amino)methylidene)-5-phenyl-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

3-(((4-Hydroxy-1-naphthyl)amino)methylidene)-5-phenyl-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

3-(((1-Ethyl-1H-pyrazol-5-yl)amino)methylidene)-5-phenyl-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

3-[(1H-Benzimidazol-2-ylamino)methylidene]-5-phenyl-1H-pyrrolo[2,3-b]pyridin-2-one;

3-(((6-(4-Morpholinyl)-3-pyridinyl)amino)methylidene)-5-phenyl-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

5-(2-Furyl)-3-(((1-methyl-3-phenyl-1H-pyrazol-5-yl)amino)methylidene)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

5-(2-Furyl)-3-(((4-hydroxy-1-naphthyl)amino)methylidene)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

3-(((1-Ethyl-1H-pyrazol-5-yl)amino)methylidene)-5-(2-furyl)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

3-[(1H-Benzimidazol-2-ylamino)methylidene]-5-(2-furyl)-1H-pyrrolo[2,3-b]pyridin-2-one;

5-(2-Furyl)-3-(((6-(4-morpholinyl)-3-pyridinyl)amino)methylidene)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

3-(((1-Methyl-3-phenyl-1H-pyrazol-5-yl)amino)methylidene)-5-(3-thienyl)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

3-(((4-Hydroxy-1-naphthyl)amino)methylidene)-5-(3-thienyl)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

3-(((1-Ethyl-1H-pyrazol-5-yl)amino)methylidene)-5-(3-thienyl)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

3-(((1H-Benzimidazol-2-ylamino)methylidene)-5-(3-thienyl)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

3-(((6-(4-Morpholinyl)-3-pyridinyl)amino)methylidene)-5-(3-thienyl)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

5-Bromo-3-(((1-methyl-3-phenyl-1H-pyrazol-5-yl)amino)methylidene)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

5-Bromo-3-(((4-hydroxy-1-naphthyl)amino)methylidene)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

5-Bromo-3-(((1-ethyl-1H-pyrazol-5-yl)amino)methylidene)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

3-(((1H-Benzimidazol-2-ylamino)methylidene)-5-bromo-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

5-Bromo-3-(((6-(4-morpholinyl)-3-pyridinyl)amino)methylidene)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

6-Chloro-3-(((1-methyl-3-phenyl-1H-pyrazol-5-yl)amino)methylidene)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

6-Chloro-3-(((4-hydroxy-1-naphthyl)amino)methylidene)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

6-Chloro-3-(((1-ethyl-1H-pyrazol-5-yl)amino)methylidene)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

3-(((1H-Benzimidazol-2-ylamino)methylidene)-6-chloro-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

6-Chloro-3-(((6-(4-morpholinyl)-3-pyridinyl)amino)methylidene)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

Ethyl 3-(((1-methyl-3-phenyl-1H-pyrazol-5-yl)amino)methylidene)-2-oxo-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine-5-carboxylate;

Ethyl 3-(((4-hydroxy-1-naphthyl)amino)methylidene)-2-oxo-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine-5-carboxylate;

Ethyl 3-(((1-ethyl-1H-pyrazol-5-yl)amino)methylidene)-2-oxo-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine-5-carboxylate;

Ethyl 3-[(1H-benzimidazol-2-ylamino)methylidene]-2-oxo-1,2-dihydro-3H-pyrrolo[2,3-b]pyridine-5-carboxylate;

Ethyl 3-([6-(4-morpholinyl)-3-pyridinyl]amino)methylidene)-2-oxo-2,3-dihydro-1H-pyrrolo[2,3-b]pyridine-5-carboxylate;

3-(((1-Methyl-3-phenyl-1H-pyrazol-5-yl)amino)methylidene)-6-phenyl-1,3-dihydro-2H-indol-2-one;

3-(((4-Hydroxy-1-naphthyl)amino)methylidene)-6-phenyl-1,3-dihydro-2H-indol-2-one;

3-(((1-Ethyl-1H-pyrazol-5-yl)amino)methylidene)-6-phenyl-1,3-dihydro-2H-indol-2-one;

3-[(1H-Benzimidazol-2-ylamino)methylidene]-6-phenyl-1H-indol-2-one;

3-([6-(4-Morpholinyl)-3-pyridinyl]amino)methylidene)-6-phenyl-1,3-dihydro-2H-indol-2-one;

6-(2-Furyl)-3-(((1-methyl-3-phenyl-1H-pyrazol-5-yl)amino)methylidene)-1,3-dihydro-2H-indol-2-one;

6-(2-Furyl)-3-([(4-hydroxy-1-naphthyl)amino)methylidene]-1,3-dihydro-2H-indol-2-one;

3-(((1-Ethyl-1H-pyrazol-5-yl)amino)methylidene)-6-(2-furyl)-1,3-dihydro-2H-indol-2-one;

3-[(1H-Benzimidazol-2-ylamino)methylidene]-6-(2-furyl)-1H-indol-2-one;

6-(2-Furyl)-3-([6-(4-morpholinyl)-3-pyridinyl]amino)methylidene)-1,3-dihydro-2H-indol-2-one;

3-(((1-Methyl-3-phenyl-1H-pyrazol-5-yl)amino)methylidene)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

3-([(4-Hydroxy-1-naphthyl)amino)methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

3-([(1-Ethyl-1H-pyrazol-5-yl)amino)methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

3-[(1H-Benzimidazol-2-ylamino)methylidene]-1H-pyrrolo[2,3-b]pyridin-2-one;

3-([(6-(4-Morpholinyl)-3-pyridinyl]amino)methylidene)-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one;

3-[(1-Methyl-3-phenyl-1H-pyrazol-5-yl)amino)methylidene]-1,3-dihydro-2H-pyrrolo[3,2-b]pyridin-2-one;

3-[(4-Hydroxy-1-naphthyl)amino)methylidene]-1,3-dihydro-2H-pyrrolo[3,2-b]pyridin-2-one;

3-[(1-Ethyl-1H-pyrazol-5-yl)amino)methylidene]-1,3-dihydro-2H-pyrrolo[3,2-b]pyridin-2-one;

3-[(1H-Benzimidazol-2-ylamino)methylidene]-1H-pyrrolo[3,2-b]pyridin-2-one;

3-([(6-(4-Morpholinyl)-3-pyridinyl]amino)methylidene)-1,3-dihydro-2H-pyrrolo[3,2-b]pyridin-2-one;

3-[(4-Hydroxy-1-naphthyl)amino)methylidene]-1,3-dihydro-2H-pyrrolo[2,3-b]pyridin-2-one; and

3-[(4-Hydroxy-1-naphthyl)amino)methylidene]-1,3-dihydro-2H-pyrrolo[3,2-b]pyridin-2-one;

and pharmaceutically active salts, solvates, hydrates, esters, amides, carbamates, affinity reagents and/or prodrugs thereof.

55. A pharmaceutical composition comprising a compound as claimed in any one of claims 1 to 54 and one or more pharmaceutically acceptable carriers, diluents and/or excipients.

5

56. A method of inhibiting a kinase comprising the step of binding said kinase with a compound as claimed in any one of claims 1 to 54.

57. The method of claim 56 wherein the kinase is selected from the group
 10 consisting of: abl, ARaf, ATK, ATM, bcr-abl, Blk, BRaf, Brk, Btk, CDK1, CDK2, CDK3, CDK4, CDK5, CDK6, CDK7, CDK8, CDK9, cfms, c-fms, c-kit, c-met, cRaf1, CSF1R, CSK, c-src, EGFR, ErbB2, ErbB3, ErbB4, ERK,

ERK1, ERK2, Fak, fes, FGFR1, FGFR2, FGFR3, FGFR4, FGFR5, Fgr, FLK-4, Fps, Frk, Fyn, GSK, gsk3a, gsk3b, Hck, IGF-1R, IKK, IKK1, IKK2, IKK3, INS-R, Integrin-linkedkinase, Jak, JAK1, JAK2, JAK3, JNK, JNK, Lck, Lyn, MEK, MEK1, MEK2, p38, PDGFR, PIK, PKB1, PKB2, PKB3, PKC, PKC α ,
5 PKC β , PKC δ , PKC ϵ , PKC γ , PKC λ , PKC μ , PKC ζ , PLK1, Polo-like kinase, PYK2, tie1, tie2, TrkA, TrkB, TrkC, UL13, UL97, VEGF-R1, VEGF-R2, Yes and Zap70.

58. The method of claim 56 wherein the kinase is selected from the group
10 consisting of: VEGF-R2, CDK1, CDK2, CDK4, Lck, Zap70, and c-Fms.

59. The method of claim 56 wherein the kinase is VEGF-R2.

60. A compound as claimed in any one of claims 1 to 54 for use in therapy.
15

61. A method of treating a disease or disorder in an animal which comprises administering to said animal a therapeutically effective amount of the compound as claimed in any one of claims 1 to 54.

20 62. The method according to claim 61 wherein the compound or physiologically acceptable salt thereof is administered in an amount of from about 0.1 to about 100 mg/kg of body weight per day.

63. The method according to claim 61 wherein the compound or
25 physiologically acceptable salt thereof is administered in an amount of from about 0.1 to about 10 mg/kg of body weight per day.

64. The method according to claim 61 wherein the animal is a human.

65. The method according to claim 61 wherein the disease or disorder comprises at least one condition selected from the group consisting of: organ transplant rejection, tumor growth, chemotherapy-induced mucositis, radiation-induced mucositis, plantar-palmar syndrome, chemotherapy-
5 induced alopecia, chemotherapy-induced thrombocytopenia, chemotherapy-induced leukopenia and hirsutism .
66. The method according to claim 61 wherein the disease or disorder comprises at least one condition selected from the group consisting of:
10 mucocitis, restenosis, atherosclerosis, rheumatoid arthritis, angiogenesis, hepatic cirrhosis, glomerulonephritis, diabetic nephropathy, malignant nephrosclerosis, chronic obstructive pulmonary disease, thrombotic microangiopathy, aglomerulopathy, psoriasis, diabetes mellitus, inflammation, a neurodegenerative disease, macular degeneration, actinic
15 keratosis and hyperproliferative disorders.
67. The method of claim 61 wherein the disease or disorder comprises at least one condition mediated by a kinase.
- 20 68. The method of claim 61 wherein the kinase is as defined in any one of claims 57 to 59.

INTERNATIONAL SEARCH REPORT

Int. Application No
PCT/US 00/05057

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D209/40 C07D487/06 C07D513/06 C07D405/04 C07D401/04
C07D405/14 C07D409/04 A61K31/40 A61K31/415 A61K31/425
A61K31/41 A61K31/47 A61P35/00 //(C07D487/06,209:00,

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,Y	WO 99 52869 A (BOEHRINGER INGELHEIM PHARMA KG) 21 October 1999 (1999-10-21) claims 1-11 ---	1-68
P,Y	WO 99 15500 A (GLAXO GROUP LTD.) 1 April 1999 (1999-04-01) claims 1-26 ---	1-68
E	WO 00 18734 A (BOEHRINGER INGELHEIM PHARMA KG) 6 April 2000 (2000-04-06) claims 1-11 ---	1-68
P,Y	DE 198 15 020 A (BOEHRINGER INGELHEIM PHARMA KG) 7 October 1999 (1999-10-07) claims 1-10 --- -/--	1-68

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

18 July 2000

Date of mailing of the international search report

04/09/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Herz, C

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/05057

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 235:00), (C07D487/06, 209:00, 253:00), (C07D513/06, 209:00, 277:00),
(C07D487/06, 209:00, 221:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	H. J. KALLMAYER: "Substituierte Isatin-phenylimine" ARCH. PHARM. (WEINHEIM, GER.), vol. 308, 1975, pages 742-748, XP000916680 * Compound 11a * table 1 -----	1

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

18 July 2000

Date of mailing of the international search report

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Herz, C

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. Jonal Application No

PCT/US 00/05057

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9952869 A	21-10-1999	DE 19816624 A AU 3814999 A	21-10-1999 01-11-1999
WO 9915500 A	01-04-1999	AU 9740798 A EP 1009738 A	12-04-1999 21-06-2000
WO 0018734 A	06-04-2000	DE 19844003 A AU 6086399 A	30-03-2000 17-04-2000
DE 19815020 A	07-10-1999	AU 3703499 A WO 9951590 A US 6043254 A	25-10-1999 14-10-1999 28-03-2000